



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 17, 2024 – 04:56 AM EST

PDB ID : 4OHS
Title : The structure of a far-red fluorescent protein, AQ143
Authors : Wannier, T.M.; Mayo, S.L.
Deposited on : 2014-01-17
Resolution : 2.19 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	238	 2% 86% 6% • 7%
1	B	238	 2% 86% 7% • 5%
1	C	238	 2% 78% 13% • 6%
1	D	238	 % 85% 7% 8%
1	E	238	 % 79% 12% • 8%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	238	<div><div></div><div>4%</div><div>84%</div><div>9%</div><div>• 5%</div></div>
1	G	238	<div><div></div><div>3%</div><div>82%</div><div>8%</div><div>• 9%</div></div>
1	H	238	<div><div></div><div>3%</div><div>74%</div><div>16%</div><div>• 8%</div></div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 14612 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FAR-RED FLUORESCENT PROTEIN AQ143.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	220	Total	C	N	O	S	0	5	0
			1789	1140	304	330	15			
1	B	226	Total	C	N	O	S	0	4	0
			1826	1164	310	337	15			
1	C	222	Total	C	N	O	S	1	4	0
			1789	1140	301	333	15			
1	D	218	Total	C	N	O	S	0	5	0
			1780	1135	302	328	15			
1	E	218	Total	C	N	O	S	0	3	0
			1765	1125	297	328	15			
1	F	226	Total	C	N	O	S	0	3	0
			1811	1154	307	335	15			
1	G	215	Total	C	N	O	S	0	3	0
			1734	1107	292	320	15			
1	H	217	Total	C	N	O	S	0	4	0
			1759	1122	297	325	15			

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		
2	B	1	Total	Cl	0	0
			1	1		
2	C	1	Total	Cl	0	0
			1	1		
2	D	1	Total	Cl	0	0
			1	1		



- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	63	Total 63	O 63	0	0
3	B	59	Total 59	O 59	0	0
3	C	44	Total 44	O 44	0	0
3	D	39	Total 39	O 39	0	0
3	E	46	Total 46	O 46	0	0
3	F	49	Total 49	O 49	0	0
3	G	26	Total 26	O 26	0	0
3	H	29	Total 29	O 29	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: FAR-RED FLUORESCENT PROTEIN AQ143





• Molecule 1: FAR-RED FLUORESCENT PROTEIN AQ143




• Molecule 1: FAR-RED FLUORESCENT PROTEIN AQ143




• Molecule 1: FAR-RED FLUORESCENT PROTEIN AQ143





• Molecule 1: FAR-RED FLUORESCENT PROTEIN AQ143









• Molecule 1: FAR-RED FLUORESCENT PROTEIN AQ143



• Molecule 1: FAR-RED FLUORESCENT PROTEIN AQ143



• Molecule 1: FAR-RED FLUORESCENT PROTEIN AQ143



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	50.97Å 68.11Å 132.79Å 98.38° 90.74° 110.47°	Depositor
Resolution (Å)	37.33 – 2.19 37.33 – 2.19	Depositor EDS
% Data completeness (in resolution range)	86.1 (37.33-2.19) 86.1 (37.33-2.19)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 2.20Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.190 , 0.221 0.193 , 0.222	Depositor DCC
R_{free} test set	3651 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	41.0	Xtriage
Anisotropy	0.780	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 47.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.029 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14612	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, NRQ, CH6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/1800	0.46	0/2428
1	B	0.35	0/1836	0.48	1/2477 (0.0%)
1	C	0.44	1/1801 (0.1%)	0.59	4/2431 (0.2%)
1	D	0.32	0/1791	0.46	0/2415
1	E	0.38	0/1769	0.48	1/2386 (0.0%)
1	F	0.39	0/1817	0.52	2/2453 (0.1%)
1	G	0.32	0/1737	0.47	1/2343 (0.0%)
1	H	0.40	1/1767 (0.1%)	0.51	2/2385 (0.1%)
All	All	0.37	2/14318 (0.0%)	0.50	11/19318 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	77	PRO	N-CD	5.28	1.55	1.47
1	C	225	PRO	N-CD	5.25	1.55	1.47

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	61	CYS	C-N-CA	8.18	142.16	121.70
1	B	61	CYS	C-N-CA	7.47	140.38	121.70
1	E	61	CYS	C-N-CA	6.04	136.81	121.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	22	HIS	C-N-CA	-6.04	106.59	121.70
1	C	224	ALA	C-N-CD	5.60	140.17	128.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	23[A]	HIS	Mainchain
1	C	23[B]	HIS	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1789	0	1723	9	0
1	B	1826	0	1756	18	0
1	C	1789	0	1717	21	0
1	D	1780	0	1713	9	0
1	E	1765	0	1693	14	0
1	F	1811	0	1734	27	0
1	G	1734	0	1661	11	0
1	H	1759	0	1683	40	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	63	0	0	0	0
3	B	59	0	0	0	0
3	C	44	0	0	2	0
3	D	39	0	0	2	0
3	E	46	0	0	0	0
3	F	49	0	0	1	0
3	G	26	0	0	1	0
3	H	29	0	0	1	0
All	All	14612	0	13680	146	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 146 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:203:LYS:HE3	1:B:211:GLU:OE1	1.40	1.19
1:H:11:ILE:HD11	1:H:41:GLU:HG3	1.21	1.16
1:H:203:LYS:HE2	1:H:205:GLU:HB2	1.33	1.10
1:F:203:LYS:HA	1:F:204:ALA:HB3	1.06	1.05
1:C:203:LYS:HD2	1:C:211:GLU:OE1	1.61	0.99

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	218/238 (92%)	215 (99%)	3 (1%)	0	100	100
1	B	223/238 (94%)	217 (97%)	6 (3%)	0	100	100
1	C	219/238 (92%)	212 (97%)	7 (3%)	0	100	100
1	D	216/238 (91%)	209 (97%)	7 (3%)	0	100	100
1	E	214/238 (90%)	209 (98%)	4 (2%)	1 (0%)	25	28
1	F	222/238 (93%)	216 (97%)	5 (2%)	1 (0%)	25	28
1	G	209/238 (88%)	208 (100%)	1 (0%)	0	100	100
1	H	214/238 (90%)	207 (97%)	6 (3%)	1 (0%)	25	28
All	All	1735/1904 (91%)	1693 (98%)	39 (2%)	3 (0%)	44	52

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	184	SER
1	F	206	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	203	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/198 (96%)	189 (100%)	0	100	100
1	B	193/198 (98%)	187 (97%)	6 (3%)	35	47
1	C	190/198 (96%)	184 (97%)	6 (3%)	34	45
1	D	189/198 (96%)	188 (100%)	1 (0%)	86	93
1	E	187/198 (94%)	183 (98%)	4 (2%)	48	63
1	F	190/198 (96%)	186 (98%)	4 (2%)	48	63
1	G	183/198 (92%)	181 (99%)	2 (1%)	70	82
1	H	186/198 (94%)	181 (97%)	5 (3%)	40	53
All	All	1507/1584 (95%)	1479 (98%)	28 (2%)	52	67

5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	143	SER
1	H	203	LYS
1	F	14	THR
1	H	73	VAL
1	E	209	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	39	GLN
1	E	157	GLN
1	F	206	GLN
1	G	197	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

16 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	NRQ	C	63[B]	-	24,24,25	3.26	4 (16%)	24,32,34	2.57	8 (33%)
1	CH6	H	63[A]	1	23,24,25	3.11	4 (17%)	28,32,34	2.71	11 (39%)
1	CH6	F	63[A]	1	23,24,25	3.10	4 (17%)	28,32,34	2.56	11 (39%)
1	CH6	G	63[A]	1	23,24,25	2.90	5 (21%)	28,32,34	2.79	10 (35%)
1	NRQ	G	63[B]	-	24,24,25	3.14	3 (12%)	24,32,34	2.90	8 (33%)
1	CH6	E	63[A]	1	23,24,25	3.06	5 (21%)	28,32,34	2.96	10 (35%)
1	NRQ	F	63[B]	-	24,24,25	3.19	3 (12%)	24,32,34	2.54	7 (29%)
1	CH6	C	63[A]	1	23,24,25	2.95	4 (17%)	28,32,34	2.54	13 (46%)
1	CH6	A	63[A]	1	23,24,25	2.95	4 (17%)	28,32,34	3.11	11 (39%)
1	NRQ	A	63[B]	-	24,24,25	3.19	3 (12%)	24,32,34	2.85	9 (37%)
1	CH6	B	63[A]	1	23,24,25	2.94	4 (17%)	28,32,34	2.74	9 (32%)
1	CH6	D	63[A]	1	23,24,25	3.10	5 (21%)	28,32,34	2.79	9 (32%)
1	NRQ	D	63[B]	-	24,24,25	3.24	3 (12%)	24,32,34	2.55	8 (33%)
1	NRQ	B	63[B]	-	24,24,25	3.30	4 (16%)	24,32,34	2.81	9 (37%)
1	NRQ	H	63[B]	-	24,24,25	3.16	3 (12%)	24,32,34	2.67	9 (37%)
1	NRQ	E	63[B]	-	24,24,25	3.30	3 (12%)	24,32,34	2.78	9 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	NRQ	C	63[B]	-	-	5/9/31/32	0/2/2/2
1	CH6	H	63[A]	1	-	5/12/31/32	0/2/2/2
1	CH6	F	63[A]	1	-	5/12/31/32	0/2/2/2
1	CH6	G	63[A]	1	-	6/12/31/32	0/2/2/2
1	NRQ	G	63[B]	-	-	5/9/31/32	0/2/2/2
1	CH6	E	63[A]	1	-	5/12/31/32	0/2/2/2
1	NRQ	F	63[B]	-	-	5/9/31/32	0/2/2/2
1	CH6	C	63[A]	1	-	5/12/31/32	0/2/2/2
1	CH6	A	63[A]	1	-	7/12/31/32	0/2/2/2
1	NRQ	A	63[B]	-	-	5/9/31/32	0/2/2/2
1	CH6	B	63[A]	1	-	4/12/31/32	0/2/2/2
1	CH6	D	63[A]	1	-	5/12/31/32	0/2/2/2
1	NRQ	D	63[B]	-	-	6/9/31/32	0/2/2/2
1	NRQ	B	63[B]	-	-	6/9/31/32	0/2/2/2
1	NRQ	H	63[B]	-	-	5/9/31/32	0/2/2/2
1	NRQ	E	63[B]	-	-	5/9/31/32	0/2/2/2

The worst 5 of 61 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	63[B]	NRQ	CB2-CA2	15.12	1.49	1.35
1	B	63[B]	NRQ	CB2-CA2	15.03	1.49	1.35
1	D	63[B]	NRQ	CB2-CA2	14.85	1.49	1.35
1	C	63[B]	NRQ	CB2-CA2	14.79	1.49	1.35
1	F	63[B]	NRQ	CB2-CA2	14.54	1.49	1.35

The worst 5 of 151 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	63[A]	CH6	CG2-CB2-CA2	-8.80	119.40	129.87
1	E	63[A]	CH6	CG2-CB2-CA2	-8.65	119.58	129.87
1	B	63[A]	CH6	CG2-CB2-CA2	-8.57	119.68	129.87
1	A	63[B]	NRQ	CG2-CB2-CA2	-8.23	120.08	129.87
1	G	63[B]	NRQ	CG2-CB2-CA2	-8.13	120.20	129.87

There are no chirality outliers.

5 of 84 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	63[A]	CH6	N2-CA2-CB2-CG2
1	A	63[A]	CH6	C2-CA2-CB2-CG2
1	A	63[B]	NRQ	N2-CA2-CB2-CG2
1	A	63[B]	NRQ	C2-CA2-CB2-CG2
1	B	63[A]	CH6	N2-CA2-CB2-CG2

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	H	63[A]	CH6	1	0
1	G	63[A]	CH6	3	0
1	C	63[A]	CH6	1	0
1	A	63[A]	CH6	1	0
1	B	63[A]	CH6	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	219/238 (92%)	0.13	4 (1%) 67 64	21, 49, 84, 127	4 (1%)
1	B	225/238 (94%)	0.17	4 (1%) 67 64	29, 51, 90, 122	3 (1%)
1	C	221/238 (92%)	0.26	5 (2%) 61 57	34, 57, 87, 115	3 (1%)
1	D	217/238 (91%)	0.15	2 (0%) 81 78	26, 56, 84, 96	4 (1%)
1	E	217/238 (91%)	0.17	2 (0%) 81 78	28, 58, 87, 125	2 (0%)
1	F	225/238 (94%)	0.45	9 (4%) 43 39	36, 66, 101, 132	2 (0%)
1	G	214/238 (89%)	0.56	7 (3%) 49 46	41, 74, 108, 127	2 (0%)
1	H	216/238 (90%)	0.57	7 (3%) 50 47	40, 72, 105, 131	3 (1%)
All	All	1754/1904 (92%)	0.31	40 (2%) 61 57	21, 59, 100, 132	23 (1%)

The worst 5 of 40 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	202	LEU	4.1
1	H	4	LEU	3.4
1	C	223	ALA	3.4
1	G	207	GLY	3.2
1	A	2	ALA	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

MODRES-RSR INFOmissingINFO

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CL	D	301	1/1	0.87	0.12	70,70,70,70	0
2	CL	B	301	1/1	0.89	0.11	74,74,74,74	0
2	CL	C	301	1/1	0.90	0.10	63,63,63,63	0
2	CL	A	301	1/1	0.92	0.09	60,60,60,60	0

6.5 Other polymers [i](#)

There are no such residues in this entry.