



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 16, 2025 – 12:09 PM EDT

PDB ID : 9OH6 / pdb\_00009oh6  
Title : H117A/M121H Azurin with Cu(II), pH 7.7  
Authors : Van Stappen, C.; Lu, Y.  
Deposited on : 2025-05-02  
Resolution : 2.04 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.006 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.44


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*





The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2096 (2.04-2.04)
Clashscore	180529	2229 (2.04-2.04)
Ramachandran outliers	177936	2217 (2.04-2.04)
Sidechain outliers	177891	2217 (2.04-2.04)
RSRZ outliers	164620	2096 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	128	
1	E	128	
1	H	128	
1	K	128	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4478 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Azurin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	128	Total	C	N	O	S	1	0	0
			971	605	164	194	8			
1	E	128	Total	C	N	O	S	1	0	0
			971	605	164	194	8			
1	H	128	Total	C	N	O	S	0	0	0
			971	605	164	194	8			
1	K	128	Total	C	N	O	S	1	0	0
			971	605	164	194	8			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	117	ALA	HIS	engineered mutation	UNP P00282
B	121	HIS	MET	engineered mutation	UNP P00282
E	117	ALA	HIS	engineered mutation	UNP P00282
E	121	HIS	MET	engineered mutation	UNP P00282
H	117	ALA	HIS	engineered mutation	UNP P00282
H	121	HIS	MET	engineered mutation	UNP P00282
K	117	ALA	HIS	engineered mutation	UNP P00282
K	121	HIS	MET	engineered mutation	UNP P00282

- Molecule 2 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C<sub>4</sub>H<sub>12</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			8	4	1	3		
2	E	1	Total	C	N	O	0	0
			8	4	1	3		
2	H	1	Total	C	N	O	0	0
			8	4	1	3		
2	K	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 3 is COPPER (II) ION (CCD ID: CU) (formula: Cu) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Cu	0	0
			2	2		
3	E	2	Total	Cu	0	0
			2	2		
3	H	2	Total	Cu	0	0
			2	2		
3	K	2	Total	Cu	0	0
			2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	149	Total	O	0	0
			149	149		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	142	Total 142	O 142	0	0
4	H	129	Total 129	O 129	0	0
4	K	134	Total 134	O 134	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Azurin

Chain B:  91% 9%



- Molecule 1: Azurin

Chain E:  88% 12%





- Molecule 1: Azurin

Chain H:  91% 8% •



- Molecule 1: Azurin

Chain K:  84% 16%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.91Å 48.84Å 81.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	81.31 – 2.04 81.31 – 2.04	Depositor EDS
% Data completeness (in resolution range)	99.0 (81.31-2.04) 99.2 (81.31-2.04)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.06 (at 2.03Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, $R_{free}$	0.176 , 0.233 0.177 , 0.233	Depositor DCC
$R_{free}$ test set	1475 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.5	Xtriage
Anisotropy	0.049	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 36.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.447 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4478	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.66 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.0706e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CU, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	B	0.30	0/988	0.47	0/1332
1	E	0.30	0/988	0.47	0/1332
1	H	0.30	0/988	0.52	0/1332
1	K	0.29	0/988	0.48	0/1332
All	All	0.30	0/3952	0.49	0/5328

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	971	0	948	8	1
1	E	971	0	948	10	1
1	H	971	0	948	6	1
1	K	971	0	948	13	1
2	B	8	0	11	0	1
2	E	8	0	11	0	1
2	H	8	0	11	0	1
2	K	8	0	11	1	1
3	B	2	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	2	0	0	0	0
3	H	2	0	0	0	0
3	K	2	0	0	0	0
4	B	149	0	0	5	3
4	E	142	0	0	1	2
4	H	129	0	0	2	0
4	K	134	0	0	3	3
All	All	4478	0	3836	37	9

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:13:MET:O	4:B:301:HOH:O	2.02	0.77
1:B:13:MET:HG2	4:B:419:HOH:O	1.88	0.73
1:B:98:ASP:OD1	4:B:304:HOH:O	2.11	0.69
1:H:76:ASP:OD2	4:H:301:HOH:O	2.13	0.66
1:H:69:ASP:OD1	4:H:302:HOH:O	2.15	0.64
1:E:12:GLN:NE2	4:E:302:HOH:O	2.30	0.64
1:B:117:ALA:O	4:B:305:HOH:O	2.16	0.62
1:E:21:THR:HB	1:E:128:LYS:HD3	1.83	0.60
1:K:55:ASP:OD1	4:K:301:HOH:O	2.17	0.59
1:B:21:THR:HB	1:B:128:LYS:HD3	1.85	0.56
1:K:33:LEU:O	1:K:92:LYS:HD2	2.06	0.55
1:H:40:PRO:HD2	1:H:43:VAL:CG1	2.37	0.54
1:B:11:ASP:O	1:B:44:MET:HE3	2.09	0.52
1:K:24:LYS:HE3	1:K:128:LYS:C	2.35	0.52
1:K:1:ALA:O	2:K:201:TRS:H21	2.13	0.49
1:K:64:MET:HG2	1:K:113:THR:O	2.13	0.48
1:K:92:LYS:NZ	1:K:94:SER:OG	2.33	0.48
1:E:48:TRP:O	1:E:83:HIS:HA	2.15	0.47
1:K:31:VAL:HG21	1:K:48:TRP:CZ2	2.50	0.47
1:K:9:GLY:O	1:K:36:PRO:HD2	2.15	0.46
1:B:121:HIS:HB2	4:B:301:HOH:O	2.15	0.45
1:E:7:ILE:HD12	1:E:17:THR:HG21	1.98	0.45
1:H:64:MET:HG2	1:H:113:THR:O	2.16	0.44
1:K:13:MET:HG3	4:K:352:HOH:O	2.18	0.44
1:E:6:ASP:O	1:E:7:ILE:HD13	2.18	0.44
1:K:11:ASP:HA	1:K:44:MET:HG2	2.00	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:24:LYS:HG3	1:E:128:LYS:O	2.18	0.43
1:E:6:ASP:C	1:E:7:ILE:HD13	2.44	0.43
1:H:70:LYS:HD3	1:H:75:PRO:HG3	2.01	0.43
1:E:4:SER:HA	1:E:30:THR:O	2.19	0.42
1:K:49:VAL:O	1:K:110:PHE:HA	2.19	0.42
1:B:119:ALA:O	1:B:122:LYS:HE2	2.19	0.42
1:K:21:THR:HG22	4:K:306:HOH:O	2.20	0.41
1:K:9:GLY:HA3	1:K:46:HIS:CD2	2.56	0.41
1:E:64:MET:HE2	1:E:115:PRO:HA	2.02	0.41
1:H:7:ILE:HD11	1:H:20:ILE:HD11	2.03	0.41
1:E:8:GLN:HG2	1:E:34:SER:OG	2.20	0.40

All (9) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:83:HIS:NE2	2:K:201:TRS:O3[2_554]	1.30	0.90
1:B:83:HIS:NE2	2:B:201:TRS:O3[2_545]	1.31	0.89
1:E:83:HIS:NE2	2:E:201:TRS:O3[2_454]	1.31	0.89
1:H:83:HIS:NE2	2:H:201:TRS:O3[2_445]	1.31	0.89
4:B:337:HOH:O	4:E:302:HOH:O[1_655]	1.77	0.43
4:B:323:HOH:O	4:B:363:HOH:O[2_545]	1.90	0.30
4:E:365:HOH:O	4:K:369:HOH:O[1_465]	2.07	0.13
4:K:366:HOH:O	4:K:407:HOH:O[2_544]	2.15	0.05
4:B:388:HOH:O	4:K:364:HOH:O[1_565]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	126/128 (98%)	124 (98%)	2 (2%)	0	<b>100</b> <b>100</b>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	126/128 (98%)	125 (99%)	1 (1%)	0	100	100
1	H	126/128 (98%)	125 (99%)	1 (1%)	0	100	100
1	K	126/128 (98%)	125 (99%)	1 (1%)	0	100	100
All	All	504/512 (98%)	499 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	109/109 (100%)	109 (100%)	0	100	100
1	E	109/109 (100%)	108 (99%)	1 (1%)	75	77
1	H	109/109 (100%)	108 (99%)	1 (1%)	75	77
1	K	109/109 (100%)	109 (100%)	0	100	100
All	All	436/436 (100%)	434 (100%)	2 (0%)	86	88

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	104	GLU
1	H	43	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	32	ASN
1	E	12	GLN
1	E	32	ASN
1	H	107	GLN
1	K	12	GLN
1	K	28	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	K	32	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	TRS	K	201	3	7,7,7	0.24	0	9,9,9	0.85	0
2	TRS	B	201	3	7,7,7	0.35	0	9,9,9	1.53	2 (22%)
2	TRS	E	201	3	7,7,7	0.35	0	9,9,9	0.82	0
2	TRS	H	201	3	7,7,7	0.28	0	9,9,9	0.93	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRS	K	201	3	-	6/9/9/9	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRS	B	201	3	-	3/9/9/9	-
2	TRS	E	201	3	-	9/9/9/9	-
2	TRS	H	201	3	-	5/9/9/9	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	201	TRS	C3-C-C1	-3.11	102.37	110.66
2	B	201	TRS	C3-C-N	2.07	113.45	108.17

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	201	TRS	N-C-C1-O1
2	H	201	TRS	C2-C-C3-O3
2	K	201	TRS	C2-C-C3-O3
2	B	201	TRS	C1-C-C3-O3
2	E	201	TRS	C2-C-C1-O1
2	E	201	TRS	C3-C-C2-O2
2	H	201	TRS	C1-C-C3-O3
2	K	201	TRS	C1-C-C2-O2
2	K	201	TRS	C1-C-C3-O3
2	B	201	TRS	C2-C-C3-O3
2	E	201	TRS	N-C-C2-O2
2	E	201	TRS	C2-C-C3-O3
2	E	201	TRS	N-C-C3-O3
2	K	201	TRS	N-C-C2-O2
2	E	201	TRS	C1-C-C2-O2
2	E	201	TRS	C1-C-C3-O3
2	K	201	TRS	C3-C-C2-O2
2	B	201	TRS	N-C-C3-O3
2	E	201	TRS	C3-C-C1-O1
2	H	201	TRS	N-C-C3-O3
2	K	201	TRS	N-C-C3-O3
2	H	201	TRS	C2-C-C1-O1
2	H	201	TRS	C1-C-C2-O2

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	K	201	TRS	1	1
2	B	201	TRS	0	1
2	E	201	TRS	0	1
2	H	201	TRS	0	1

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	128/128 (100%)	-1.43	0 100 100	13, 19, 34, 44	1 (0%)
1	E	128/128 (100%)	-1.43	0 100 100	12, 20, 34, 42	1 (0%)
1	H	128/128 (100%)	-1.44	0 100 100	13, 20, 28, 35	0
1	K	128/128 (100%)	-1.45	0 100 100	13, 20, 29, 36	1 (0%)
All	All	512/512 (100%)	-1.44	0 100 100	12, 20, 32, 44	3 (0%)

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	TRS	B	201	8/8	0.98	0.06	20,22,27,28	0
2	TRS	E	201	8/8	0.99	0.04	19,22,26,27	0
2	TRS	H	201	8/8	0.99	0.03	17,23,28,30	0
2	TRS	K	201	8/8	0.99	0.04	16,22,25,26	0

*Continued on next page...*

*Continued from previous page...*




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CU	B	202	1/1	1.00	0.03	24,24,24,24	1
3	CU	B	203	1/1	1.00	0.01	22,22,22,22	1
3	CU	E	202	1/1	1.00	0.01	24,24,24,24	1
3	CU	E	203	1/1	1.00	0.01	19,19,19,19	1
3	CU	H	202	1/1	1.00	0.03	24,24,24,24	1
3	CU	H	203	1/1	1.00	0.01	15,15,15,15	1
3	CU	K	202	1/1	1.00	0.01	25,25,25,25	1
3	CU	K	203	1/1	1.00	0.01	17,17,17,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around CU B 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around CU B 203:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around CU E 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around CU E 203:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around CU H 202:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around CU H 203:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around CU K 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around CU K 203:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers ⓘ

There are no such residues in this entry.