



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 13, 2025 – 10:11 AM EST

PDB ID : 9OHZ / pdb\_00009ohz  
Title : Crystal structure of CD1a presenting ganglioside GD3  
Authors : Cao, T.P.; Ciacchi, L.; Rossjohn, J.; Shahine, A.  
Deposited on : 2025-05-05  
Resolution : 2.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.46


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	3585 (2.16-2.12)
Ramachandran outliers	177936	3554 (2.16-2.12)
Sidechain outliers	177891	3553 (2.16-2.12)
RSRZ outliers	164620	3337 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	285	
2	B	106	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3208 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T-cell surface glycoprotein CD1a.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	270	Total	C	N	O	S	0	2	0
			2162	1387	374	393	8			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	ALA	-	expression tag	UNP P06126
A	2	THR	-	expression tag	UNP P06126
A	51	TRP	CYS	variant	UNP P06126
A	279	GLY	-	expression tag	UNP P06126
A	280	SER	-	expression tag	UNP P06126
A	281	LEU	-	expression tag	UNP P06126
A	282	VAL	-	expression tag	UNP P06126
A	283	PRO	-	expression tag	UNP P06126
A	284	ARG	-	expression tag	UNP P06126
A	285	GLY	-	expression tag	UNP P06126

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	105	Total	C	N	O	S	0	0	0
			857	546	144	164	3			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	100	GLY	-	expression tag	UNP P61769
B	101	SER	-	expression tag	UNP P61769
B	102	LEU	-	expression tag	UNP P61769
B	103	VAL	-	expression tag	UNP P61769
B	104	PRO	-	expression tag	UNP P61769
B	105	ARG	-	expression tag	UNP P61769

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is N-{(2S,3R)-1-[(4-O-beta-D-galactopyranosyl-beta-D-glucopyranosyl)oxy]-3-hydroxyoctadecan-2-yl}tetracosanamide (CCD ID: A1CB9) (formula:  $C_{54}H_{105}NO_{13}$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			68	54	1	13		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	55	Total	O	0	0
			55	55		
5	B	52	Total	O	0	0
			52	52		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: T-cell surface glycoprotein CD1a



- Molecule 2: Beta-2-microglobulin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	42.39Å 90.35Å 107.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.18 – 2.14 46.18 – 2.14	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.18-2.14) 99.9 (46.18-2.14)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.12 (at 2.14Å)	Xtriage
Refinement program	PHENIX (1.21.2_5419: ???)	Depositor
R, $R_{free}$	0.217 , 0.247 0.218 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	42.2	Xtriage
Anisotropy	0.282	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 42.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3208	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.52% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, A1CB9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.29	0/2237	0.74	2/3048 (0.1%)
2	B	0.23	0/881	0.64	0/1197
All	All	0.27	0/3118	0.71	2/4245 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	82	ARG	CA-CB-CG	5.42	124.94	114.10
1	A	82	ARG	CB-CG-CD	-5.03	99.74	111.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2162	0	2020	37	0
2	B	857	0	812	6	0
3	A	14	0	13	0	0
4	A	68	0	0	0	0
5	A	55	0	0	1	0
5	B	52	0	0	2	0
All	All	3208	0	2845	42	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:278:GLU:HA	1:A:282:VAL:HG23	1.67	0.75
1:A:14:TRP:HD1	1:A:28:VAL:HG22	1.56	0.70
1:A:278:GLU:HG3	1:A:279:GLY:N	2.11	0.65
2:B:102:LEU:HD22	2:B:102:LEU:H	1.63	0.63
1:A:69:LEU:HD21	1:A:161:LEU:HD11	1.81	0.62
1:A:20:ASN:H	1:A:24:LYS:HE3	1.63	0.62
1:A:68:THR:O	1:A:72:ILE:HG12	2.00	0.62
1:A:278:GLU:HG3	1:A:279:GLY:H	1.64	0.61
1:A:187:PRO:HB3	1:A:211:PHE:HB3	1.82	0.61
1:A:278:GLU:CD	1:A:284:ARG:HE	2.08	0.60
1:A:278:GLU:OE1	1:A:284:ARG:NH2	2.30	0.60
1:A:205:VAL:HG22	1:A:248:THR:HG22	1.84	0.59
1:A:82:ARG:CZ	1:A:82:ARG:HB3	2.33	0.59
1:A:278:GLU:OE2	1:A:284:ARG:HG3	2.05	0.56
1:A:217:TRP:HB3	1:A:264:LYS:HB2	1.90	0.54
1:A:86:HIS:HB2	5:A:404:HOH:O	2.08	0.53
1:A:73:ARG:HH21	1:A:154:GLU:HG3	1.75	0.52
1:A:256:ALA:HA	1:A:259:LEU:HD12	1.90	0.52
1:A:278:GLU:CG	1:A:279:GLY:H	2.23	0.51
1:A:49:PHE:HB3	1:A:54:SER:HB2	1.94	0.49
1:A:167:PRO:O	1:A:171:LEU:HG	2.13	0.49
2:B:1:ILE:O	5:B:201:HOH:O	2.21	0.46
2:B:17:ASN:HD21	2:B:74:GLU:HB3	1.81	0.46
1:A:64:LYS:O	1:A:68:THR:HG23	2.15	0.46
1:A:246:ARG:HD2	5:B:236:HOH:O	2.16	0.45
1:A:82:ARG:HB3	1:A:82:ARG:NH1	2.32	0.45
1:A:43:ASN:OD1	1:A:44:SER:N	2.50	0.44
1:A:47:ILE:HG13	1:A:67:GLU:HG2	1.99	0.44
1:A:221:MET:HE3	1:A:221:MET:HB3	1.92	0.44
1:A:278:GLU:CG	1:A:279:GLY:N	2.78	0.44
1:A:88:LEU:O	1:A:140:MET:HE3	2.18	0.43
1:A:19:TYR:HB2	1:A:23:TRP:O	2.18	0.43
1:A:263:VAL:HB	1:A:273:ILE:HB	2.00	0.43
1:A:23:TRP:O	1:A:24:LYS:HD2	2.18	0.43
1:A:39:THR:HB	1:A:48:VAL:HG13	2.02	0.42
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.55	0.42
2:B:51:HIS:HA	2:B:65:LEU:O	2.19	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:14:TRP:CD1	1:A:28:VAL:HG22	2.46	0.41
1:A:277:TRP:O	1:A:278:GLU:HB2	2.20	0.41
2:B:91:LYS:HE3	2:B:91:LYS:HB2	1.97	0.41
1:A:20:ASN:N	1:A:24:LYS:HE3	2.33	0.40
1:A:10:PHE:H	1:A:170:ILE:HD12	1.86	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	266/285 (93%)	248 (93%)	12 (4%)	6 (2%)	5	1
2	B	103/106 (97%)	101 (98%)	2 (2%)	0	100	100
All	All	369/391 (94%)	349 (95%)	14 (4%)	6 (2%)	8	3

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	258	ASP
1	A	110	SER
1	A	43	ASN
1	A	128	ASN
1	A	257	ALA
1	A	278	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/246 (93%)	222 (97%)	7 (3%)	35	34
2	B	96/99 (97%)	95 (99%)	1 (1%)	73	77
All	All	325/345 (94%)	317 (98%)	8 (2%)	42	43

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	LYS
1	A	25	GLN
1	A	48	VAL
1	A	75	ILE
1	A	86	HIS
1	A	246	ARG
1	A	250	GLU
2	B	102	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	115	GLN
1	A	129	ASN
2	B	2	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	A1CB9	A	302	-	69,69,69	1.60	12 (17%)	80,83,83	2.99	7 (8%)
3	NAG	A	301	1	14,14,15	0.40	0	17,19,21	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	A1CB9	A	302	-	-	36/59/99/99	0/2/2/2
3	NAG	A	301	1	-	4/6/23/26	0/1/1/1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	302	A1CB9	C83-N82	6.57	1.48	1.34
4	A	302	A1CB9	C14-C15	-3.44	1.37	1.52
4	A	302	A1CB9	C15-C16	2.98	1.57	1.52
4	A	302	A1CB9	C99-C98	-2.92	1.37	1.51
4	A	302	A1CB9	O38-C31	2.84	1.50	1.43
4	A	302	A1CB9	O34-C33	2.40	1.50	1.44
4	A	302	A1CB9	O84-C83	-2.37	1.18	1.23
4	A	302	A1CB9	C32-C31	-2.33	1.46	1.52
4	A	302	A1CB9	C85-C83	2.26	1.55	1.51
4	A	302	A1CB9	OA8-C16	-2.24	1.38	1.43
4	A	302	A1CB9	C22-C23	-2.15	1.46	1.52
4	A	302	A1CB9	C18-C17	2.02	1.55	1.51

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	302	A1CB9	C14-C15-C16	24.43	154.14	114.11
4	A	302	A1CB9	O25-C24-C23	5.29	120.66	109.72
4	A	302	A1CB9	C26-C24-C23	-5.15	98.88	113.38
4	A	302	A1CB9	C29-O28-C23	-3.01	110.85	117.98
4	A	302	A1CB9	O34-C33-C32	2.61	114.40	109.70
4	A	302	A1CB9	C85-C83-N82	2.37	120.05	115.86
4	A	302	A1CB9	C35-C33-C32	-2.24	107.53	113.02

There are no chirality outliers.

All (40) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	302	A1CB9	C15-C16-C17-C18
4	A	302	A1CB9	C15-C16-C17-N82
4	A	302	A1CB9	OA8-C16-C17-C18
4	A	302	A1CB9	OA8-C16-C17-N82
4	A	302	A1CB9	C16-C17-C18-O19
4	A	302	A1CB9	N82-C17-C18-O19
4	A	302	A1CB9	C97-C98-C99-CA0
3	A	301	NAG	O5-C5-C6-O6
4	A	302	A1CB9	O25-C24-C26-O27
4	A	302	A1CB9	C03-C04-C05-C06
4	A	302	A1CB9	C85-C86-C87-C88
4	A	302	A1CB9	C23-C24-C26-O27
3	A	301	NAG	C8-C7-N2-C2
3	A	301	NAG	O7-C7-N2-C2
4	A	302	A1CB9	C11-C12-C13-C14
4	A	302	A1CB9	CA0-CA1-CA2-CA3
4	A	302	A1CB9	C09-C10-C11-C12
3	A	301	NAG	C4-C5-C6-O6
4	A	302	A1CB9	C05-C06-C07-C08
4	A	302	A1CB9	C12-C13-C14-C15
4	A	302	A1CB9	C92-C93-C94-C95
4	A	302	A1CB9	C90-C91-C92-C93
4	A	302	A1CB9	C91-C92-C93-C94
4	A	302	A1CB9	C08-C09-C10-C11
4	A	302	A1CB9	CA3-CA4-CA5-CA6
4	A	302	A1CB9	C88-C89-C90-C91
4	A	302	A1CB9	C86-C87-C88-C89
4	A	302	A1CB9	C87-C88-C89-C90
4	A	302	A1CB9	C93-C94-C95-C96
4	A	302	A1CB9	C01-C02-C03-C04
4	A	302	A1CB9	CA2-CA3-CA4-CA5

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
4	A	302	A1CB9	C96-C97-C98-C99
4	A	302	A1CB9	C21-C20-O19-C18
4	A	302	A1CB9	C95-C96-C97-C98
4	A	302	A1CB9	CA4-CA5-CA6-CA7
4	A	302	A1CB9	C94-C95-C96-C97
4	A	302	A1CB9	O25-C20-O19-C18
4	A	302	A1CB9	C02-C03-C04-C05
4	A	302	A1CB9	C16-C17-N82-C83
4	A	302	A1CB9	C10-C11-C12-C13

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	270/285 (94%)	0.98	41 (15%) <b>6</b> <b>8</b>	18, 54, 86, 101	3 (1%)
2	B	105/106 (99%)	0.62	8 (7%) <b>21</b> <b>26</b>	30, 42, 76, 98	0
All	All	375/391 (95%)	0.88	49 (13%) <b>8</b> <b>11</b>	18, 51, 86, 101	3 (0%)

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	102	LEU	4.5
1	A	257	ALA	4.3
2	B	0	ALA	4.2
2	B	100	GLY	4.2
1	A	23	TRP	4.1
1	A	75	ILE	3.9
2	B	103	VAL	3.7
1	A	110	SER	3.5
2	B	1	ILE	3.5
1	A	109	VAL	3.5
1	A	66	LEU	3.3
2	B	101	SER	3.2
2	B	104	PRO	3.2
1	A	278	GLU	3.1
1	A	70	PHE	2.9
1	A	63	TRP	2.9
1	A	171	LEU	2.9
2	B	99	MET	2.8
1	A	40	TRP	2.8
1	A	58	PHE	2.7
1	A	78	PHE	2.7
1	A	139	ASN	2.7
1	A	72	ILE	2.7
1	A	181	LEU	2.6

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	47	ILE	2.6
1	A	254	GLY	2.5
1	A	56	GLY	2.5
1	A	61	GLU	2.4
1	A	84	TYR	2.4
1	A	111	GLY	2.4
1	A	49	PHE	2.3
1	A	282	VAL	2.3
1	A	9	SER	2.3
1	A	147	VAL	2.2
1	A	172	GLY	2.2
1	A	83	ARG	2.1
1	A	33	SER	2.1
1	A	86	HIS	2.1
1	A	19	TYR	2.1
1	A	279	GLY	2.1
1	A	57	ASN	2.1
1	A	92	TYR	2.1
1	A	45	SER	2.1
1	A	69	LEU	2.0
1	A	88	LEU	2.0
1	A	132	LEU	2.0
1	A	80	GLY	2.0
1	A	82	ARG	2.0
1	A	53	TRP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	A	301	14/15	0.29	0.21	72,87,93,94	0
4	A1CB9	A	302	68/68	0.76	0.18	53,73,109,118	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.