



wwPDB EM Validation Summary Report ⓘ

Jul 8, 2024 – 09:02 am BST

PDB ID : 7ORI
EMDB ID : EMD-13038
Title : La Crosse virus polymerase at replication late-elongation stage
Authors : Arragain, B.; Durieux Trouilleteau, Q.; Baudin, F.; Cusack, S.; Schoehn, G.;
Malet, H.
Deposited on : 2021-06-06
Resolution : 3.90 Å(reported)
Based on initial model : 6Z8K

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92
Mogul : 1.8.4, CSD as541be (2020)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	17	
2	T	30	
3	P	30	
4	A	2276	

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 17326 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (5'-R(P*AP*CP*GP*AP*GP*UP*GP*UP*CP*GP*U)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
1	H	11	Total	C	N	O	P	0	0
			237	105	42	79	11		

- Molecule 2 is a RNA chain called RNA (5'-R(P*AP*AP*CP*GP*UP*UP*AP*UP*CP*UP*AP*UP*AP*AP*CP*AP*CP*UP*AP*CP*U)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
2	T	21	Total	C	N	O	P	0	0
			439	198	74	146	21		

- Molecule 3 is a RNA chain called RNA (5'-R(P*AP*UP*AP*GP*AP*UP*AP*AP*CP*GP*UP*U)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
3	P	12	Total	C	N	O	P	0	0
			256	115	46	83	12		

- Molecule 4 is a protein called La Crosse virus polymerase.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	A	2007	Total	C	N	O	S	0	0
			16383	10499	2717	3059	108		

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	34	LYS	HIS	engineered mutation	UNP A5HC98
A	1028G	SER	-	insertion	UNP A5HC98
A	1028H	GLY	-	insertion	UNP A5HC98
A	1028I	TRP	-	insertion	UNP A5HC98
A	1028J	SER	-	insertion	UNP A5HC98
A	1028K	HIS	-	insertion	UNP A5HC98

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	1028L	PRO	-	insertion	UNP A5HC98
A	1028M	GLN	-	insertion	UNP A5HC98
A	1028N	PHE	-	insertion	UNP A5HC98
A	1028O	GLU	-	insertion	UNP A5HC98
A	1028P	LYS	-	insertion	UNP A5HC98
A	1028Q	GLY	-	insertion	UNP A5HC98
A	1028R	SER	-	insertion	UNP A5HC98
A	1028S	GLY	-	insertion	UNP A5HC98


- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	AltConf
5	A	1	Total Zn 1 1	0

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	AltConf
6	A	1	Total Mg 1 1	0

- Molecule 7 is PYROPHOSPHATE 2- (three-letter code: POP) (formula: H₂O₇P₂).





Mol	Chain	Residues	Atoms	AltConf
7	A	1	Total O P 9 7 2	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RNA (5'-R(P*AP*CP*GP*AP*GP*UP*GP*UP*CP*GP*U)-3')

Chain H: 




- Molecule 2: RNA (5'-R(P*AP*AP*CP*GP*UP*UP*AP*UP*CP*UP*AP*UP*AP*AP*CP*A P*CP*UP*AP*CP*U)-3')

Chain T: 




- Molecule 3: RNA (5'-R(P*AP*UP*AP*GP*AP*UP*AP*AP*CP*GP*UP*U)-3')

Chain P: 



- Molecule 4: La Crosse virus polymerase

Chain A: 



R2205	K643	L1144	L1298	R1424	P1599	I1730	LYS	ASP	Y2100	R2205
T2216	R653	Q145	I1298	R1439	P1599	K1731	ILE	MET	S2101	T2216
P2217	Y654	G146	P1317	K1440	R1602	Y1736	GLN	GLY	R2102	P2217
P2218	M655	N147	P1317	L1441	M1605	Q1743	THR	ASN	I1982	P2218
N2225	I656	F148	A1323	E1466	M1605	L1744	PRO	GLY	L2116	N2225
F2226	D688	SER	R1326	K1467	C1607	L1745	ASN	THR	L2117	F2226
K2227	Y669	TRP	K1327	K1467	S1608	L1745	GLY	THR	L2118	K2227
K2228	K797	PRO	D1328	L1498	S1608	S1759	THR	THR	E2119	K2228
E2121	I798	PHE	K1327	L1498	S1608	E1760	ILE	GLY	R2120	E2121
T2122	L799	GLN	D1328	A1504	P1611	E1760	THR	THR	R1999	T2122
E2126	K805	LYS	K1340	A1504	P1611	A1764	TYR	TYR	I2000	E2126
E2127	R813	GLY	L1341	F1507	K1617	A1764	ASN	ASN	T2009	E2127
A2128	L817	SER	S1342	I1508	L1618	C1767	ARG	ARG	T2010	A2128
F2129	L821	GLY	M1343	I1511	H1621	F1768	LYS	SER	K2011	F2129
T2130	F695	TYR	T1346	T1511	L1625	L1769	ASP	THR	K2012	T2130
N2134	S842	SER	V1347	K1516	L1625	L1770	ILE	ARG	G2023	N2134
G2135	C943	ASN	G1348	F1517	V1626	L1771	THR	THR	P2024	G2135
F2136	L844	LEU	L1349	V1517	L1627	T1772	GLY	GLN	P2024	F2136
N2141	Y710	GLY	L1354	V1518	D1641	H1773	GLU	VAL	L2030	N2141
G2145	L711	LYS	L1357	Y1528	P1642	L1790	ASN	THR	L2030	G2145
V2148	I716	ILE	L1357	I1529	T1643	I1791	LYS	GLN	T2034	V2148
V2149	Q718	GLY	I1358	N1530	E1644	I1791	ILE	ILE	E2035	V2149
L2152	K720	VAL	K1358	LEU	M1645	E1794	THR	HIS	K2038	L2152
N2159	I721	VAL	L1360	HIS	A1646	F1795	TYR	SER	S2039	N2159
W2160	L724	LYS	T1365	ASP	R1647	S1796	ALA	GLY	V2148	W2160
K2166	N724	GLN	T1365	ARG	H1651	Y1797	LEU	LEU	V2149	K2166
H2169	I730	LYS	R1371	ALA	H1651	L1805	CYS	ILE	N2046	H2169
L2172	W730	ILE	E1372	LEU	K1666	Y1813	THR	THR	N2049	L2172
M2177	W731	LEU	S1373	GLY	V1667	Y1813	THR	THR	F2050	M2177
H2179	F732	LEU	E1373	LYS	R1668	P1820	LYS	ARG	R2051	H2179
Y2180	V736	VAL	Q1377	PRO	M1671	P1820	THR	GLY	L2055	Y2180
F2184	T737	GLN	I1378	ASP	K1686	L1828	PRO	GLU	T2062	F2184
D2185	L738	SER	K1382	ILE	K1686	R1829	GLU	ASN	C2063	D2185
F2190	Y746	ARG	T1390	ILE	R1690	L1833	ILE	ILE	C2064	F2190
MET	P748	LEU	E1393	Q1571	Q1693	D1837	THR	THR	N2071	MET
GLY	F751	ALA	K1398	Y1572	Q1693	D1837	THR	THR	D2072	GLY
ASN	K754	ASN	R1401	I1573	Y1696	K1840	GLY	ARG	P2081	ASN
PRD	M763	PRD	Y1402	Y1574	E1697	SER	ARG	ILE	T2085	PRD
MET	Q1018	MET	L1272	S1575	Y1699	HIS	LYS	THR	E2086	MET
PHE	K1019	PHE	A1273	Y1576	K1700	GLU	ASN	ASN	G2087	PHE
VAL	N1020	VAL	L1403	I1577	S1701	VAL	LEU	ILE	T2087	VAL
THR	R1021	THR	S1274	T1578	T1702	THR	GLY	THR	T2087	THR
ASP	D1024	ASP	R1275	L1579	T1702	THR	SER	ARG	I2090	ASP
GLN	I1027	GLN	S1276	N1580	F1709	THR	ARG	HIS	H2091	GLN
VAL	E1028	VAL	S1277	D1581	S1715	THR	GLY	GLY	S2092	VAL
CYS	ALA	CYS	K1284	M1584	T1718	GLN	LYS	THR	T2093	CYS
LEU	LEU	LEU	P1289	S1419	T1718	THR	VAL	VAL	P2094	LEU
GLU	ALA	GLU	M1293	T1586	L1724	SER	PHE	ASN	I2095	GLU
VAL	ALA	VAL	V1293	I1587	L1724	ARG	GLU	VAL	F2096	VAL
GLY	THR	GLY	V1294	T1590	L1729	HIS	ASN	ALA	H2097	GLY
HIS	I781	THR	S1295	T1590	L1729	LEU	MET	THR	I2098	HIS
									Y2099	

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	24151	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS GLACIOS	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	60	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	36000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.028	Depositor
Minimum map value	-0.012	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.005	Depositor
Map size (Å)	297.69998, 297.69998, 297.69998	wwPDB
Map dimensions	260, 260, 260	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.145, 1.145, 1.145	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, POP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	H	0.69	1/264 (0.4%)	0.67	0/408
2	T	0.22	0/488	0.79	0/753
3	P	0.23	0/286	0.72	0/443
4	A	0.27	0/16721	0.45	0/22547
All	All	0.28	1/17759 (0.0%)	0.48	0/24151

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	1	A	OP3-P	-10.66	1.48	1.61

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	237	0	119	4	0
2	T	439	0	226	14	0
3	P	256	0	129	10	0
4	A	16383	0	16426	289	0
5	A	1	0	0	0	0
6	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	9	0	0	0	0
All	All	17326	0	16900	305	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 305 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:813:ARG:NH1	4:A:986:PRO:HD3	1.67	1.09
4:A:2064:CYS:HB2	4:A:2169:HIS:HE1	1.15	1.03
4:A:2064:CYS:CB	4:A:2169:HIS:HE1	1.76	0.98
4:A:2064:CYS:HB2	4:A:2169:HIS:CE1	1.97	0.98
4:A:813:ARG:HH12	4:A:986:PRO:HD3	1.35	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
4	A	1991/2276 (88%)	1923 (97%)	68 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
4	A	1852/2097 (88%)	1829 (99%)	23 (1%)	71 83

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	A	1390	THR
4	A	2051	ARG
4	A	1602	ARG
4	A	2072	ASP
4	A	655	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
4	A	622	ASN
4	A	760	HIS
4	A	1580	ASN
4	A	1704	HIS
4	A	1743	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	H	10/17 (58%)	4 (40%)	0
2	T	19/30 (63%)	9 (47%)	0
3	P	11/30 (36%)	1 (9%)	0
All	All	40/77 (51%)	14 (35%)	0

5 of 14 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	H	6	U
1	H	7	G
1	H	8	U
1	H	9	C
2	T	2	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	POP	A	2303	-	6,8,8	0.80	0	13,13,13	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	POP	A	2303	-	-	0/6/6/6	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-13038. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 130



Y Index: 130




Z Index: 130


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 133



Y Index: 116




Z Index: 134


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.005. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 201 nm^3 ; this corresponds to an approximate mass of 182 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.256 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.256 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.90	-	-
Author-provided FSC curve	3.88	4.37	3.94
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-13038 and PDB model 7ORI. Per-residue inclusion information can be found in section [3](#) on page [5](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.005 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.005).

9.4 Atom inclusion [i](#)



At the recommended contour level, 99% of all backbone atoms, 90% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.005) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.9040	<div></div> 0.4290
A	<div></div> 0.9000	<div></div> 0.4280
H	<div></div> 0.9870	<div></div> 0.4610
P	<div></div> 0.9730	<div></div> 0.4260
T	<div></div> 0.9570	<div></div> 0.4240

