



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 30, 2025 – 02:06 PM EDT

PDB ID : 9P6K / pdb_00009p6k
Title : Colorado Potato Beetle Glutathione S-transferase Sigma Class member 2
Authors : Moural, T.W.; Zhu, F.
Deposited on : 2025-06-19
Resolution : 2.84 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1367 (2.86-2.82)
Clashscore	180529	1455 (2.86-2.82)
Ramachandran outliers	177936	1422 (2.86-2.82)
Sidechain outliers	177891	1423 (2.86-2.82)
RSRZ outliers	164620	1368 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	221	
1	B	221	
1	C	221	
1	D	221	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 13968 atoms, of which 7080 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called glutathione transferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	210	Total	C	H	N	O	S	0	0	0
			3492	1118	1770	284	314	6			
1	B	210	Total	C	H	N	O	S	0	0	0
			3492	1118	1770	284	314	6			
1	C	210	Total	C	H	N	O	S	0	0	0
			3492	1118	1770	284	314	6			
1	D	210	Total	C	H	N	O	S	0	0	0
			3492	1118	1770	284	314	6			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	156	ILE	PHE	conflict	UNP A0A1P8PEX1
A	191	SER	PHE	conflict	UNP A0A1P8PEX1
A	205	LEU	PHE	conflict	UNP A0A1P8PEX1
A	206	SER	PHE	conflict	UNP A0A1P8PEX1
A	214	LEU	-	expression tag	UNP A0A1P8PEX1
A	215	GLU	-	expression tag	UNP A0A1P8PEX1
A	216	HIS	-	expression tag	UNP A0A1P8PEX1
A	217	HIS	-	expression tag	UNP A0A1P8PEX1
A	218	HIS	-	expression tag	UNP A0A1P8PEX1
A	219	HIS	-	expression tag	UNP A0A1P8PEX1
A	220	HIS	-	expression tag	UNP A0A1P8PEX1
A	221	HIS	-	expression tag	UNP A0A1P8PEX1
B	156	ILE	PHE	conflict	UNP A0A1P8PEX1
B	191	SER	PHE	conflict	UNP A0A1P8PEX1
B	205	LEU	PHE	conflict	UNP A0A1P8PEX1
B	206	SER	PHE	conflict	UNP A0A1P8PEX1
B	214	LEU	-	expression tag	UNP A0A1P8PEX1
B	215	GLU	-	expression tag	UNP A0A1P8PEX1
B	216	HIS	-	expression tag	UNP A0A1P8PEX1
B	217	HIS	-	expression tag	UNP A0A1P8PEX1
B	218	HIS	-	expression tag	UNP A0A1P8PEX1

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	219	HIS	-	expression tag	UNP A0A1P8PEX1
B	220	HIS	-	expression tag	UNP A0A1P8PEX1
B	221	HIS	-	expression tag	UNP A0A1P8PEX1
C	156	ILE	PHE	conflict	UNP A0A1P8PEX1
C	191	SER	PHE	conflict	UNP A0A1P8PEX1
C	205	LEU	PHE	conflict	UNP A0A1P8PEX1
C	206	SER	PHE	conflict	UNP A0A1P8PEX1
C	214	LEU	-	expression tag	UNP A0A1P8PEX1
C	215	GLU	-	expression tag	UNP A0A1P8PEX1
C	216	HIS	-	expression tag	UNP A0A1P8PEX1
C	217	HIS	-	expression tag	UNP A0A1P8PEX1
C	218	HIS	-	expression tag	UNP A0A1P8PEX1
C	219	HIS	-	expression tag	UNP A0A1P8PEX1
C	220	HIS	-	expression tag	UNP A0A1P8PEX1
C	221	HIS	-	expression tag	UNP A0A1P8PEX1
D	156	ILE	PHE	conflict	UNP A0A1P8PEX1
D	191	SER	PHE	conflict	UNP A0A1P8PEX1
D	205	LEU	PHE	conflict	UNP A0A1P8PEX1
D	206	SER	PHE	conflict	UNP A0A1P8PEX1
D	214	LEU	-	expression tag	UNP A0A1P8PEX1
D	215	GLU	-	expression tag	UNP A0A1P8PEX1
D	216	HIS	-	expression tag	UNP A0A1P8PEX1
D	217	HIS	-	expression tag	UNP A0A1P8PEX1
D	218	HIS	-	expression tag	UNP A0A1P8PEX1
D	219	HIS	-	expression tag	UNP A0A1P8PEX1
D	220	HIS	-	expression tag	UNP A0A1P8PEX1
D	221	HIS	-	expression tag	UNP A0A1P8PEX1

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: glutathione transferase

Chain A: 




- Molecule 1: glutathione transferase

Chain B: 



- Molecule 1: glutathione transferase

Chain C: 



- Molecule 1: glutathione transferase

Chain D: 





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	49.37Å 88.78Å 97.90Å 90.00° 99.73° 90.00°	Depositor
Resolution (Å)	29.20 – 2.84 29.20 – 2.84	Depositor EDS
% Data completeness (in resolution range)	97.3 (29.20-2.84) 95.7 (29.20-2.84)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.41 (at 2.85Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.221 , 0.268 0.221 , 0.268	Depositor DCC
R_{free} test set	1920 reflections (9.34%)	wwPDB-VP
Wilson B-factor (Å ²)	76.2	Xtriage
Anisotropy	0.437	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 81.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13968	wwPDB-VP
Average B, all atoms (Å ²)	117.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	0/1759	0.92	3/2370 (0.1%)
1	B	0.64	0/1759	0.94	3/2370 (0.1%)
1	C	0.56	0/1759	0.85	2/2370 (0.1%)
1	D	0.56	0/1759	0.90	4/2370 (0.2%)
All	All	0.61	0/7036	0.90	12/9480 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	75	LYS	CA-CB-CG	6.70	127.50	114.10
1	A	100	LYS	CB-CG-CD	6.57	126.41	111.30
1	D	51	GLN	CA-CB-CG	-6.13	101.85	114.10
1	D	145	LEU	N-CA-C	6.10	119.11	111.24
1	C	12	PHE	N-CA-CB	-5.97	102.12	111.56

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	51	GLN	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1722	1770	1769	50	0
1	B	1722	1770	1769	69	1
1	C	1722	1770	1769	62	1
1	D	1722	1770	1769	56	0
All	All	6888	7080	7076	233	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 233 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:12:PHE:HD1	1:B:202:ASN:ND2	1.38	1.22
1:B:12:PHE:CZ	1:B:205:LEU:HD21	1.92	1.04
1:B:103:VAL:HG12	1:B:167:MET:HE3	1.39	1.04
1:A:89:ILE:HD11	1:A:147:LEU:HD11	1.37	1.02
1:B:12:PHE:CD1	1:B:202:ASN:ND2	2.29	0.99

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:198:LYS:NZ	1:C:139:GLU:OE2[2_656]	2.15	0.05

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	208/221 (94%)	202 (97%)	6 (3%)	0	100	100
1	B	208/221 (94%)	201 (97%)	7 (3%)	0	100	100
1	C	208/221 (94%)	198 (95%)	10 (5%)	0	100	100
1	D	208/221 (94%)	202 (97%)	6 (3%)	0	100	100
All	All	832/884 (94%)	803 (96%)	29 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/201 (95%)	191 (100%)	0	100	100
1	B	191/201 (95%)	189 (99%)	2 (1%)	73	87
1	C	191/201 (95%)	190 (100%)	1 (0%)	86	94
1	D	191/201 (95%)	190 (100%)	1 (0%)	86	94
All	All	764/804 (95%)	760 (100%)	4 (0%)	86	94

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	173	LEU
1	B	180	THR
1	C	52	LEU
1	D	52	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	ASN
1	A	141	ASN
1	A	212	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	74	ASN
1	D	74	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	210/221 (95%)	-0.48	0 100 100	54, 87, 149, 204	0
1	B	210/221 (95%)	-0.46	2 (0%) 79 77	62, 105, 158, 197	0
1	C	210/221 (95%)	-0.28	0 100 100	65, 133, 194, 260	0
1	D	210/221 (95%)	-0.35	0 100 100	64, 125, 186, 230	0
All	All	840/884 (95%)	-0.39	2 (0%) 92 93	54, 111, 183, 260	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	12	PHE	2.9
1	B	205	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.