



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2024 – 01:30 AM EDT

PDB ID : 6P8E  
Title : Crystal structure of CDK4 in complex with CyclinD1 and P27  
Authors : Guiley, K.Z.; Stevenson, J.W.; Lou, K.; Barkovich, K.J.; Bunch, K.; Tripathi, S.M.; Shokat, K.M.; Rubin, S.M.  
Deposited on : 2019-06-07  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	249	<div> <div>92%</div> <div>8%</div> </div>
2	B	302	<div> <div>3%</div> <div>77%</div> <div>8%</div> <div>14%</div> </div>
3	C	72	<div> <div>3%</div> <div>61%</div> <div>18%</div> <div>21%</div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4709 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called G1/S-specific cyclin-D1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	247	Total	C	N	O	S	0	0	0
			1965	1248	335	362	20			

- Molecule 2 is a protein called Cyclin-dependent kinase 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	260	Total	C	N	O	S	0	0	0
			2057	1317	359	371	10			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	GLY	-	expression tag	UNP P11802
B	0	GLU	-	expression tag	UNP P11802
B	1	PHE	-	expression tag	UNP P11802
B	?	-	GLY	deletion	UNP P11802
B	?	-	GLY	deletion	UNP P11802
B	?	-	GLY	deletion	UNP P11802
B	43	GLU	GLY	engineered mutation	UNP P11802
B	44	GLU	GLY	engineered mutation	UNP P11802

- Molecule 3 is a protein called Cyclin-dependent kinase inhibitor 1B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	57	Total	C	N	O	S	0	0	0
			491	307	88	93	3			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	22	GLY	-	expression tag	UNP P46527
C	23	GLU	-	expression tag	UNP P46527

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	24	PHE	-	expression tag	UNP P46527

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total 2	Mg 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	105	Total 105	O 105	0	0
5	B	78	Total 78	O 78	0	0
5	C	11	Total 11	O 11	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: G1/S-specific cyclin-D1



- Molecule 2: Cyclin-dependent kinase 4



- Molecule 3: Cyclin-dependent kinase inhibitor 1B



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.41 Å   67.49 Å   187.28 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	93.64 – 2.30 93.64 – 2.30	Depositor EDS
% Data completeness (in resolution range)	94.0 (93.64-2.30) 93.2 (93.64-2.30)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.69 (at 2.29 Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, $R_{free}$	0.172   ,   0.222 0.174   ,   0.220	Depositor DCC
$R_{free}$ test set	1839 reflections (5.21%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.0	Xtriage
Anisotropy	0.248	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 49.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4709	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.45	0/1998	0.55	0/2699
2	B	0.38	0/2109	0.54	0/2865
3	C	0.37	0/505	0.47	0/678
All	All	0.41	0/4612	0.54	0/6242

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1965	0	2016	11	0
2	B	2057	0	2053	15	0
3	C	491	0	449	9	0
4	B	2	0	0	0	0
5	A	105	0	0	0	0
5	B	78	0	0	2	0
5	C	11	0	0	1	0
All	All	4709	0	4518	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:188:TYR:HB2	2:B:192:VAL:HG11	1.78	0.65
2:B:41:ASN:HB2	2:B:84:ILE:HG13	1.79	0.63
2:B:98:ARG:NH2	2:B:203:GLU:OE2	2.33	0.61
1:A:261:GLN:O	1:A:264:GLN:HG3	2.07	0.54
2:B:70:ARG:NH2	5:B:503:HOH:O	2.41	0.53
2:B:184:LEU:HB3	2:B:239:VAL:HG23	1.91	0.53
1:A:28:LEU:HD21	1:A:236:VAL:HB	1.91	0.52
1:A:173:GLU:H	1:A:173:GLU:CD	2.12	0.52
1:A:190:ALA:HA	1:A:195:PHE:CG	2.45	0.51
1:A:19:ASP:N	1:A:19:ASP:OD1	2.42	0.51
3:C:26:PRO:HD2	3:C:29:CYS:SG	2.53	0.49
2:B:274:THR:HG21	2:B:279:LYS:HG2	1.94	0.49
1:A:189:CYS:SG	1:A:251:ILE:HG12	2.53	0.48
2:B:200:ILE:O	2:B:204:MET:HG3	2.13	0.47
2:B:34:LEU:CD2	2:B:89:VAL:HG22	2.45	0.47
2:B:65:HIS:O	2:B:70:ARG:NH2	2.48	0.47
3:C:57:GLN:HG3	3:C:63:ASP:HA	1.97	0.47
3:C:42:THR:O	3:C:46:GLU:HG3	2.14	0.46
1:A:22:LEU:O	1:A:26:ARG:HG3	2.16	0.46
3:C:24:PHE:HB3	5:C:103:HOH:O	2.17	0.44
3:C:57:GLN:HG3	3:C:62:PHE:O	2.18	0.44
2:B:135:HIS:O	2:B:136:ARG:HB2	2.18	0.44
2:B:152:LYS:NZ	5:B:503:HOH:O	2.36	0.44
1:A:233:LEU:HA	1:A:233:LEU:HD23	1.73	0.43
2:B:260:GLU:H	2:B:260:GLU:CD	2.21	0.43
2:B:180:PRO:O	2:B:184:LEU:HD13	2.20	0.42
1:A:190:ALA:HA	1:A:195:PHE:HB2	2.02	0.42
3:C:68:LYS:HA	3:C:68:LYS:HD3	1.92	0.42
3:C:69:PRO:O	3:C:70:LEU:HD23	2.21	0.41
1:A:37:THR:HG21	2:B:62:ALA:HB1	2.02	0.41
3:C:67:HIS:HB3	3:C:68:LYS:HE2	2.03	0.41
1:A:131:SER:OG	3:C:31:ASN:HB2	2.21	0.40
2:B:295:ASP:N	2:B:295:ASP:OD1	2.46	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	245/249 (98%)	242 (99%)	3 (1%)	0	100	100
2	B	254/302 (84%)	245 (96%)	9 (4%)	0	100	100
3	C	55/72 (76%)	54 (98%)	1 (2%)	0	100	100
All	All	554/623 (89%)	541 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	222/224 (99%)	218 (98%)	4 (2%)	59	75
2	B	226/260 (87%)	222 (98%)	4 (2%)	59	75
3	C	53/66 (80%)	53 (100%)	0	100	100
All	All	501/550 (91%)	493 (98%)	8 (2%)	62	78

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	188	LEU
1	A	245	ARG
1	A	254	LEU
1	A	263	GLN
2	B	38	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	43	GLU
2	B	70	ARG
2	B	256	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	247/249 (99%)	0.04	3 (1%) 79 83	21, 41, 74, 134	0
2	B	260/302 (86%)	0.18	10 (3%) 40 47	30, 59, 104, 145	0
3	C	57/72 (79%)	0.22	2 (3%) 44 51	43, 77, 117, 141	0
All	All	564/623 (90%)	0.12	15 (2%) 54 62	21, 51, 104, 145	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	27	HIS	5.5
3	C	71	GLU	4.8
2	B	157	GLY	4.6
1	A	23	LEU	3.9
2	B	238	ASP	3.3
2	B	239	VAL	3.2
2	B	83	GLU	3.0
1	A	261	GLN	2.9
2	B	19	THR	2.6
3	C	24	PHE	2.6
2	B	28	SER	2.5
2	B	232	GLU	2.5
1	A	21	ASN	2.4
2	B	187	THR	2.3
2	B	237	ARG	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	MG	B	402	1/1	0.51	0.21	83,83,83,83	0
4	MG	B	401	1/1	0.92	0.09	50,50,50,50	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.