



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 22, 2024 – 08:55 AM EDT

PDB ID : 3PU9
Title : Crystal structure of serine/threonine phosphatase *Sphaerobacter thermophilus* DSM 20745
Authors : Nocek, B.; Tesar, C.; Clancy, S.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2010-12-03
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1935 (1.56-1.56)
Clashscore	180529	2073 (1.56-1.56)
Ramachandran outliers	177936	2037 (1.56-1.56)
Sidechain outliers	177891	2034 (1.56-1.56)
RSRZ outliers	164620	1935 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	242	<div> <div>4%</div> <div>88%</div> <div>9%</div> <div>.</div> </div>
1	B	242	<div> <div>14%</div> <div>81%</div> <div>14%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	501	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3867 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein serine/threonine phosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	236	Total	C	N	O	Se	1	4	0
			1750	1089	328	327	6			
1	B	236	Total	C	N	O	Se	0	6	0
			1773	1104	330	333	6			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP D1C2D8
A	-1	ASN	-	expression tag	UNP D1C2D8
A	0	ALA	-	expression tag	UNP D1C2D8
B	-2	SER	-	expression tag	UNP D1C2D8
B	-1	ASN	-	expression tag	UNP D1C2D8
B	0	ALA	-	expression tag	UNP D1C2D8

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		
2	B	2	Total	Mg	0	0
			2	2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	189	Total	O	0	0
			189	189		
4	B	144	Total	O	0	1
			145	145		



3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein serine/threonine phosphatase



- Molecule 1: Protein serine/threonine phosphatase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.17Å 54.71Å 63.26Å 90.00° 99.51° 90.00°	Depositor
Resolution (Å)	40.00 – 1.55 40.00 – 1.55	Depositor EDS
% Data completeness (in resolution range)	98.1 (40.00-1.55) 98.1 (40.00-1.55)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.83 (at 1.55Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.178 , 0.217 0.185 , 0.226	Depositor DCC
R_{free} test set	2988 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	18.0	Xtriage
Anisotropy	0.162	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 55.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.007 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3867	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 53.33 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.2827e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.00	2/1773 (0.1%)	0.90	3/2392 (0.1%)
1	B	0.96	1/1802 (0.1%)	1.01	9/2430 (0.4%)
All	All	0.98	3/3575 (0.1%)	0.96	12/4822 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	151	GLU	CB-CG	-5.64	1.41	1.52
1	B	9	GLU	CB-CG	-5.18	1.42	1.52
1	A	43	ASP	CB-CG	5.11	1.62	1.51

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	87	ARG	NE-CZ-NH2	11.03	125.82	120.30
1	B	111	ARG	NE-CZ-NH1	9.59	125.10	120.30
1	B	87	ARG	NE-CZ-NH1	-9.48	115.56	120.30
1	B	111	ARG	NE-CZ-NH2	-9.36	115.62	120.30
1	B	211	ASP	CB-CG-OD2	5.91	123.62	118.30
1	A	219	ARG	NE-CZ-NH2	-5.90	117.35	120.30
1	B	64	ARG	NE-CZ-NH2	5.66	123.13	120.30
1	A	25	ASP	CB-CG-OD1	5.53	123.27	118.30
1	B	134	ARG	NE-CZ-NH1	5.47	123.03	120.30
1	B	29	LEU	CB-CG-CD1	5.35	120.10	111.00
1	A	53	ARG	NE-CZ-NH2	-5.32	117.64	120.30
1	B	208	ARG	NE-CZ-NH2	-5.11	117.75	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1750	0	1793	33	0
1	B	1773	0	1826	62	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	6	0	7	4	0
4	A	189	0	0	5	0
4	B	145	0	0	10	0
All	All	3867	0	3626	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:238:LEU:O	1:B:102:LEU:HD13	1.47	1.11
1:A:211:ASP:HB2	4:A:347:HOH:O	1.52	1.07
3:A:501:GOL:H11	1:B:230:ARG:HD2	1.51	0.90
1:B:96:ASP:OD1	1:B:111:ARG:HD2	1.77	0.85
1:B:232[B]:MSE:HG2	4:B:310:HOH:O	1.73	0.84
1:A:80:HIS:HD2	1:A:136:GLY:H	1.28	0.82
1:B:83:LEU:HD11	1:B:91[C]:VAL:CG2	2.10	0.82
1:A:10:LEU:CD2	1:A:29:LEU:HD11	2.10	0.81
1:A:238:LEU:CB	1:B:102:LEU:HD22	2.13	0.79
1:B:30:VAL:CG2	1:B:67[A]:ASN:OD1	2.31	0.79
1:B:87:ARG:HB3	1:B:91[A]:VAL:HG12	1.69	0.74
1:A:80:HIS:CD2	1:A:136:GLY:H	2.07	0.72
1:A:238:LEU:O	1:B:102:LEU:CD1	2.33	0.72
1:A:238:LEU:N	1:A:239:GLY:HA2	2.05	0.72
1:A:238:LEU:CB	1:B:102:LEU:CD2	2.71	0.69
1:A:12:VAL:HB	1:B:232[A]:MSE:HG2	1.74	0.67
1:B:83:LEU:HD11	1:B:91[B]:VAL:CG1	2.25	0.67
1:B:87:ARG:CB	1:B:91[A]:VAL:CG1	2.75	0.65
1:B:83:LEU:HD11	1:B:91[C]:VAL:HG23	1.78	0.65
1:B:30:VAL:HG21	1:B:67[A]:ASN:OD1	1.97	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:87:ARG:CB	1:B:91[A]:VAL:HG12	2.25	0.65
3:A:501:GOL:H11	1:B:230:ARG:CD	2.26	0.65
1:B:131:SER:HB3	4:B:326:HOH:O	1.96	0.64
1:A:238:LEU:CA	1:B:102:LEU:HD22	2.28	0.64
1:B:233:ALA:N	4:B:310:HOH:O	2.31	0.64
1:B:87:ARG:HG3	1:B:91[A]:VAL:HG11	1.79	0.63
1:B:4:MSE:HA	1:B:4:MSE:HE2	1.80	0.63
1:A:22:GLU:HA	3:A:501:GOL:H31	1.80	0.61
1:B:91[B]:VAL:O	1:B:91[B]:VAL:HG13	2.01	0.60
1:A:238:LEU:CA	1:B:102:LEU:CD2	2.79	0.60
1:B:87:ARG:HB2	1:B:91[B]:VAL:HG12	1.82	0.60
1:A:54:LEU:HD22	1:A:91[A]:VAL:HG11	1.84	0.59
1:A:238:LEU:CB	1:B:6:VAL:CG2	2.82	0.58
1:A:211:ASP:CB	4:A:347:HOH:O	2.26	0.57
1:B:87:ARG:HB2	1:B:91[B]:VAL:CG1	2.34	0.57
1:B:83:LEU:HD11	1:B:91[B]:VAL:HG11	1.86	0.57
1:B:83:LEU:HD11	1:B:91[B]:VAL:HG13	1.87	0.57
1:A:238:LEU:HA	1:B:102:LEU:CD2	2.35	0.57
1:A:238:LEU:CB	1:B:6:VAL:HG23	2.36	0.56
1:B:36:GLY:C	1:B:37:PHE:CD1	2.80	0.54
1:A:4:MSE:N	4:A:391:HOH:O	2.39	0.54
1:B:114:GLU:HA	4:B:326:HOH:O	2.08	0.53
1:B:64:ARG:HH21	1:B:64:ARG:CG	2.23	0.52
1:B:169:ARG:HB3	4:B:357:HOH:O	2.09	0.51
1:A:238:LEU:HA	1:B:102:LEU:CD1	2.41	0.50
1:B:30:VAL:HG23	1:B:67[A]:ASN:OD1	2.10	0.50
1:A:54:LEU:HD11	1:A:83:LEU:HD11	1.94	0.49
1:B:131:SER:CB	4:B:326:HOH:O	2.57	0.49
1:B:108:LEU:HD23	1:B:147:GLU:HG3	1.95	0.49
1:B:64:ARG:NH1	4:B:251:HOH:O	2.46	0.49
1:B:23:SER:O	1:B:52:LYS:HD3	2.13	0.48
1:B:87:ARG:HB2	1:B:91[A]:VAL:CG1	2.43	0.48
1:A:238:LEU:H	1:A:239:GLY:HA2	1.77	0.47
1:A:54:LEU:CD2	1:A:91[A]:VAL:CG1	2.92	0.47
1:A:171:GLY:HA2	4:A:425:HOH:O	2.14	0.47
1:A:54:LEU:HD22	1:A:91[A]:VAL:CG1	2.45	0.46
1:A:80:HIS:HD2	1:A:136:GLY:N	2.06	0.46
1:B:64:ARG:HH21	1:B:64:ARG:HG3	1.80	0.46
1:A:10:LEU:HD22	1:A:29:LEU:HD11	1.95	0.46
1:B:142:ARG:HH11	1:B:142:ARG:HG3	1.81	0.46
1:B:114:GLU:OE1	4:B:435:HOH:O	2.20	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:54:LEU:HD21	1:B:91[C]:VAL:HG21	1.98	0.45
1:A:127:GLN:C	1:A:128:ILE:HD13	2.37	0.45
1:B:107:VAL:HG21	1:B:154:LEU:HD23	1.98	0.45
1:B:49:ALA:O	1:B:53:ARG:HG3	2.17	0.45
1:A:211:ASP:O	1:A:212:ASP:C	2.55	0.44
1:A:238:LEU:HA	1:B:102:LEU:HD21	1.99	0.44
1:B:87:ARG:CB	1:B:91[B]:VAL:CG1	2.96	0.44
1:B:64:ARG:HG3	1:B:64:ARG:NH2	2.32	0.43
1:A:238:LEU:C	1:B:102:LEU:HD22	2.39	0.43
3:A:501:GOL:H32	4:A:317:HOH:O	2.18	0.43
1:B:115:VAL:HG13	1:B:164:ILE:HG13	2.01	0.43
1:B:227:GLU:HG3	1:B:229:ILE:HG13	2.01	0.43
1:A:10:LEU:HD23	1:A:29:LEU:HD11	1.97	0.42
1:A:238:LEU:CB	1:B:102:LEU:HD21	2.47	0.41
1:B:83:LEU:CD1	1:B:91[B]:VAL:CG1	2.98	0.41
1:B:87:ARG:CB	1:B:91[B]:VAL:HG12	2.47	0.41
1:B:91[B]:VAL:HG12	4:B:245:HOH:O	2.21	0.41
1:B:142:ARG:HH11	1:B:142:ARG:CG	2.34	0.40
1:B:114:GLU:HG3	1:B:114:GLU:O	2.21	0.40
1:B:64:ARG:CG	1:B:64:ARG:NH2	2.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	238/242 (98%)	234 (98%)	4 (2%)	0	100	100
1	B	241/242 (100%)	237 (98%)	4 (2%)	0	100	100
All	All	479/484 (99%)	471 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	177/176 (101%)	175 (99%)	2 (1%)	70	49
1	B	182/176 (103%)	177 (97%)	5 (3%)	40	12
All	All	359/352 (102%)	352 (98%)	7 (2%)	50	24

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	LEU
1	A	226	VAL
1	B	29	LEU
1	B	47	SER
1	B	64	ARG
1	B	106	SER
1	B	142	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	80	HIS
1	A	122	ASN
1	A	191	GLN
1	B	191	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	501	-	5,5,5	0.87	0	5,5,5	2.36	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	501	-	-	2/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	GOL	O2-C2-C3	4.45	127.61	109.18
3	A	501	GOL	O3-C3-C2	2.76	122.80	110.38

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	GOL	C1-C2-C3-O3
3	A	501	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	GOL	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	231/242 (95%)	0.50	10 (4%) 40 48	13, 24, 42, 65	3 (1%)
1	B	231/242 (95%)	0.90	35 (15%) 6 7	7, 28, 54, 99	5 (2%)
All	All	462/484 (95%)	0.70	45 (9%) 15 17	7, 26, 51, 99	8 (1%)

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	132	GLY	8.6
1	B	133	GLY	7.0
1	B	102	LEU	5.4
1	B	131	SER	4.8
1	B	103	ALA	4.3
1	B	145	VAL	4.0
1	B	128	ILE	3.5
1	B	223	VAL	3.5
1	B	129	SER	3.4
1	B	185	GLY	3.2
1	B	146	LEU	3.2
1	B	144	ARG	3.1
1	B	47	SER	3.1
1	B	122	ASN	3.1
1	A	229	ILE	3.1
1	A	237	PRO	3.0
1	A	211	ASP	2.9
1	B	233	ALA	2.8
1	B	19	GLY	2.7
1	B	105	ARG	2.6
1	A	226	VAL	2.6
1	A	223	VAL	2.5
1	B	114	GLU	2.5
1	B	46	GLY	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	34	GLY	2.4
1	A	224	GLU	2.4
1	B	184	ALA	2.4
1	B	227	GLU	2.3
1	B	237	PRO	2.3
1	B	104	SER	2.3
1	A	171	GLY	2.3
1	B	238	LEU	2.3
1	B	226	VAL	2.3
1	B	48	GLY	2.3
1	A	225	GLU	2.2
1	B	130	GLU	2.2
1	B	106	SER	2.2
1	B	64	ARG	2.2
1	B	37	PHE	2.2
1	A	238	LEU	2.1
1	B	183	VAL	2.1
1	B	224	GLU	2.1
1	A	239	GLY	2.0
1	B	138	TYR	2.0
1	B	115	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MG	B	302	1/1	0.88	0.22	27,27,27,27	1
3	GOL	A	501	6/6	0.91	0.15	15,17,23,26	6

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MG	A	302	1/1	0.97	0.19	13,13,13,13	0
2	MG	A	301	1/1	0.99	0.22	12,12,12,12	1
2	MG	B	301	1/1	0.99	0.19	6,6,6,6	1

6.5 Other polymers [i](#)

There are no such residues in this entry.