



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2024 – 11:35 PM EDT

PDB ID : 5PWU  
Title : PanDDA analysis group deposition – Crystal Structure of SP100 after initial refinement with no ligand modelled (structure 18)  
Authors : Pearce, N.M.; Krojer, T.; Talon, R.; Bradley, A.R.; Fairhead, M.; Sethi, R.; Wright, N.; MacLean, E.; Collins, P.; Brandao-Neto, J.; Douangamath, A.; Renjie, Z.; Dias, A.; Ng, J.; Brennan, P.E.; Cox, O.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.; von Delft, F.  
Deposited on : 2017-02-08  
Resolution : 1.44 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2021 (1.46-1.42)
Clashscore	141614	2086 (1.46-1.42)
Ramachandran outliers	138981	2047 (1.46-1.42)
Sidechain outliers	138945	2047 (1.46-1.42)
RSRZ outliers	127900	1993 (1.46-1.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	180	
1	B	180	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

Validation Pipeline (wwPDB-VP) : 2.37.1

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ZN	A	903	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3620 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nuclear autoantigen Sp-100.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	178	Total	C	N	O	S	0	10	0
			1536	971	268	278	19			
1	B	177	Total	C	N	O	S	0	7	0
			1480	940	258	263	19			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Zn	0	0
			3	3		
2	B	2	Total	Zn	0	0
			2	2		

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	311	Total	O	0	3
			311	311		
5	B	268	Total	O	0	0
			268	268		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Nuclear autoantigen Sp-100

Chain A:  3% 86% 12% ..



#### • Molecule 1: Nuclear autoantigen Sp-100

Chain B:  13% 84% 13% ..



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	127.69Å 45.34Å 83.36Å 90.00° 102.35° 90.00°	Depositor
Resolution (Å)	36.67 – 1.44 36.67 – 1.44	Depositor EDS
% Data completeness (in resolution range)	97.8 (36.67-1.44) 97.9 (36.67-1.44)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.26 (at 1.44Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.175 , 0.203 0.184 , 0.211	Depositor DCC
$R_{free}$ test set	4179 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.4	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 48.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3620	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, MES, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.42	7/1603 (0.4%)	1.29	13/2154 (0.6%)
1	B	1.39	4/1538 (0.3%)	1.35	11/2071 (0.5%)
All	All	1.41	11/3141 (0.4%)	1.32	24/4225 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	742	CYS	CB-SG	-14.68	1.57	1.82
1	A	763	GLU	CD-OE1	-9.91	1.14	1.25
1	A	797	SER	CB-OG	-7.98	1.31	1.42
1	A	803[A]	ARG	CZ-NH2	-6.19	1.25	1.33
1	A	803[B]	ARG	CZ-NH2	-6.19	1.25	1.33
1	B	865	GLU	CG-CD	5.80	1.60	1.51
1	A	723	SER	CA-CB	5.39	1.61	1.52
1	B	823	ASN	CG-ND2	-5.38	1.19	1.32
1	B	780	GLU	CD-OE2	5.33	1.31	1.25
1	A	723	SER	CB-OG	5.28	1.49	1.42
1	B	711	TRP	CE3-CZ3	5.18	1.47	1.38

All (24) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	829[A]	ARG	NE-CZ-NH2	-16.48	112.06	120.30
1	B	829[B]	ARG	NE-CZ-NH2	-16.48	112.06	120.30
1	A	851	ASP	CB-CG-OD1	9.64	126.98	118.30
1	A	803[A]	ARG	NE-CZ-NH1	8.41	124.51	120.30
1	A	803[B]	ARG	NE-CZ-NH1	8.41	124.51	120.30
1	B	855	ARG	NE-CZ-NH1	8.34	124.47	120.30
1	A	742	CYS	CA-CB-SG	7.53	127.56	114.00
1	A	787	TYR	CB-CG-CD2	7.42	125.45	121.00
1	A	803[A]	ARG	NE-CZ-NH2	-6.66	116.97	120.30
1	A	803[B]	ARG	NE-CZ-NH2	-6.66	116.97	120.30
1	B	855	ARG	NE-CZ-NH2	-6.66	116.97	120.30
1	A	838	ARG	NE-CZ-NH1	-6.27	117.17	120.30
1	A	780	GLU	OE1-CD-OE2	6.23	130.78	123.30
1	B	789	ASP	CB-CG-OD2	-6.07	112.84	118.30
1	B	829[A]	ARG	NH1-CZ-NH2	6.05	126.06	119.40
1	B	829[B]	ARG	NH1-CZ-NH2	6.05	126.06	119.40
1	B	787	TYR	CB-CG-CD1	5.98	124.59	121.00
1	A	855	ARG	NE-CZ-NH2	-5.79	117.41	120.30
1	B	827	TYR	CB-CG-CD1	5.77	124.46	121.00
1	B	746	ARG	NE-CZ-NH2	-5.60	117.50	120.30
1	B	794	PHE	CB-CG-CD1	-5.59	116.89	120.80
1	A	722	ARG	NE-CZ-NH2	-5.54	117.53	120.30
1	A	801	TYR	CB-CG-CD1	-5.14	117.91	121.00
1	A	744	PHE	CB-CG-CD2	-5.06	117.26	120.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	755	PRO	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1536	0	1496	10	0
1	B	1480	0	1414	11	0
2	A	3	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	2	0	0	0	0
3	A	12	0	12	2	0
4	A	4	0	6	2	0
4	B	4	0	6	1	0
5	A	311	0	0	11	2
5	B	268	0	0	8	4
All	All	3620	0	2934	25	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:803[A]:ARG:NH2	5:A:1001:HOH:O	1.66	1.24
1:A:803[B]:ARG:NH1	5:A:1003:HOH:O	1.86	1.08
1:A:829[A]:ARG:NH1	5:A:1002:HOH:O	1.85	1.08
3:A:904:MES:H81	5:A:1225:HOH:O	1.65	0.96
1:B:826[B]:MET:HE2	5:B:1197:HOH:O	1.72	0.89
1:A:756:GLU:HG3	5:A:1089:HOH:O	1.77	0.82
1:B:750:ILE:HD11	5:B:1001:HOH:O	1.86	0.74
1:B:826[B]:MET:SD	5:B:1015:HOH:O	2.46	0.72
1:A:862[A]:ASP:OD1	5:A:1004:HOH:O	2.07	0.71
5:A:1001:HOH:O	1:B:773:PRO:HG2	1.93	0.68
4:B:903:EDO:O2	5:B:1001:HOH:O	2.09	0.68
1:B:716[A]:CYS:SG	5:B:1156:HOH:O	2.53	0.66
1:B:849:ARG:O	1:B:854[B]:THR:HG21	1.98	0.63
1:B:757:SER:O	1:B:759:SER:N	2.35	0.59
1:A:802:ASN:O	3:A:904:MES:H32	2.02	0.58
4:A:905:EDO:H21	5:A:1054:HOH:O	2.04	0.57
1:B:817:LYS:NZ	5:B:1006:HOH:O	2.39	0.55
4:A:905:EDO:C2	5:A:1054:HOH:O	2.57	0.53
1:A:824:GLU:HB2	1:A:826[B]:MET:HE2	1.91	0.51
1:A:762:GLN:NE2	5:A:1008:HOH:O	2.43	0.51
1:B:785:LYS:HE3	5:B:1223:HOH:O	2.12	0.48
1:B:854[A]:THR:HG21	5:B:1007:HOH:O	2.13	0.47
1:A:810:GLN:NE2	5:A:1009:HOH:O	2.48	0.46
1:B:757:SER:C	1:B:759:SER:N	2.70	0.45
1:A:849:ARG:O	1:A:854[B]:THR:HG21	2.17	0.44

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:1224:HOH:O	5:B:1224:HOH:O[2_555]	1.60	0.60
5:A:1103:HOH:O	5:A:1240:HOH:O[1_545]	1.88	0.32
5:B:1020:HOH:O	5:B:1020:HOH:O[2_556]	1.91	0.29
5:A:1161:HOH:O	5:B:1018:HOH:O[3_545]	2.13	0.07
5:B:1041:HOH:O	5:B:1156:HOH:O[2_556]	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	186/180 (103%)	182 (98%)	4 (2%)	0	100	100
1	B	182/180 (101%)	173 (95%)	7 (4%)	2 (1%)	14	2
All	All	368/360 (102%)	355 (96%)	11 (3%)	2 (0%)	25	8

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	757	SER
1	B	758	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/171 (104%)	178 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	B	165/171 (96%)	160 (97%)	5 (3%)	41 9
All	All	343/342 (100%)	338 (98%)	5 (2%)	78 34

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	711	TRP
1	B	850[A]	GLU
1	B	850[B]	GLU
1	B	862[A]	ASP
1	B	862[B]	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 5 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	MES	A	904	-	12,12,12	0.65	0	14,16,16	2.89	5 (35%)
4	EDO	A	905	-	3,3,3	0.55	0	2,2,2	0.04	0
4	EDO	B	903	-	3,3,3	0.74	0	2,2,2	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MES	A	904	-	-	1/6/14/14	0/1/1/1
4	EDO	A	905	-	-	1/1/1/1	-
4	EDO	B	903	-	-	0/1/1/1	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	904	MES	C5-N4-C3	6.72	123.95	108.83
3	A	904	MES	C7-N4-C3	4.97	123.94	111.23
3	A	904	MES	C6-C5-N4	4.09	116.31	110.10
3	A	904	MES	C2-C3-N4	3.58	115.54	110.10
3	A	904	MES	C6-O1-C2	3.46	121.45	109.89

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	904	MES	C8-C7-N4-C3
4	A	905	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	904	MES	2	0
4	A	905	EDO	2	0
4	B	903	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	178/180 (98%)	0.03	6 (3%) 45 46	13, 20, 49, 97	0
1	B	177/180 (98%)	0.37	23 (12%) 3 3	14, 23, 76, 128	0
All	All	355/360 (98%)	0.20	29 (8%) 11 12	13, 22, 65, 128	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	806	SER	9.7
1	B	757	SER	8.8
1	A	735	ALA	8.5
1	B	800	TYR	5.9
1	B	805	GLY	5.7
1	B	759	SER	5.2
1	B	758	GLN	5.1
1	B	804	GLU	4.9
1	A	736	ASN	4.2
1	B	760	GLY	4.0
1	B	876	GLU	3.9
1	B	875	GLN	3.7
1	B	803	ARG	3.4
1	A	703	ASN	3.1
1	B	701	ASN	3.1
1	B	810	GLN	3.0
1	B	807	GLN	3.0
1	A	737	LYS	2.9
1	B	700	GLU	2.9
1	B	808	GLY	2.7
1	A	878	SER	2.7
1	B	735	ALA	2.6
1	B	802	ASN	2.5
1	A	701	ASN	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	711	TRP	2.3
1	B	755	PRO	2.3
1	B	801	TYR	2.1
1	B	738	ASN	2.1
1	B	756	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ZN	A	903	1/1	0.56	0.61	95,95,95,95	1
4	EDO	A	905	4/4	0.85	0.27	43,52,56,65	0
2	ZN	A	902	1/1	0.93	0.06	27,27,27,27	0
4	EDO	B	903	4/4	0.96	0.06	27,28,29,30	0
3	MES	A	904	12/12	0.97	0.13	15,20,22,22	12
2	ZN	B	902	1/1	0.98	0.05	21,21,21,21	0
2	ZN	A	901	1/1	0.98	0.05	21,21,21,21	0
2	ZN	B	901	1/1	0.99	0.07	18,18,18,18	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.