



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 26, 2024 – 05:56 pm BST

PDB ID : 8QC9
Title : Crystal structure of the outer membrane decaheme cytochrome MtrC (A293Boc-Lys)
Authors : Nash, B.W.; Lockwood, C.J.; Edwards, M.J.; Whiting, K.; Butt, J.N.; Clarke, T.A.
Deposited on : 2023-08-25
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.002 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.38.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	679	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ACT	A	723	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5971 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Extracellular iron oxide respiratory system surface decaheme cytochrome c component MtrC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	626	Total	C	N	O	S	0	3	0
			4719	2923	812	952	32			

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	3	MET	-	initiating methionine	UNP Q8EG34
A	4	LYS	-	expression tag	UNP Q8EG34
A	5	PHE	-	expression tag	UNP Q8EG34
A	6	LYS	-	expression tag	UNP Q8EG34
A	7	LEU	-	expression tag	UNP Q8EG34
A	8	ASN	-	expression tag	UNP Q8EG34
A	9	LEU	-	expression tag	UNP Q8EG34
A	10	ILE	-	expression tag	UNP Q8EG34
A	11	THR	-	expression tag	UNP Q8EG34
A	12	LEU	-	expression tag	UNP Q8EG34
A	13	ALA	-	expression tag	UNP Q8EG34
A	14	LEU	-	expression tag	UNP Q8EG34
A	15	LEU	-	expression tag	UNP Q8EG34
A	16	ALA	-	expression tag	UNP Q8EG34
A	17	ASN	-	expression tag	UNP Q8EG34
A	18	THR	-	expression tag	UNP Q8EG34
A	19	GLY	-	expression tag	UNP Q8EG34
A	20	LEU	-	expression tag	UNP Q8EG34
A	21	ALA	-	expression tag	UNP Q8EG34
A	22	VAL	-	expression tag	UNP Q8EG34
A	23	ALA	-	expression tag	UNP Q8EG34
A	24	ALA	-	expression tag	UNP Q8EG34
A	25	ASP	-	expression tag	UNP Q8EG34
A	293	LBY	ALA	engineered mutation	UNP Q8EG34
A	672	SER	-	expression tag	UNP Q8EG34
A	673	ALA	-	expression tag	UNP Q8EG34

Continued on next page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	674	TRP	-	expression tag	UNP Q8EG34
A	675	SER	-	expression tag	UNP Q8EG34
A	676	HIS	-	expression tag	UNP Q8EG34
A	677	PRO	-	expression tag	UNP Q8EG34
A	678	GLN	-	expression tag	UNP Q8EG34
A	679	PHE	-	expression tag	UNP Q8EG34
A	680	GLU	-	expression tag	UNP Q8EG34
A	681	LYS	-	expression tag	UNP Q8EG34

- [illegible]

[illegible]


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	5	Total	Ca		
			5	5	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O		
			4	2	2	0	0
4	A	1	Total	C	O		
			4	2	2	0	0
4	A	1	Total	C	O		
			4	2	2	0	0
4	A	1	Total	C	O		
			4	2	2	0	0
4	A	1	Total	C	O		
			4	2	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	785	Total	O	0	0
			785	785		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Extracellular iron oxide respiratory system surface decaheme cytochrome c component MtrC



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	53.08Å 90.02Å 154.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	58.63 – 2.00 58.63 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (58.63-2.00) 99.8 (58.63-2.00)	Depositor EDS
R_{merge}	0.35	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 2.00Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.173 , 0.214 0.173 , 0.214	Depositor DCC
R_{free} test set	2634 reflections (5.17%)	wwPDB-VP
Wilson B-factor (Å ²)	22.3	Xtriage
Anisotropy	0.338	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 51.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5971	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, LBY, HEC, EDO, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/4794	0.60	0/6509

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4719	0	4536	36	0
2	A	430	0	301	16	0
3	A	5	0	0	0	0
4	A	24	0	36	3	0
5	A	8	0	6	4	0
6	A	785	0	0	11	2
All	All	5971	0	4879	47	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 47 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:51:LEU:H	5:A:723:ACT:H1	1.21	1.05
1:A:473:LEU:O	6:A:801:HOH:O	1.85	0.94
1:A:465:LYS:NZ	6:A:802:HOH:O	2.04	0.89
1:A:571:LEU:HD11	2:A:701:HEC:HMD3	1.65	0.77
1:A:289:ILE:HD11	2:A:705:HEC:HBA1	1.71	0.73

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:1389:HOH:O	6:A:1564:HOH:O[3_554]	1.93	0.27
6:A:1173:HOH:O	6:A:1455:HOH:O[1_455]	2.19	0.01

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	626/679 (92%)	611 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	516/550 (94%)	511 (99%)	5 (1%)	73	78

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	125	LYS
1	A	198	HIS
1	A	521	LEU
1	A	595	ASN
1	A	625	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	251	ASN
1	A	511	GLN
1	A	512	HIS
1	A	552	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	LBY	A	293	1	14,15,16	1.05	0	14,19,21	1.20	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LBY	A	293	1	-	6/14/15/17	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	293	LBY	O2-CZ-O1	-2.95	120.24	125.62

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	293	LBY	O2-CZ-NZ-CE
1	A	293	LBY	O1-CZ-O2-CT
1	A	293	LBY	NZ-CZ-O2-CT
1	A	293	LBY	O1-CZ-NZ-CE
1	A	293	LBY	CG-CD-CE-NZ

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	293	LBY	3	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 5 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEC	A	708	1	32,50,50	2.07	6 (18%)	24,82,82	1.50	5 (20%)
5	ACT	A	722	-	3,3,3	1.52	1 (33%)	3,3,3	1.41	0
2	HEC	A	710	1	32,50,50	2.14	6 (18%)	24,82,82	1.94	7 (29%)
2	HEC	A	701	1	32,50,50	2.19	4 (12%)	24,82,82	2.02	7 (29%)
2	HEC	A	703	1	32,50,50	2.01	5 (15%)	24,82,82	1.53	4 (16%)
4	EDO	A	718	-	3,3,3	0.47	0	2,2,2	1.14	0
2	HEC	A	704	1	32,50,50	2.02	3 (9%)	24,82,82	1.84	6 (25%)
2	HEC	A	707	1	32,50,50	2.07	5 (15%)	24,82,82	1.56	3 (12%)
2	HEC	A	702	1	32,50,50	1.99	3 (9%)	24,82,82	2.18	7 (29%)
2	HEC	A	709	1	32,50,50	2.12	5 (15%)	24,82,82	1.63	6 (25%)
4	EDO	A	719	-	3,3,3	0.46	0	2,2,2	0.27	0
4	EDO	A	717	-	3,3,3	0.56	0	2,2,2	0.15	0
4	EDO	A	720	-	3,3,3	0.54	0	2,2,2	0.37	0
4	EDO	A	721	-	3,3,3	0.46	0	2,2,2	0.88	0
5	ACT	A	723	3	3,3,3	0.85	0	3,3,3	1.78	2 (66%)
4	EDO	A	716	-	3,3,3	0.49	0	2,2,2	0.39	0
2	HEC	A	706	1	32,50,50	1.94	3 (9%)	24,82,82	1.81	3 (12%)
2	HEC	A	705	1	32,50,50	2.23	6 (18%)	24,82,82	1.49	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEC	A	708	1	-	4/10/54/54	-
2	HEC	A	710	1	-	0/10/54/54	-
2	HEC	A	701	1	-	3/10/54/54	-
2	HEC	A	703	1	-	2/10/54/54	-
4	EDO	A	718	-	-	1/1/1/1	-
2	HEC	A	704	1	-	1/10/54/54	-
2	HEC	A	707	1	-	4/10/54/54	-
2	HEC	A	702	1	-	2/10/54/54	-
2	HEC	A	709	1	-	1/10/54/54	-
4	EDO	A	719	-	-	1/1/1/1	-
4	EDO	A	717	-	-	0/1/1/1	-
4	EDO	A	720	-	-	1/1/1/1	-
4	EDO	A	721	-	-	1/1/1/1	-
4	EDO	A	716	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEC	A	706	1	-	1/10/54/54	-
2	HEC	A	705	1	-	2/10/54/54	-

The worst 5 of 47 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	709	HEC	C3C-C2C	-6.64	1.33	1.40
2	A	701	HEC	C2B-C3B	-6.59	1.33	1.40
2	A	705	HEC	C2B-C3B	-6.32	1.34	1.40
2	A	708	HEC	C2B-C3B	-6.12	1.34	1.40
2	A	706	HEC	C2B-C3B	-6.12	1.34	1.40

The worst 5 of 54 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	706	HEC	CMC-C2C-C1C	-6.45	118.54	128.46
2	A	702	HEC	CMC-C2C-C1C	-5.60	119.86	128.46
2	A	701	HEC	CMC-C2C-C1C	-4.90	120.93	128.46
2	A	710	HEC	CBA-CAA-C2A	-4.72	104.65	112.60
2	A	701	HEC	CBD-CAD-C3D	-4.50	104.94	112.62

There are no chirality outliers.

5 of 24 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	701	HEC	C2A-CAA-CBA-CGA
2	A	704	HEC	C2A-CAA-CBA-CGA
4	A	721	EDO	O1-C1-C2-O2
4	A	719	EDO	O1-C1-C2-O2
2	A	709	HEC	C2A-CAA-CBA-CGA

There are no ring outliers.

13 monomers are involved in 22 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	708	HEC	1	0
5	A	722	ACT	1	0
2	A	710	HEC	1	0
2	A	701	HEC	2	0
2	A	703	HEC	2	0
4	A	718	EDO	1	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	704	HEC	1	0
2	A	707	HEC	2	0
2	A	702	HEC	2	0
4	A	719	EDO	2	0
5	A	723	ACT	3	0
2	A	706	HEC	2	0
2	A	705	HEC	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


























5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	625/679 (92%)	-0.40	2 (0%) 90 89	9, 22, 38, 61	3 (0%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	263	GLU	2.7
1	A	242	ASP	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LBY	A	293	16/17	0.75	0.20	34,57,100,120	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	ACT	A	722	4/4	0.74	0.25	23,36,39,48	0
4	EDO	A	720	4/4	0.76	0.18	42,50,57,61	0
4	EDO	A	721	4/4	0.84	0.15	32,38,39,44	0
4	EDO	A	717	4/4	0.85	0.16	36,39,40,42	0
5	ACT	A	723	4/4	0.85	0.13	23,31,33,35	0
4	EDO	A	718	4/4	0.89	0.10	18,23,26,31	0
4	EDO	A	716	4/4	0.90	0.09	28,30,34,36	0
4	EDO	A	719	4/4	0.92	0.08	30,31,32,37	0
3	CA	A	715	1/1	0.96	0.09	45,45,45,45	0
2	HEC	A	705	43/43	0.97	0.08	15,22,32,38	0
2	HEC	A	706	43/43	0.97	0.07	14,18,31,48	0
3	CA	A	714	1/1	0.97	0.17	33,33,33,33	0
2	HEC	A	701	43/43	0.97	0.07	10,16,39,46	0
2	HEC	A	702	43/43	0.98	0.06	11,16,26,27	0
2	HEC	A	703	43/43	0.98	0.05	10,14,17,19	0
2	HEC	A	707	43/43	0.98	0.06	13,19,23,34	0
2	HEC	A	708	43/43	0.98	0.06	12,15,21,23	0
2	HEC	A	709	43/43	0.98	0.07	12,16,30,44	0
2	HEC	A	710	43/43	0.98	0.06	12,17,22,29	0
3	CA	A	712	1/1	0.98	0.03	25,25,25,25	0
3	CA	A	713	1/1	0.98	0.05	30,30,30,30	0
2	HEC	A	704	43/43	0.98	0.07	11,18,35,43	0
3	CA	A	711	1/1	0.99	0.02	18,18,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around HEC A 705:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 706:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 702:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 703:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 707:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 708:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 709:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEC A 710:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEC A 704:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.