



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 8, 2025 – 03:04 pm GMT

PDB ID : 9QD1 / pdb_00009qdl
Title : Structure of J paramyxovirus virus receptor binding protein
Authors : Stelfox, A.J.; Bowden, T.A.
Deposited on : 2025-03-05
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	435	<div> <div>8%</div> <div>91%</div> <div>9%</div> </div>
1	B	435	<div> <div>5%</div> <div>92%</div> <div>8%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7050 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Attachment glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	435	Total	C	N	O	S	0	0	0
			3382	2150	568	643	21			
1	B	435	Total	C	N	O	S	0	0	0
			3382	2150	568	643	21			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	147	THR	-	expression tag	UNP Q49HN3
A	148	GLY	-	expression tag	UNP Q49HN3
B	147	THR	-	expression tag	UNP Q49HN3
B	148	GLY	-	expression tag	UNP Q49HN3


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	122	Total	O	0	0
			122	122		
2	B	164	Total	O	0	0
			164	164		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Attachment glycoprotein



• Molecule 1: Attachment glycoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	164.24Å 164.24Å 112.47Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.40 – 2.20 46.40 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.40-2.20) 100.0 (46.40-2.20)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 2.20Å)	Xtriage
Refinement program	PHENIX (1.21.1_5286: ???)	Depositor
R, R_{free}	0.189 , 0.207 0.190 , 0.207	Depositor DCC
R_{free} test set	4454 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	42.7	Xtriage
Anisotropy	0.393	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.033 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7050	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.15	0/3476	0.36	0/4744
1	B	0.16	0/3476	0.37	0/4744
All	All	0.15	0/6952	0.37	0/9488

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3382	0	3252	21	0
1	B	3382	0	3250	18	0
2	A	122	0	0	0	0
2	B	164	0	0	2	0
All	All	7050	0	6502	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:350:ARG:NH1	1:B:450:ASP:OD1	2.26	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:278:GLU:O	2:B:601:HOH:O	2.16	0.61
1:A:469:PRO:O	1:A:470:HIS:ND1	2.41	0.54
1:B:429:TRP:CG	1:B:469:PRO:HA	2.43	0.53
1:A:250:MET:HE2	1:A:302:LEU:HD11	1.92	0.52
1:A:521:THR:HG22	1:B:186:PRO:HB3	1.92	0.51
1:A:230:ASP:OD1	1:A:538:LYS:NZ	2.37	0.49
1:A:269:SER:OG	1:A:318:PHE:HA	2.14	0.48
1:B:382:HIS:HD2	2:B:607:HOH:O	1.98	0.47
1:B:504:PRO:HB2	1:B:520:ILE:HD12	1.97	0.47
1:B:572:ASP:OD1	1:B:574:SER:OG	2.33	0.46
1:A:196:SER:HB2	1:A:323:SER:HB2	1.97	0.46
1:A:429:TRP:CG	1:A:469:PRO:HA	2.49	0.46
1:A:174:VAL:HG11	1:A:235:VAL:HG11	1.99	0.45
1:A:191:ARG:HD3	1:A:213:CYS:HB3	1.98	0.45
1:A:496:ASP:HB3	1:A:503:ASN:ND2	2.32	0.44
1:A:461:PRO:HB2	1:A:470:HIS:CD2	2.52	0.44
1:B:196:SER:HB2	1:B:323:SER:HB2	1.99	0.44
1:A:244:TRP:CD2	1:A:298:LYS:HD3	2.53	0.44
1:A:396:ARG:HD3	1:A:442:TRP:CD1	2.52	0.44
1:B:529:TYR:HB2	1:B:548:GLU:CD	2.42	0.43
1:A:419:LYS:NZ	1:A:485:VAL:O	2.51	0.43
1:B:269:SER:OG	1:B:318:PHE:HA	2.17	0.43
1:A:162:ASN:HA	1:A:573:THR:OG1	2.18	0.43
1:B:348:MET:HE1	1:B:448:TRP:HD1	1.83	0.43
1:B:425:ARG:HD3	1:B:427:SER:OG	2.19	0.43
1:A:303:TYR:CE2	1:A:390:LYS:HE2	2.54	0.43
1:A:529:TYR:HB2	1:A:548:GLU:CD	2.44	0.43
1:B:384:LYS:HE2	1:B:384:LYS:HB3	1.67	0.42
1:A:250:MET:CE	1:A:302:LEU:HD11	2.50	0.42
1:B:207:ALA:HB2	1:B:221:TYR:CZ	2.55	0.42
1:B:205:ALA:HA	1:B:222:TYR:O	2.19	0.41
1:A:396:ARG:HD3	1:A:442:TRP:NE1	2.36	0.41
1:A:543:CYS:HB2	1:A:564:TYR:CE2	2.56	0.41
1:B:348:MET:HE1	1:B:448:TRP:CD1	2.56	0.41
1:B:191:ARG:HD3	1:B:213:CYS:HB3	2.04	0.40
1:B:419:LYS:NZ	1:B:485:VAL:O	2.53	0.40
1:A:280:THR:HG22	1:A:372:GLY:HA3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	433/435 (100%)	415 (96%)	18 (4%)	0	100	100
1	B	433/435 (100%)	418 (96%)	15 (4%)	0	100	100
All	All	866/870 (100%)	833 (96%)	33 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	378/378 (100%)	374 (99%)	4 (1%)	70	82
1	B	378/378 (100%)	374 (99%)	4 (1%)	70	82
All	All	756/756 (100%)	748 (99%)	8 (1%)	70	82

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	183	SER
1	A	349	CYS
1	A	453	SER
1	A	487	TYR
1	B	349	CYS
1	B	453	SER
1	B	487	TYR
1	B	500	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	172	ASN
1	A	231	HIS
1	A	341	HIS
1	A	434	GLN
1	B	341	HIS
1	B	388	ASN
1	B	415	HIS
1	B	571	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	435/435 (100%)	0.49	36 (8%)	19 16	31, 51, 82, 113	0
1	B	435/435 (100%)	0.06	23 (5%)	33 30	28, 46, 77, 95	0
All	All	870/870 (100%)	0.27	59 (6%)	25 22	28, 48, 80, 113	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	581	LEU	5.4
1	A	580	PRO	5.0
1	B	581	LEU	4.5
1	A	579	THR	4.3
1	A	513	GLU	4.3
1	A	577	SER	3.9
1	A	578	LEU	3.9
1	A	576	GLY	3.8
1	A	164	LYS	3.8
1	A	349	CYS	3.7
1	B	349	CYS	3.7
1	B	275	PHE	3.6
1	A	364	TYR	3.6
1	A	283	GLU	3.5
1	B	216	SER	3.5
1	A	472	CYS	3.5
1	A	468	CYS	3.4
1	A	575	THR	3.3
1	B	579	THR	3.3
1	B	364	TYR	3.3
1	A	147	THR	3.2
1	B	161	THR	3.2
1	B	213	CYS	3.1
1	B	472	CYS	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	570	CYS	3.1
1	B	147	THR	3.1
1	A	162	ASN	2.9
1	B	274	SER	2.9
1	A	573	THR	2.9
1	B	468	CYS	2.8
1	B	512	GLN	2.8
1	A	574	SER	2.8
1	A	344	ASN	2.8
1	B	580	PRO	2.8
1	B	355	GLN	2.8
1	B	575	THR	2.7
1	A	216	SER	2.7
1	B	356	SER	2.6
1	A	442	TRP	2.6
1	A	169	GLN	2.5
1	A	348	MET	2.5
1	A	512	GLN	2.5
1	A	370	GLU	2.4
1	A	345	VAL	2.4
1	A	159	THR	2.4
1	A	361	LEU	2.4
1	A	358	THR	2.4
1	B	370	GLU	2.4
1	B	578	LEU	2.2
1	A	157	MET	2.2
1	A	347	ALA	2.2
1	A	362	CYS	2.2
1	B	158	CYS	2.2
1	A	572	ASP	2.2
1	B	576	GLY	2.1
1	A	357	PRO	2.1
1	A	471	GLN	2.1
1	B	273	GLY	2.1
1	B	361	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.