



# Full wwPDB NMR Structure Validation Report ⓘ

Oct 28, 2024 – 02:42 pm GMT

PDB ID : 1QLZ  
Title : Human prion protein  
Authors : Zahn, R.; Liu, A.; Luhrs, T.; Wuthrich, K.  
Deposited on : 1999-09-20

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	210492	14027
Ramachandran outliers	207382	12486
Sidechain outliers	206894	12463

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	210	

## 2 Ensemble composition and analysis

This entry contains 20 models. Model 19 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:128-A:228 (101)	0.73	19

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 2 single-model clusters were found.

Cluster number	Models
1	2, 3, 4, 5, 6, 7, 8, 11, 14, 15, 17, 19, 20
2	1, 10, 12, 13, 16
Single-model clusters	9; 18

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1688 atoms, of which 811 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called PRION PROTEIN.

Mol	Chain	Residues	Atoms						Trace
1	A	104	Total	C	H	N	O	S	0
			1688	544	811	153	171	9	


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	21	GLY	-	cloning artifact	UNP P04156
A	22	SER	-	cloning artifact	UNP P04156




### 4.2.2 Score per residue for model 2

- Molecule 1: PRION PROTEIN




### 4.2.3 Score per residue for model 3

- Molecule 1: PRION PROTEIN




#### 4.2.4 Score per residue for model 4

- Molecule 1: PRION PROTEIN




#### 4.2.5 Score per residue for model 5

- Molecule 1: PRION PROTEIN




#### 4.2.6 Score per residue for model 6

- Molecule 1: PRION PROTEIN




#### 4.2.7 Score per residue for model 7


- Molecule 1: PRION PROTEIN



#### 4.2.8 Score per residue for model 8



- Molecule 1: PRION PROTEIN







#### 4.2.9 Score per residue for model 9

- Molecule 1: PRION PROTEIN





#### 4.2.10 Score per residue for model 10

- Molecule 1: PRION PROTEIN




#### 4.2.11 Score per residue for model 11

- Molecule 1: PRION PROTEIN











#### 4.2.12 Score per residue for model 12

- Molecule 1: PRION PROTEIN





#### 4.2.13 Score per residue for model 13

- Molecule 1: PRION PROTEIN



#### 4.2.14 Score per residue for model 14



- Molecule 1: PRION PROTEIN







#### 4.2.15 Score per residue for model 15

- Molecule 1: PRION PROTEIN




#### 4.2.16 Score per residue for model 16

- Molecule 1: PRION PROTEIN




#### 4.2.17 Score per residue for model 17

- Molecule 1: PRION PROTEIN




### 4.2.18 Score per residue for model 18

- Molecule 1: PRION PROTEIN




### 4.2.19 Score per residue for model 19 (medoid)

- Molecule 1: PRION PROTEIN



### 4.2.20 Score per residue for model 20

- Molecule 1: PRION PROTEIN



## 5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *LEAST RESTRAINT VIOLATION*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
OPALp	refinement	
DYANA	structure solution	

No chemical shift data was provided.

## 6 Model quality i

### 6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.69±0.01	0±0/881 ( 0.0± 0.0%)	1.03±0.03	1±1/1189 ( 0.1± 0.1%)
All	All	0.69	0/17620 ( 0.0%)	1.03	22/23780 ( 0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.7±1.3
All	All	0	34

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	151	ARG	NE-CZ-NH2	-8.56	116.02	120.30	8	4
1	A	148	ARG	NE-CZ-NH2	-7.08	116.76	120.30	5	1
1	A	163	TYR	CB-CG-CD2	-6.55	117.07	121.00	3	1
1	A	148	ARG	NE-CZ-NH1	6.47	123.54	120.30	5	1
1	A	128	TYR	CA-CB-CG	6.20	125.18	113.40	1	1
1	A	156	ARG	NE-CZ-NH2	-5.92	117.34	120.30	2	2
1	A	208	ARG	NE-CZ-NH2	-5.56	117.52	120.30	11	1
1	A	148	ARG	CD-NE-CZ	5.56	131.38	123.60	5	1
1	A	144	ASP	CB-CG-OD1	5.54	123.29	118.30	2	1
1	A	164	ARG	NE-CZ-NH2	-5.54	117.53	120.30	11	2
1	A	151	ARG	CD-NE-CZ	5.50	131.31	123.60	8	1
1	A	228	ARG	NE-CZ-NH1	5.30	122.95	120.30	17	1
1	A	136	ARG	NE-CZ-NH1	5.25	122.93	120.30	11	1
1	A	209	VAL	CA-CB-CG1	5.24	118.75	110.90	12	1
1	A	156	ARG	NE-CZ-NH1	5.08	122.84	120.30	2	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	150	TYR	CB-CG-CD1	-5.05	117.97	121.00	4	1
1	A	226	TYR	CB-CG-CD2	-5.01	117.99	121.00	6	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	151	ARG	Sidechain	4
1	A	136	ARG	Sidechain	4
1	A	145	TYR	Sidechain	3
1	A	164	ARG	Sidechain	3
1	A	157	TYR	Sidechain	3
1	A	208	ARG	Sidechain	3
1	A	228	ARG	Sidechain	2
1	A	149	TYR	Sidechain	2
1	A	134	MET	Peptide	2
1	A	148	ARG	Sidechain	2
1	A	169	TYR	Sidechain	2
1	A	163	TYR	Peptide	1
1	A	220	ARG	Sidechain	1
1	A	198	PHE	Sidechain	1
1	A	162	TYR	Sidechain	1

## 6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	861	794	797	2±1
All	All	17220	15880	15940	40

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:130:LEU:HD11	1:A:160:GLN:HB2	0.62	1.72	18	20
1:A:217:GLN:HA	1:A:220:ARG:CG	0.54	2.32	15	1
1:A:200:GLU:HG2	1:A:204:LYS:HE3	0.52	1.82	13	1
1:A:175:PHE:CD2	1:A:218:TYR:CE1	0.48	3.01	9	1
1:A:183:THR:HB	1:A:210:VAL:HG11	0.48	1.84	8	1
1:A:130:LEU:HD12	1:A:161:VAL:O	0.47	2.10	4	4
1:A:161:VAL:HG12	1:A:210:VAL:HG13	0.47	1.85	8	1
1:A:191:THR:HG22	1:A:196:GLU:CG	0.45	2.42	12	1
1:A:129:MET:HE2	1:A:129:MET:H	0.44	1.72	1	1
1:A:217:GLN:HA	1:A:220:ARG:HG3	0.43	1.91	15	1
1:A:150:TYR:CZ	1:A:154:MET:SD	0.43	3.11	12	1
1:A:129:MET:HE2	1:A:129:MET:CA	0.42	2.44	1	1
1:A:200:GLU:CG	1:A:204:LYS:HE3	0.42	2.43	13	1
1:A:172:GLN:CD	1:A:172:GLN:H	0.41	2.18	14	1
1:A:129:MET:HE2	1:A:129:MET:N	0.40	2.30	1	1
1:A:176:VAL:HA	1:A:179:CYS:SG	0.40	2.56	8	1
1:A:137:PRO:HD2	1:A:209:VAL:HG13	0.40	1.94	11	1
1:A:130:LEU:HD21	1:A:160:GLN:NE2	0.40	2.32	16	1

## 6.3 Torsion angles

### 6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	100/210 (48%)	90±3 (90±3%)	9±2 (9±2%)	1±1 (1±1%)	15	64
All	All	2000/4200 (48%)	1793 (90%)	186 (9%)	21 (1%)	15	64

All 11 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	167	ASP	4
1	A	172	GLN	3
1	A	131	GLY	2
1	A	171	ASN	2
1	A	169	TYR	2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	170	SER	2
1	A	154	MET	2
1	A	196	GLU	1
1	A	166	MET	1
1	A	155	HIS	1
1	A	132	SER	1

### 6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	96/159 (60%)	80±3 (83±3%)	16±3 (17±3%)	4	39
All	All	1920/3180 (60%)	1599 (83%)	321 (17%)	4	39

All 57 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	197	ASN	16
1	A	150	TYR	16
1	A	160	GLN	15
1	A	206	MET	14
1	A	163	TYR	14
1	A	208	ARG	14
1	A	135	SER	13
1	A	132	SER	12
1	A	187	HIS	12
1	A	136	ARG	11
1	A	217	GLN	10
1	A	179	CYS	10
1	A	128	TYR	9
1	A	147	ASP	8
1	A	174	ASN	8
1	A	129	MET	7
1	A	186	GLN	7
1	A	227	GLN	7
1	A	148	ARG	6

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	204	LYS	6
1	A	144	ASP	6
1	A	173	ASN	5
1	A	164	ARG	5
1	A	172	GLN	5
1	A	171	ASN	5
1	A	200	GLU	5
1	A	167	ASP	4
1	A	168	GLU	4
1	A	169	TYR	4
1	A	221	GLU	4
1	A	177	HIS	4
1	A	154	MET	4
1	A	155	HIS	4
1	A	214	CYS	4
1	A	175	PHE	3
1	A	159	ASN	3
1	A	228	ARG	3
1	A	170	SER	3
1	A	188	THR	3
1	A	226	TYR	3
1	A	151	ARG	3
1	A	153	ASN	2
1	A	222	SER	2
1	A	156	ARG	2
1	A	220	ARG	2
1	A	191	THR	2
1	A	178	ASP	2
1	A	182	ILE	1
1	A	202	ASP	1
1	A	225	TYR	1
1	A	134	MET	1
1	A	138	ILE	1
1	A	194	LYS	1
1	A	181	ASN	1
1	A	211	GLU	1
1	A	166	MET	1
1	A	196	GLU	1

### 6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided