



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 12, 2024 – 08:47 AM EDT

PDB ID : 1QPN  
Title : Quinolinate Phosphoribosyl Transferase from Mycobacterium Tuberculosis in Complex with NCNN  
Authors : Sharma, V.; Grubmeyer, C.; Sacchettini, J.C.; TB Structural Genomics Consortium (TBSGC)  
Deposited on : 1998-11-20  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtrriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	284	
1	B	284	
1	C	284	
1	D	284	
1	E	284	
1	F	284	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NCN	C	2905	-	-	X	-
2	NCN	F	2906	-	-	X	-

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 12965 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (QUINOLINATE PHOSPHORIBOSYL TRANSFERASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	284	Total	C	N	O	S	0	0	0
			2095	1301	378	411	5			
1	B	284	Total	C	N	O	S	0	0	0
			2095	1301	378	411	5			
1	C	284	Total	C	N	O	S	0	0	0
			2095	1301	378	411	5			
1	D	284	Total	C	N	O	S	0	0	0
			2095	1301	378	411	5			
1	E	284	Total	C	N	O	S	0	0	0
			2095	1301	378	411	5			
1	F	284	Total	C	N	O	S	0	0	0
			2095	1301	378	411	5			

- Molecule 2 is NICOTINATE MONONUCLEOTIDE (three-letter code: NCN) (formula:  $C_{11}H_{14}NO_9P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 22	C 11	N 1	O 9	P 1	0	0
2	B	1	Total 22	C 11	N 1	O 9	P 1	0	0
2	C	1	Total 22	C 11	N 1	O 9	P 1	0	0
2	D	1	Total 22	C 11	N 1	O 9	P 1	0	0
2	E	1	Total 22	C 11	N 1	O 9	P 1	0	0
2	F	1	Total 22	C 11	N 1	O 9	P 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	46	Total 46	O 46	0	0
3	B	48	Total 48	O 48	0	0
3	C	54	Total 54	O 54	0	0
3	D	40	Total 40	O 40	0	0
3	E	45	Total 45	O 45	0	0
3	F	30	Total 30	O 30	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


#### • Molecule 1: PROTEIN (QUINOLINATE PHOSPHORIBOSYL TRANSFERASE)

Chain A: 




#### • Molecule 1: PROTEIN (QUINOLINATE PHOSPHORIBOSYL TRANSFERASE)

Chain B: 




#### • Molecule 1: PROTEIN (QUINOLINATE PHOSPHORIBOSYL TRANSFERASE)

Chain C: 







• Molecule 1: PROTEIN (QUINOLINATE PHOSPHORIBOSYL TRANSFERASE)



• Molecule 1: PROTEIN (QUINOLINATE PHOSPHORIBOSYL TRANSFERASE)



• Molecule 1: PROTEIN (QUINOLINATE PHOSPHORIBOSYL TRANSFERASE)



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	100.38Å 100.38Å 144.61Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	8.00 – 2.60	Depositor
% Data completeness (in resolution range)	96.3 (8.00-2.60)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.185 , 0.269	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	12965	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NCN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.70	0/2120	0.89	2/2891 (0.1%)
1	B	0.71	0/2120	0.91	2/2891 (0.1%)
1	C	0.73	0/2120	0.88	1/2891 (0.0%)
1	D	0.69	0/2120	0.89	2/2891 (0.1%)
1	E	0.70	0/2120	0.89	1/2891 (0.0%)
1	F	0.69	0/2120	0.89	1/2891 (0.0%)
All	All	0.70	0/12720	0.89	9/17346 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	D	1718	LEU	CA-CB-CG	7.92	133.51	115.30
1	F	2718	LEU	CA-CB-CG	7.90	133.47	115.30
1	B	718	LEU	CA-CB-CG	7.38	132.27	115.30
1	C	1218	LEU	CA-CB-CG	6.93	131.23	115.30
1	A	154	ARG	NE-CZ-NH2	-6.89	116.85	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	1525	TYR	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2095	0	2135	81	0
1	B	2095	0	2135	79	0
1	C	2095	0	2135	87	0
1	D	2095	0	2135	70	0
1	E	2095	0	2135	87	0
1	F	2095	0	2135	84	0
2	A	22	0	12	5	0
2	B	22	0	12	6	0
2	C	22	0	12	8	0
2	D	22	0	12	6	0
2	E	22	0	12	6	0
2	F	22	0	12	7	0
3	A	46	0	0	5	0
3	B	48	0	0	2	0
3	C	54	0	0	5	0
3	D	40	0	0	1	0
3	E	45	0	0	3	0
3	F	30	0	0	2	0
All	All	12965	0	12882	465	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 465 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:2903:NCN:H2	3:E:3006:HOH:O	1.57	1.04
1:F:2661:HIS:ND1	2:F:2906:NCN:H4	1.75	1.01
1:D:1748:SER:OG	2:D:2904:NCN:H3'	1.62	0.97
1:A:248:SER:OG	2:A:2901:NCN:H3'	1.65	0.96
1:E:2248:SER:OG	2:E:2903:NCN:H3'	1.66	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/284 (99%)	274 (97%)	6 (2%)	2 (1%)	22	43
1	B	282/284 (99%)	273 (97%)	8 (3%)	1 (0%)	34	57
1	C	282/284 (99%)	272 (96%)	8 (3%)	2 (1%)	22	43
1	D	282/284 (99%)	273 (97%)	8 (3%)	1 (0%)	34	57
1	E	282/284 (99%)	273 (97%)	7 (2%)	2 (1%)	22	43
1	F	282/284 (99%)	273 (97%)	8 (3%)	1 (0%)	34	57
All	All	1692/1704 (99%)	1638 (97%)	45 (3%)	9 (0%)	29	52

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	586	PRO
1	A	86	PRO
1	C	1086	PRO
1	D	1586	PRO
1	F	2586	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	216/216 (100%)	175 (81%)	41 (19%)	1	2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	216/216 (100%)	177 (82%)	39 (18%)	1	2
1	C	216/216 (100%)	176 (82%)	40 (18%)	1	2
1	D	216/216 (100%)	176 (82%)	40 (18%)	1	2
1	E	216/216 (100%)	174 (81%)	42 (19%)	1	2
1	F	216/216 (100%)	180 (83%)	36 (17%)	2	3
All	All	1296/1296 (100%)	1058 (82%)	238 (18%)	1	2

5 of 238 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	1251	LEU
1	F	2631	THR
1	D	1634	LYS
1	F	2610	LEU
1	F	2772	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	1675	HIS
1	E	2160	ASN
1	E	2149	GLN
1	E	2175	HIS
1	B	728	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NCN	A	2901	-	21,23,23	3.64	12 (57%)	27,34,34	2.04	7 (25%)
2	NCN	D	2904	-	21,23,23	3.65	14 (66%)	27,34,34	2.00	7 (25%)
2	NCN	B	2902	-	21,23,23	3.68	12 (57%)	27,34,34	2.07	8 (29%)
2	NCN	F	2906	-	21,23,23	3.47	9 (42%)	27,34,34	2.06	8 (29%)
2	NCN	E	2903	-	21,23,23	3.32	10 (47%)	27,34,34	2.17	8 (29%)
2	NCN	C	2905	-	21,23,23	3.38	11 (52%)	27,34,34	2.08	9 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NCN	A	2901	-	-	7/14/30/30	0/2/2/2
2	NCN	D	2904	-	-	8/14/30/30	0/2/2/2
2	NCN	B	2902	-	-	6/14/30/30	0/2/2/2
2	NCN	F	2906	-	-	8/14/30/30	0/2/2/2
2	NCN	E	2903	-	-	8/14/30/30	0/2/2/2
2	NCN	C	2905	-	-	8/14/30/30	0/2/2/2

The worst 5 of 68 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2904	NCN	O4'-C1'	10.29	1.54	1.40
2	B	2902	NCN	O4'-C1'	10.25	1.54	1.40
2	A	2901	NCN	O4'-C1'	9.57	1.53	1.40
2	F	2906	NCN	O4'-C1'	9.45	1.53	1.40
2	E	2903	NCN	O4'-C1'	9.41	1.53	1.40

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	2903	NCN	C4'-O4'-C1'	-5.76	104.65	109.92
2	E	2903	NCN	C6-N1-C2	-5.26	117.40	121.88
2	B	2902	NCN	C4'-O4'-C1'	-5.14	105.22	109.92
2	D	2904	NCN	C4'-O4'-C1'	-5.00	105.35	109.92
2	A	2901	NCN	C4'-O4'-C1'	-4.85	105.49	109.92

There are no chirality outliers.

5 of 45 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	2905	NCN	C5'-O5'-P-O1P
2	D	2904	NCN	C5'-O5'-P-O1P
2	E	2903	NCN	C5'-O5'-P-O1P
2	F	2906	NCN	C5'-O5'-P-O1P
2	F	2906	NCN	C5'-O5'-P-O3P


There are no ring outliers.

6 monomers are involved in 38 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2901	NCN	5	0
2	D	2904	NCN	6	0
2	B	2902	NCN	6	0
2	F	2906	NCN	7	0
2	E	2903	NCN	6	0
2	C	2905	NCN	8	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


## Ligand NCN A 2901




## Ligand NCN D 2904




## Ligand NCN B 2902



## Ligand NCN F 2906







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.