



wwPDB EM Validation Summary Report ⓘ

Nov 15, 2025 – 06:11 am GMT

PDB ID : 9QTB / pdb_00009qtb
EMDB ID : EMD-53349
Title : Apo form of the L protein from Rift Valley Fever Virus
Authors : Kral, M.; Das, A.R.; Kotacka, T.; Blahosova, A.; Hodek, J.; Konvalinka, J.;
Demo, G.; Kozisek, M.
Deposited on : 2025-04-08
Resolution : 3.50 Å (reported)
Based on initial models : ., 7EEI

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev129
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
EM percentile statistics : 202505.v01 (Using data in the EMD archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY


The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	210492	15764	-
Ramachandran outliers	207382	16835	-
Sidechain outliers	206894	16415	-
Q-score	-	25397	13950 (3.00 - 4.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	2123	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 10289 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-directed RNA polymerase L.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	1283	Total	C	N	O	S	0	0
			10289	6568	1755	1890	76		

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	103	ALA	ASP	variant	UNP A2SZS1
A	2093	GLY	-	expression tag	UNP A2SZS1
A	2094	SER	-	expression tag	UNP A2SZS1
A	2095	ALA	-	expression tag	UNP A2SZS1
A	2096	TRP	-	expression tag	UNP A2SZS1
A	2097	SER	-	expression tag	UNP A2SZS1
A	2098	HIS	-	expression tag	UNP A2SZS1
A	2099	PRO	-	expression tag	UNP A2SZS1
A	2100	GLN	-	expression tag	UNP A2SZS1
A	2101	PHE	-	expression tag	UNP A2SZS1
A	2102	GLU	-	expression tag	UNP A2SZS1
A	2103	LYS	-	expression tag	UNP A2SZS1
A	2104	GLY	-	expression tag	UNP A2SZS1
A	2105	GLY	-	expression tag	UNP A2SZS1
A	2106	GLY	-	expression tag	UNP A2SZS1
A	2107	SER	-	expression tag	UNP A2SZS1
A	2108	GLY	-	expression tag	UNP A2SZS1
A	2109	GLY	-	expression tag	UNP A2SZS1
A	2110	GLY	-	expression tag	UNP A2SZS1
A	2111	SER	-	expression tag	UNP A2SZS1
A	2112	GLY	-	expression tag	UNP A2SZS1
A	2113	GLY	-	expression tag	UNP A2SZS1
A	2114	SER	-	expression tag	UNP A2SZS1
A	2115	ALA	-	expression tag	UNP A2SZS1
A	2116	TRP	-	expression tag	UNP A2SZS1
A	2117	SER	-	expression tag	UNP A2SZS1
A	2118	HIS	-	expression tag	UNP A2SZS1
A	2119	PRO	-	expression tag	UNP A2SZS1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	2120	GLN	-	expression tag	UNP A2SZS1
A	2121	PHE	-	expression tag	UNP A2SZS1
A	2122	GLU	-	expression tag	UNP A2SZS1
A	2123	LYS	-	expression tag	UNP A2SZS1

ARG	THR	ASN	VAL	LYS	GLN	ASP	CYS	GLU	PHE	LEU	R1511
LEU	LEU	LEU	LEU	GLY	ALA	MET	GLN	SER	MET	LEU	T1512
GLN	PHE	GLN	GLN	SER	GLN	ASP	SER	SER	GLY	CYS	T1513
ARG	GLY	MET	PHE	MET	PHE	VAL	PRO	PRO	THR	ILE	K1519
GLU	ALA	SER	PHE	PHE	ARG	ARG	ASP	LEU	VAL	GLN	T1520
LEU	HIS	PHE	GLU	GLU	GLU	LEU	GLY	GLY	VAL	GLY	L1521
ILE	TYR	LEU	VAL	VAL	PRO	ARG	LYS	ILE	PHE	PRO	R1522
ARG	ASP	ASN	ASN	ALA	SER	GLU	PRO	PRO	ILE	SER	D1531
ILE	ILE	VAL	VAL	VAL	ASN	GLU	ILE	ILE	LYS	GLU	T1536
ASP	SER	GLU	SER	GLU	GLU	VAL	ARG	ARG	CYS	SER	R1540
LEU	LEU	GLU	TRP	GLU	VAL	MET	TRP	GLN	GLN	TYR	E1549
GLU	SER	HIS	ARG	ARG	LYS	VAL	ALA	SER	LYS	LEU	GLN
LEU	MET	ASP	CYS	GLY	VAL	ARG	GLU	ASP	VAL	ILE	P1572
TYR	ILE	VAL	VAL	VAL	ALA	GLY	MET	GLY	VAL	GLN	LEU
GLU	GLU	GLU	HIS	TYR	VAL	SER	GLY	GLN	TYR	TYR	ASN
THR	PRO	LEU	ALA	ALA	PRO	THR	VAL	THR	LYS	VAL	ASN
ASP	LEU	ASP	ALA	LEU	LYS	ASN	ASN	ASN	VAL	LEU	ILE
VAL	MET	MET	MET	LEU	LYS	ASN	ASN	ASN	VAL	SER	GLN
ASP	ASP	LYS	ASP	ASP	PHE	LYS	THR	ASP	TYR	VAL	V1579
PHE	PHE	PHE	LEU	LEU	VAL	LEU	GLY	GLY	GLY	SER	R1580
MET	VAL	VAL	MET	ALA	ALA	LYS	TYR	LYS	VAL	ASN	N1581
LEU	ASP	ILE	ILE	ILE	ILE	HIS	SER	SER	GLY	VAL	F1582
GLY	TYR	GLU	GLU	GLU	LEU	HIS	SER	LYS	TRP	ILE	R1591
ALA	ILE	ALA	ALA	ASP	ALA	SER	LYS	ILE	ILE	GLU	P1599
ALA	LYS	SER	LYS	ASN	GLY	GLU	ASP	ARG	GLY	GLU	VAL
MET	GLY	ASN	ASN	ASN	GLY	LEU	GLY	GLY	PHE	SER	LYS
HIS	GLY	ALA	ALA	LYS	LYS	HIS	ALA	ALA	MET	ASP	GLY
PRO	ARG	PHE	PHE	PHE	ARG	ILE	ARG	ILE	ASP	GLY	LYS
GLN	ALA	SER	ALA	SER	GLN	LEU	TYR	GLN	TYR	THR	ARG
PHE	SER	VAL	VAL	ILE	ILE	ILE	SER	TRP	THR	LYS	GLY
GLU	VAL	SER	SER	VAL	LYS	TYR	MET	GLN	VAL	THR	VAL
LYS	ARG	VAL	VAL	VAL	GLY	THR	HIS	VAL	VAL	ASN	SER
GLY	LYS	ASP	ASP	ASP	ILE	ALA	ASP	PHE	HIS	ASN	K1608
GLY	VAL	VAL	VAL	CYS	ASN	SER	SER	SER	GLY	LEU	V1626
GLY	LEU	LEU	ILE	ILE	THR	ASP	ARG	ARG	ILE	ILE	GLY
SER	GLU	GLU	THR	LEU	ARG	ASP	MET	MET	ILE	GLU	GLU
GLY	GLY	ASP	GLY	ASP	LEU	LEU	GLY	GLY	ASN	LEU	ASP
GLY	ARG	VAL	VAL	VAL	SER	SER	PRO	PRO	PHE	GLN	GLN
SER	CYS	SER	SER	GLY	GLU	PRO	LYS	LYS	ILE	ARG	LEU
GLY	SER	SER	ILE	ILE	VAL	GLY	GLY	GLY	PHE	GLY	ALA
GLY	SER	SER	SER	PRO	VAL	ILE	PRO	GLN	LEU	THR	PHE
ALA	LYS	ASP	ASP	TYR	ILE	LYS	PHE	PRO	ASP	GLY	THR
TRP	TYR	ASP	GLU	MET	CYS	SER	CYS	ARG	ASP	GLY	ARG
SER	ASP	GLU	SER	GLU	SER	ILE	PRO	LEU	HIS	VAL	VAL
HIS	LEU	LEU	LEU	SER	GLU	ASP	THR	ASN	VAL	SER	ALA
PRO	SER	PHE	SER	PHE	SER	ASP	TYR	ASN	VAL	VAL	ALA
GLN	LYS	ASP	LYS	ASP	SER	GLU	ILE	ILE	VAL	PRO	GLU
PHE	VAL	VAL	VAL	THR	ARG	GLY	ILE	THR	THR	ILE	ILE
GLU	LEU	SER	LEU	SER	ARG	VAL	ILE	VAL	THR	LEU	LEU
LYS	PHE	LEU	PHE	ASP	VAL	VAL	ILE	THR	ASN	GLU	LYS
THR	ASN	THR	THR	GLY	SER	ALA	GLY	THR	THR	THR	HTS

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	185424	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	1300	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	105000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.633	Depositor
Minimum map value	-0.251	Depositor
Average map value	0.002	Depositor
Map value standard deviation	0.031	Depositor
Recommended contour level	0.12	Depositor
Map size (Å)	213.504, 213.504, 213.504	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.834, 0.834, 0.834	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.16	0/10517	0.34	0/14182

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	10289	0	10284	97	0
All	All	10289	0	10284	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 97 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:309:TYR:OH	1:A:696:PRO:O	1.98	0.80
1:A:250:PHE:O	1:A:813:LYS:NZ	2.21	0.73
1:A:1531:ASP:OD1	1:A:1536:THR:OG1	2.13	0.67
1:A:971:HIS:HE2	1:A:1140:SER:HG	1.42	0.66
1:A:214:LEU:O	1:A:219:ARG:NH1	2.29	0.66

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1267/2123 (60%)	1213 (96%)	53 (4%)	1 (0%)	48 79

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1215	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	1140/1878 (61%)	1140 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1100	HIS
1	A	1247	GLN
1	A	1310	GLN
1	A	978	GLN
1	A	334	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-53349. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 118




Y Index: 116




Z Index: 143


6.3.2 Raw map



X Index: 118



Y Index: 116




Z Index: 143


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.12. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 124 nm^3 ; this corresponds to an approximate mass of 112 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.286 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.286 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.50	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.12	4.66	4.18

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.12 differs from the reported value 3.5 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-53349 and PDB model 9QTB. Per-residue inclusion information can be found in [section 3](#) on [page 5](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.12 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.12).

9.4 Atom inclusion [i](#)



At the recommended contour level, 100% of all backbone atoms, 88% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.12) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div><div></div></div> 0.8850	<div><div></div></div> 0.3800
A	<div><div></div></div> 0.8850	<div><div></div></div> 0.3800

