



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 07:23 AM EDT

PDB ID : 2R7G  
Title : Structure of the retinoblastoma protein pocket domain in complex with adenovirus E1A CR1 domain  
Authors : Liu, X.; Marmorstein, R.  
Deposited on : 2007-09-07  
Resolution : 1.67 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6780 (1.70-1.66)
Clashscore	141614	7310 (1.70-1.66)
Ramachandran outliers	138981	7173 (1.70-1.66)
Sidechain outliers	138945	7172 (1.70-1.66)
RSRZ outliers	127900	6661 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	
1	C	347	
2	B	10	
2	D	10	
2	E	10	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6267 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Retinoblastoma-associated protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	336	Total	C	N	O	S	0	0	0
			2760	1779	463	498	20			
1	C	337	Total	C	N	O	S	0	0	0
			2771	1785	467	499	20			

- Molecule 2 is a protein called Early E1A 32 kDa protein.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	10	Total	C	N	O	0	0	0
			84	56	12	16			
2	D	10	Total	C	N	O	0	0	0
			84	56	12	16			
2	E	9	Total	C	N	O	0	0	0
			76	50	11	15			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	232	Total	O	0	0
			232	232		
4	B	7	Total	O	0	0
			7	7		
4	C	234	Total	O	0	0
			234	234		
4	D	7	Total	O	0	0
			7	7		
4	E	2	Total	O	0	0
			2	2		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Retinoblastoma-associated protein



- Molecule 1: Retinoblastoma-associated protein



- Molecule 2: Early E1A 32 kDa protein




- Molecule 2: Early E1A 32 kDa protein

Chain D:  100%

There are no outlier residues recorded for this chain.

- Molecule 2: Early E1A 32 kDa protein

Chain E:  20% 60% 30% 10%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.71Å 57.71Å 92.27Å 94.34° 92.69° 105.79°	Depositor
Resolution (Å)	40.82 – 1.67 40.84 – 1.67	Depositor EDS
% Data completeness (in resolution range)	95.2 (40.82-1.67) 87.5 (40.84-1.67)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.07 (at 1.67Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.195 , 0.218 0.194 , 0.218	Depositor DCC
$R_{free}$ test set	12140 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.9	Xtriage
Anisotropy	0.090	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 50.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.012 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6267	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.57	0/2814	0.63	0/3790
1	C	0.57	0/2825	0.63	0/3804
2	B	0.50	0/87	0.76	0/119
2	D	0.54	0/87	0.62	0/119
2	E	0.38	0/79	0.59	0/108
All	All	0.56	0/5892	0.63	0/7940

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2760	0	2814	49	0
1	C	2771	0	2827	37	0
2	B	84	0	80	4	0
2	D	84	0	80	0	0
2	E	76	0	69	4	0
3	A	5	0	0	0	0
3	C	5	0	0	0	0
4	A	232	0	0	6	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	7	0	0	0	0
4	C	234	0	0	4	0
4	D	7	0	0	0	0
4	E	2	0	0	0	0
All	All	6267	0	5870	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:525:ALA:HB1	1:A:557:ILE:HD12	1.42	1.00
1:A:514:PHE:HZ	1:A:557:ILE:HD11	1.33	0.93
1:C:486:LEU:HD23	1:C:531:VAL:HG11	1.52	0.91
1:C:422:ILE:HD11	1:C:484:MET:SD	2.12	0.88
1:A:555:HIS:HA	1:A:558:MET:HE3	1.56	0.86
1:C:528:PHE:CE2	1:C:532:ILE:HD11	2.13	0.84
1:C:663:ASN:O	1:C:667:GLU:HG2	1.78	0.83
1:A:753:ILE:HD12	2:E:48:ASP:HB3	1.61	0.81
1:A:528:PHE:CE2	1:A:532:ILE:HD11	2.17	0.79
1:A:676:LEU:HD11	1:A:717:ILE:HD13	1.63	0.78
1:A:493:VAL:O	1:A:497:THR:HG23	1.85	0.77
1:A:555:HIS:HA	1:A:558:MET:CE	2.16	0.75
1:A:645:THR:HG23	2:B:45:GLU:O	1.87	0.74
1:A:425:ILE:HG13	1:C:425:ILE:HD11	1.69	0.74
1:A:514:PHE:CZ	1:A:557:ILE:HD11	2.20	0.74
1:C:555:HIS:HA	1:C:558:MET:HE2	1.70	0.72
1:C:555:HIS:HA	1:C:558:MET:CE	2.21	0.71
1:C:493:VAL:O	1:C:497:THR:HG23	1.91	0.70
1:A:557:ILE:HD13	1:A:561:LEU:HD12	1.73	0.70
1:A:714:VAL:HG13	2:E:43:LEU:HD21	1.73	0.69
1:C:422:ILE:CD1	1:C:484:MET:SD	2.83	0.67
1:C:428:GLU:HG2	4:C:955:HOH:O	1.95	0.66
1:A:537:LYS:HD2	4:A:952:HOH:O	1.96	0.65
1:A:453:TYR:HD1	1:A:486:LEU:HD22	1.62	0.65
1:C:678:HIS:HE1	1:C:780:SER:O	1.80	0.64
1:A:753:ILE:CD1	2:E:48:ASP:HB3	2.27	0.64
1:A:652:LYS:HE2	4:A:915:HOH:O	1.99	0.62
1:A:678:HIS:HE1	1:A:780:SER:O	1.83	0.62
1:C:529:TYR:OH	1:C:653:LYS:HD2	2.03	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:497:THR:HG22	4:A:1117:HOH:O	2.03	0.57
1:A:425:ILE:CG1	1:C:425:ILE:HD11	2.35	0.56
1:C:652:LYS:HE3	1:C:656:ARG:HH22	1.69	0.56
1:A:685:GLN:HE22	1:A:689:GLN:HE21	1.52	0.56
1:A:453:TYR:CD1	1:A:486:LEU:HD22	2.40	0.56
2:B:47:TYR:O	2:B:49:LEU:HD13	2.06	0.54
1:C:685:GLN:HE22	1:C:689:GLN:NE2	2.05	0.54
1:C:709:TYR:CE1	1:C:713:LYS:HE2	2.44	0.53
1:A:380:ASN:HB2	4:A:1021:HOH:O	2.09	0.52
1:A:456:VAL:HG13	1:A:538:ALA:HB3	1.90	0.52
1:A:643:LYS:NZ	4:A:964:HOH:O	2.43	0.52
1:A:685:GLN:HE22	1:A:689:GLN:NE2	2.09	0.51
1:A:486:LEU:HD23	1:A:486:LEU:O	2.10	0.51
1:C:531:VAL:HG12	1:C:531:VAL:O	2.09	0.51
1:A:488:ALA:HB2	1:A:521:LEU:HD23	1.93	0.50
1:A:691:GLU:HG3	1:A:764:LEU:HD21	1.92	0.50
1:C:497:THR:HG22	4:C:1105:HOH:O	2.13	0.48
1:C:486:LEU:HD23	1:C:531:VAL:CG1	2.33	0.48
1:A:744:ILE:HD12	1:A:744:ILE:N	2.29	0.47
1:A:553:CYS:O	1:A:557:ILE:HG12	2.14	0.47
1:C:691:GLU:HG3	1:C:764:LEU:HD21	1.97	0.46
1:A:652:LYS:HE3	1:A:656:ARG:HH12	1.80	0.46
1:A:529:TYR:HB2	1:A:557:ILE:HG21	1.98	0.46
1:C:500:ARG:HG2	1:C:501:SER:N	2.32	0.45
1:C:652:LYS:CE	1:C:656:ARG:HH22	2.27	0.45
1:A:744:ILE:HG22	1:A:745:LYS:HD3	1.98	0.45
1:C:425:ILE:HD12	1:C:425:ILE:N	2.31	0.45
1:A:548:LYS:HE3	1:A:548:LYS:HB2	1.65	0.44
1:A:517:ILE:HG13	1:A:521:LEU:HD22	1.99	0.44
1:A:475:LYS:NZ	1:C:575:GLN:HE21	2.14	0.44
1:C:558:MET:HE3	1:C:657:LEU:HD22	2.00	0.44
1:C:558:MET:HE3	1:C:558:MET:HB2	1.66	0.44
1:A:519:ASN:HB3	4:A:959:HOH:O	2.18	0.44
1:C:456:VAL:HG13	1:C:538:ALA:HB3	1.99	0.44
1:C:555:HIS:HA	1:C:558:MET:HE3	1.99	0.43
1:C:731:LEU:HB2	1:C:734:ALA:HB2	2.00	0.43
1:A:500:ARG:HG2	1:A:501:SER:N	2.34	0.43
1:A:555:HIS:ND1	1:A:558:MET:CE	2.81	0.43
1:C:557:ILE:HA	1:C:561:LEU:HB2	2.00	0.43
1:A:761:MET:HG2	1:A:765:LYS:HB3	1.99	0.43
1:A:531:VAL:HG12	1:A:531:VAL:O	2.19	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:484:MET:HE3	1:C:521:LEU:HG	2.01	0.42
1:C:784:HIS:O	1:C:785:ILE:HG13	2.19	0.42
1:C:489:CYS:SG	1:C:532:ILE:HD13	2.59	0.42
1:A:709:TYR:CE1	1:A:713:LYS:HG3	2.54	0.42
1:C:484:MET:CE	1:C:521:LEU:HG	2.49	0.42
1:C:721:PHE:O	1:C:725:VAL:HB	2.19	0.42
1:C:439:VAL:HA	4:C:958:HOH:O	2.20	0.42
1:C:726:THR:O	1:C:729:LYS:HG2	2.20	0.41
1:A:774:THR:HG23	1:A:775:ARG:HD2	2.02	0.41
1:A:662:LEU:HD22	1:A:666:CYS:SG	2.60	0.41
1:A:725:VAL:HG13	1:A:739:PHE:CZ	2.56	0.41
1:A:761:MET:HE1	2:E:44:HIS:HA	2.02	0.40
1:A:475:LYS:HE3	4:C:931:HOH:O	2.19	0.40
1:A:476:LEU:HD21	2:B:43:LEU:HD23	2.03	0.40
2:B:47:TYR:HB2	2:B:49:LEU:HD22	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	330/347 (95%)	322 (98%)	7 (2%)	1 (0%)	41	23
1	C	331/347 (95%)	324 (98%)	4 (1%)	3 (1%)	17	4
2	B	8/10 (80%)	8 (100%)	0	0	100	100
2	D	8/10 (80%)	8 (100%)	0	0	100	100
2	E	7/10 (70%)	7 (100%)	0	0	100	100
All	All	684/724 (94%)	669 (98%)	11 (2%)	4 (1%)	25	10

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	501	SER
1	C	718	ASP
1	C	501	SER
1	C	719	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/325 (96%)	293 (94%)	20 (6%)	17	4
1	C	314/325 (97%)	296 (94%)	18 (6%)	20	5
2	B	10/10 (100%)	9 (90%)	1 (10%)	7	1
2	D	10/10 (100%)	10 (100%)	0	100	100
2	E	9/10 (90%)	9 (100%)	0	100	100
All	All	656/680 (96%)	617 (94%)	39 (6%)	19	5

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	380	ASN
1	A	385	LEU
1	A	487	LEU
1	A	500	ARG
1	A	502	THR
1	A	521	LEU
1	A	552	ARG
1	A	555	HIS
1	A	569	LEU
1	A	662	LEU
1	A	668	ARG
1	A	669	LEU
1	A	683	LEU
1	A	694	LEU
1	A	713	LYS
1	A	714	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	725	VAL
1	A	745	LYS
1	A	775	ARG
1	A	778	THR
2	B	49	LEU
1	C	385	LEU
1	C	486	LEU
1	C	487	LEU
1	C	500	ARG
1	C	521	LEU
1	C	552	ARG
1	C	555	HIS
1	C	569	LEU
1	C	652	LYS
1	C	662	LEU
1	C	669	LEU
1	C	672	GLU
1	C	678	HIS
1	C	683	LEU
1	C	694	LEU
1	C	714	VAL
1	C	721	PHE
1	C	745	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	380	ASN
1	A	390	ASN
1	A	399	ASN
1	A	406	ASN
1	A	471	GLN
1	A	678	HIS
1	A	689	GLN
1	A	690	ASN
1	A	716	ASN
1	A	736	GLN
1	A	767	ASN
1	A	770	GLN
1	C	399	ASN
1	C	406	ASN
1	C	471	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	575	GLN
1	C	678	HIS
1	C	689	GLN
1	C	690	ASN
1	C	762	GLN
1	C	770	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	C	901	-	4,4,4	0.56	0	6,6,6	0.42	0
3	SO4	A	902	-	4,4,4	0.22	0	6,6,6	0.46	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	336/347 (96%)	0.55	34 (10%) <b>7</b> <b>7</b>	18, 25, 42, 54	0
1	C	337/347 (97%)	0.65	37 (10%) <b>5</b> <b>5</b>	19, 25, 44, 53	0
2	B	10/10 (100%)	0.39	0 <b>100</b> <b>100</b>	23, 26, 28, 31	0
2	D	10/10 (100%)	0.30	0 <b>100</b> <b>100</b>	23, 25, 28, 29	0
2	E	9/10 (90%)	1.13	2 (22%) <b>0</b> <b>0</b>	37, 39, 42, 42	0
All	All	702/724 (96%)	0.60	73 (10%) <b>6</b> <b>6</b>	18, 25, 42, 54	0

All (73) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	502	THR	10.6
1	A	502	THR	9.1
1	C	501	SER	9.0
1	A	501	SER	9.0
1	A	775	ARG	8.8
1	A	774	THR	8.8
1	C	775	ARG	7.5
1	C	774	THR	6.7
1	C	509	GLY	6.3
1	C	785	ILE	6.0
1	C	773	SER	5.9
1	A	784	HIS	5.8
1	A	509	GLY	5.4
1	C	716	ASN	5.0
1	A	777	PRO	4.7
1	A	785	ILE	4.4
1	A	510	THR	4.4
1	A	672	GLU	4.3
1	C	777	PRO	4.3
1	C	718	ASP	4.3

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	671	SER	4.2
1	C	671	SER	4.2
1	A	732	PRO	4.0
1	C	778	THR	4.0
1	C	579	ARG	3.9
1	C	776	PRO	3.9
1	C	643	LYS	3.7
1	A	747	GLU	3.6
1	A	778	THR	3.6
1	C	668	ARG	3.6
1	A	776	PRO	3.5
2	E	48	ASP	3.5
1	C	674	PRO	3.3
1	C	746	GLU	3.2
1	C	510	THR	3.2
1	A	668	ARG	3.1
1	C	673	HIS	3.1
1	C	784	HIS	3.1
1	C	439	VAL	3.0
1	C	672	GLU	3.0
1	C	531	VAL	2.7
1	A	667	GLU	2.7
1	C	730	ASP	2.7
1	A	439	VAL	2.6
1	C	736	GLN	2.6
1	C	732	PRO	2.6
1	C	382	ILE	2.5
1	C	523	LEU	2.5
1	C	436	GLN	2.5
1	C	528	PHE	2.5
1	A	424	TYR	2.4
1	A	762	GLN	2.4
1	A	745	LYS	2.4
1	C	424	TYR	2.3
1	A	544	ARG	2.3
1	A	523	LEU	2.3
1	C	544	ARG	2.3
1	A	487	LEU	2.3
1	A	673	HIS	2.3
1	C	667	GLU	2.3
1	A	674	PRO	2.2
1	A	531	VAL	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	733	HIS	2.2
1	A	486	LEU	2.2
1	A	500	ARG	2.2
1	A	528	PHE	2.2
1	A	746	GLU	2.2
1	A	481	ILE	2.1
1	C	767	ASN	2.1
2	E	47	TYR	2.1
1	C	779	LEU	2.1
1	A	643	LYS	2.0
1	C	722	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SO4	C	901	5/5	0.98	0.08	20,22,23,24	0
3	SO4	A	902	5/5	0.99	0.09	20,21,22,22	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.