



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 22, 2024 – 05:56 AM EDT

PDB ID : 3RAG
Title : Crystal Structure of Uncharacterized protein Aaci_0196 from Alicyclobacillus acidocaldarius subsp. acidocaldarius DSM 446
Authors : Wu, R.; Bedearn, J.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2011-03-28
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7108 (1.80-1.80)
Clashscore	180529	8162 (1.80-1.80)
Ramachandran outliers	177936	8077 (1.80-1.80)
Sidechain outliers	177891	8076 (1.80-1.80)
RSRZ outliers	164620	7108 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	242	
1	B	242	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	241	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3709 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called uncharacterized protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	232	Total	C	N	O	S	Se	0	2	0
			1736	1065	320	339	3	9			
1	B	226	Total	C	N	O	S	Se	0	2	0
			1688	1042	309	325	3	9			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP C8WQT1
A	-1	ASN	-	expression tag	UNP C8WQT1
A	0	ALA	-	expression tag	UNP C8WQT1
B	-2	SER	-	expression tag	UNP C8WQT1
B	-1	ASN	-	expression tag	UNP C8WQT1
B	0	ALA	-	expression tag	UNP C8WQT1

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		
2	B	1	Total	Na	0	0
			1	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	160	Total	O	0	0
			160	160		
4	B	111	Total	O	0	0
			111	111		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: uncharacterized protein



- Molecule 1: uncharacterized protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	46.70Å 85.89Å 132.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.95 – 1.80 42.95 – 1.80	Depositor EDS
% Data completeness (in resolution range)	93.9 (42.95-1.80) 93.6 (42.95-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.99 (at 1.79Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.193 , 0.215 0.242 , 0.260	Depositor DCC
R_{free} test set	2526 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	30.3	Xtriage
Anisotropy	0.522	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 48.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3709	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	0/1747	0.60	0/2356
1	B	0.52	0/1697	0.56	0/2288
All	All	0.54	0/3444	0.58	0/4644

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1736	0	1754	61	1
1	B	1688	0	1721	43	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	12	0	16	26	0
4	A	160	0	0	3	0
4	B	111	0	0	4	0
All	All	3709	0	3491	104	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (104) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:45:GLY:N	3:A:241:GOL:H2	1.57	1.18
1:A:45:GLY:H	3:A:241:GOL:H2	0.89	1.02
1:B:182:GLU:HG2	1:B:183:ALA:H	1.22	1.01
1:A:45:GLY:H	3:A:241:GOL:C2	1.79	0.94
1:A:209:THR:HG21	4:A:256:HOH:O	1.66	0.93
1:B:67:MSE:HE2	1:B:83:MSE:HG2	1.55	0.89
1:A:52:GLN:NE2	3:A:241:GOL:H32	1.90	0.87
1:A:67:MSE:HE2	1:A:83:MSE:HG2	1.58	0.86
1:A:52:GLN:HE21	3:A:241:GOL:H32	1.41	0.84
1:B:182:GLU:HG2	1:B:183:ALA:N	1.93	0.83
1:A:52:GLN:CB	3:A:241:GOL:H11	2.09	0.82
1:A:52:GLN:HG2	3:A:241:GOL:H11	1.61	0.81
1:B:39:ASN:HD22	1:B:67:MSE:H	1.29	0.80
1:B:205:ALA:O	1:B:206:ARG:HG3	1.84	0.78
1:A:101:LEU:O	1:A:104:MSE:O	2.02	0.78
1:A:188:GLN:HE22	1:A:195:ASN:H	1.31	0.77
1:A:39:ASN:HD22	1:A:67:MSE:H	1.34	0.76
1:A:52:GLN:CG	3:A:241:GOL:H11	2.17	0.74
1:A:52:GLN:HB3	3:A:241:GOL:H11	1.68	0.74
1:A:44:VAL:HA	3:A:241:GOL:H12	1.70	0.74
1:A:52:GLN:HE21	3:A:241:GOL:C3	2.01	0.73
1:B:5:ALA:O	1:B:130[A]:VAL:HG21	1.93	0.69
1:A:188:GLN:NE2	1:A:195:ASN:H	1.89	0.69
1:A:18:SER:HA	1:A:209:THR:OG1	1.93	0.68
1:B:39:ASN:ND2	1:B:67:MSE:H	1.93	0.64
1:A:67:MSE:CE	1:A:83:MSE:HG2	2.26	0.64
1:A:44:VAL:HA	3:A:241:GOL:C1	2.28	0.62
1:B:39:ASN:HD21	1:B:87:THR:HG21	1.64	0.62
1:B:148:ILE:HA	1:B:151[A]:VAL:HG12	1.80	0.62
1:A:209:THR:HG23	4:A:263:HOH:O	1.99	0.61
1:A:58:HIS:HD2	3:A:242:GOL:O2	1.83	0.61
1:A:39:ASN:HD21	1:A:87:THR:HG21	1.65	0.61
1:B:148:ILE:HA	1:B:151[A]:VAL:CG1	2.33	0.59
1:A:49:ALA:CB	3:A:241:GOL:H31	2.33	0.58
1:A:52:GLN:HG2	3:A:241:GOL:C1	2.33	0.58
1:A:52:GLN:CG	3:A:241:GOL:H32	2.33	0.58
1:B:99:GLU:O	1:B:103:VAL:HG22	2.03	0.58
1:A:39:ASN:ND2	1:A:67:MSE:H	1.99	0.58
1:B:130[B]:VAL:HG12	1:B:131:ALA:O	2.04	0.58
1:A:101:LEU:C	1:A:104:MSE:O	2.43	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:179:LYS:HG2	1:A:185:ARG:CZ	2.35	0.56
1:A:185:ARG:HD3	1:A:202:GLU:OE1	2.05	0.55
1:A:44:VAL:HA	3:A:241:GOL:C2	2.36	0.54
1:A:141:SER:HB3	1:A:209:THR:HG22	1.89	0.54
1:B:145:ARG:HB3	1:B:205:ALA:HB3	1.88	0.54
1:B:86:GLN:HG2	1:B:90:MSE:HE2	1.89	0.54
1:A:44:VAL:HG13	3:A:241:GOL:O2	2.08	0.54
1:B:142:ALA:O	1:B:145:ARG:HG2	2.07	0.54
1:B:16:GLY:HA2	1:B:49:ALA:HB2	1.90	0.53
1:A:196:VAL:HG13	4:A:301:HOH:O	2.08	0.53
1:B:162:LEU:O	1:B:165:ARG:HG2	2.08	0.53
1:A:179:LYS:HG2	1:A:185:ARG:NH2	2.25	0.52
1:A:96:VAL:HG11	1:A:119:VAL:HG13	1.92	0.51
1:A:52:GLN:HG2	3:A:241:GOL:H32	1.91	0.51
1:B:5:ALA:O	1:B:130[B]:VAL:HG11	2.11	0.51
1:A:179:LYS:HE3	1:A:185:ARG:HG3	1.93	0.50
1:B:227:ARG:CZ	4:B:383:HOH:O	2.60	0.50
1:A:89:GLN:HG3	1:A:127[A]:GLU:OE1	2.11	0.49
1:A:15:ASP:O	1:A:52:GLN:HG3	2.12	0.49
1:B:140:THR:O	1:B:205:ALA:HA	2.12	0.49
1:A:101:LEU:HA	1:A:104:MSE:O	2.13	0.48
1:B:5:ALA:C	1:B:130[A]:VAL:HG21	2.32	0.48
1:B:230:ASP:HB2	4:B:258:HOH:O	2.13	0.48
1:B:182:GLU:CG	1:B:183:ALA:N	2.62	0.48
1:B:72:GLN:HG2	4:B:354:HOH:O	2.13	0.47
1:B:93:GLN:HG3	1:B:120:MSE:HE1	1.96	0.47
1:A:106:LYS:HB3	1:A:110:ASP:HB2	1.95	0.47
1:B:2:THR:O	1:B:130[A]:VAL:HG23	2.14	0.47
1:B:6:THR:HA	1:B:130[B]:VAL:CG1	2.45	0.47
1:B:148:ILE:O	1:B:152:ARG:HG2	2.14	0.47
1:B:72:GLN:HG2	1:B:72:GLN:H	1.45	0.47
1:B:6:THR:HA	1:B:130[A]:VAL:HG22	1.97	0.47
1:A:52:GLN:CD	3:A:241:GOL:H32	2.34	0.46
1:A:44:VAL:HA	3:A:241:GOL:H2	1.96	0.46
1:A:44:VAL:C	3:A:241:GOL:H2	2.30	0.46
1:B:150:THR:O	1:B:153:GLU:HG2	2.15	0.46
1:A:90:MSE:O	1:A:94:GLN:HG3	2.15	0.46
1:B:6:THR:HA	1:B:130[B]:VAL:HG13	1.98	0.46
1:A:188:GLN:HE22	1:A:194:VAL:HA	1.81	0.46
1:A:188:GLN:HE22	1:A:195:ASN:N	2.07	0.45
1:B:67:MSE:HE1	1:B:82:MSE:SE	2.66	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:39:ASN:HB3	1:A:83:MSE:HE3	1.99	0.45
1:A:1:MSE:HB2	1:A:1:MSE:HE3	1.34	0.45
1:A:49:ALA:HB2	3:A:241:GOL:H31	1.99	0.44
1:A:179:LYS:HE2	1:A:185:ARG:HH21	1.81	0.44
1:B:94:GLN:NE2	1:B:98:GLN:OE1	2.50	0.44
1:B:52:GLN:HG3	1:B:55:GLN:H	1.82	0.44
1:A:185:ARG:HD2	1:A:202:GLU:HB3	2.00	0.44
1:B:67:MSE:CE	1:B:83:MSE:HG2	2.38	0.44
1:A:44:VAL:CA	3:A:241:GOL:H2	2.48	0.44
1:A:45:GLY:O	1:A:46:ARG:HD2	2.17	0.43
1:B:7:ILE:HG13	1:B:88:MSE:HG3	2.00	0.43
1:A:68:CYS:H	3:A:242:GOL:C3	2.32	0.42
1:B:199:LEU:HD23	1:B:199:LEU:HA	1.85	0.42
1:B:1:MSE:HE3	1:B:1:MSE:HB2	1.84	0.42
1:B:156:ARG:HD2	1:B:196:VAL:CG1	2.50	0.42
1:A:86:GLN:HG2	1:A:90:MSE:HE2	2.01	0.41
1:A:140:THR:O	1:A:205:ALA:HA	2.21	0.41
1:B:177:PRO:HA	1:B:183:ALA:O	2.20	0.41
1:B:177:PRO:HD3	1:B:208:GLY:O	2.19	0.41
1:B:30:ARG:HD3	4:B:416:HOH:O	2.21	0.41
1:A:52:GLN:NE2	3:A:241:GOL:C3	2.66	0.41
1:A:152:ARG:HG2	1:A:203:LEU:HD12	2.03	0.41
1:A:101:LEU:CA	1:A:104:MSE:O	2.68	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:109:GLU:OE1	1:B:185:ARG:NE[3_546]	2.06	0.14

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	232/242 (96%)	230 (99%)	1 (0%)	1 (0%)	30	19
1	B	222/242 (92%)	219 (99%)	3 (1%)	0	100	100
All	All	454/484 (94%)	449 (99%)	4 (1%)	1 (0%)	44	31

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	210	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	186/183 (102%)	182 (98%)	4 (2%)	47	36
1	B	182/183 (100%)	177 (97%)	5 (3%)	40	28
All	All	368/366 (100%)	359 (98%)	9 (2%)	47	32

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MSE
1	A	55[A]	GLN
1	A	55[B]	GLN
1	A	88	MSE
1	B	72	GLN
1	B	103	VAL
1	B	104	MSE
1	B	145	ARG
1	B	147	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	52	GLN
1	A	58	HIS
1	A	188	GLN
1	B	39	ASN
1	B	89	GLN
1	B	93	GLN
1	B	94	GLN
1	B	98	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	241	-	5,5,5	0.43	0	5,5,5	0.37	0
3	GOL	A	242	-	5,5,5	0.35	0	5,5,5	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	241	-	-	2/4/4/4	-
3	GOL	A	242	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	242	GOL	O1-C1-C2-C3
3	A	241	GOL	O1-C1-C2-C3
3	A	242	GOL	C1-C2-C3-O3
3	A	242	GOL	O1-C1-C2-O2
3	A	241	GOL	O1-C1-C2-O2
3	A	242	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 26 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	241	GOL	24	0
3	A	242	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	223/242 (92%)	0.81	31 (13%) 7 6	13, 33, 61, 90	2 (0%)
1	B	217/242 (89%)	1.03	23 (10%) 13 10	17, 41, 72, 92	2 (0%)
All	All	440/484 (90%)	0.92	54 (12%) 9 8	13, 38, 67, 92	4 (0%)

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	182	GLU	9.8
1	A	229	VAL	8.0
1	B	49	ALA	6.3
1	B	0	ALA	5.9
1	B	2	THR	4.9
1	A	55[A]	GLN	4.5
1	A	109	GLU	4.4
1	A	105	GLY	4.3
1	B	3	ARG	4.3
1	B	161	SER	4.2
1	B	201	ALA	3.8
1	B	229	VAL	3.5
1	A	0	ALA	3.5
1	B	74	ALA	3.4
1	A	102	ALA	3.2
1	B	230	ASP	3.2
1	A	230	ASP	3.1
1	A	224	SER	3.1
1	A	103	VAL	3.1
1	A	51	GLU	3.0
1	A	118	ARG	3.0
1	A	209	THR	3.0
1	A	3	ARG	2.8
1	A	74	ALA	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	179	LYS	2.7
1	B	103	VAL	2.6
1	A	196	VAL	2.6
1	A	45	GLY	2.6
1	B	178	GLY	2.6
1	A	100	LEU	2.5
1	B	35	GLY	2.5
1	A	85	HIS	2.5
1	B	145	ARG	2.5
1	B	109	GLU	2.5
1	A	228	ASN	2.4
1	B	143	SER	2.4
1	B	204	VAL	2.3
1	A	93	GLN	2.3
1	A	113	PRO	2.2
1	B	50	GLY	2.2
1	A	94	GLN	2.2
1	A	121	GLN	2.2
1	B	51	GLU	2.2
1	A	-1	ASN	2.2
1	B	52	GLN	2.2
1	B	4	GLU	2.1
1	A	226	ALA	2.1
1	A	205	ALA	2.1
1	A	107	SER	2.1
1	A	52	GLN	2.0
1	B	206	ARG	2.0
1	B	151[A]	VAL	2.0
1	A	101	LEU	2.0
1	A	161	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	A	241	6/6	0.81	0.25	35,50,56,68	6
3	GOL	A	242	6/6	0.86	0.13	37,52,65,67	0
2	NA	B	240	1/1	0.93	0.26	50,50,50,50	0
2	NA	A	240	1/1	0.97	0.10	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.