



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 15, 2024 – 06:18 PM EDT

PDB ID : 4S1T
Title : Crystal structure of the mutant I26A/N52A of the endoribonuclease from human coronavirus 229E
Authors : Huo, T.; Liu, X.
Deposited on : 2015-01-15
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	<div> <div>9%</div> <div>67%</div> <div>25%</div> <div>6%</div> </div>
1	B	349	<div> <div>11%</div> <div>73%</div> <div>21%</div> <div>• •</div> </div>
1	C	349	<div> <div>•</div> <div>72%</div> <div>22%</div> <div>• •</div> </div>
1	D	349	<div> <div>4%</div> <div>69%</div> <div>23%</div> <div>5% • •</div> </div>
1	E	349	<div> <div>2%</div> <div>70%</div> <div>25%</div> <div>• •</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	349	<div><div><div>%</div><div><div></div></div><div>74%</div><div>19%</div><div><div></div><div></div></div></div></div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 16623 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uridylate-specific endoribonuclease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	342	Total	C	N	O	S	0	0	0
			2679	1728	433	505	13			
1	B	342	Total	C	N	O	S	0	0	0
			2679	1728	433	505	13			
1	C	342	Total	C	N	O	S	0	0	0
			2679	1728	433	505	13			
1	D	342	Total	C	N	O	S	0	0	0
			2679	1728	433	505	13			
1	E	342	Total	C	N	O	S	0	0	0
			2679	1728	433	505	13			
1	F	342	Total	C	N	O	S	0	0	0
			2679	1728	433	505	13			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP P0C6X1
A	26	ALA	ILE	engineered mutation	UNP P0C6X1
A	52	ALA	ASN	engineered mutation	UNP P0C6X1
B	0	SER	-	expression tag	UNP P0C6X1
B	26	ALA	ILE	engineered mutation	UNP P0C6X1
B	52	ALA	ASN	engineered mutation	UNP P0C6X1
C	0	SER	-	expression tag	UNP P0C6X1
C	26	ALA	ILE	engineered mutation	UNP P0C6X1
C	52	ALA	ASN	engineered mutation	UNP P0C6X1
D	0	SER	-	expression tag	UNP P0C6X1
D	26	ALA	ILE	engineered mutation	UNP P0C6X1
D	52	ALA	ASN	engineered mutation	UNP P0C6X1
E	0	SER	-	expression tag	UNP P0C6X1
E	26	ALA	ILE	engineered mutation	UNP P0C6X1
E	52	ALA	ASN	engineered mutation	UNP P0C6X1
F	0	SER	-	expression tag	UNP P0C6X1
F	26	ALA	ILE	engineered mutation	UNP P0C6X1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	52	ALA	ASN	engineered mutation	UNP P0C6X1

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		
2	E	1	Total	O	P	0	0
			5	4	1		
2	F	1	Total	O	P	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	63	Total	O	0	0
			63	63		
3	B	49	Total	O	0	0
			49	49		
3	C	102	Total	O	0	0
			102	102		


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	109	Total 109	O 109	0	0
3	E	102	Total 102	O 102	0	0
3	F	94	Total 94	O 94	0	0




• Molecule 1: Uridylate-specific endoribonuclease




• Molecule 1: Uridylate-specific endoribonuclease



• Molecule 1: Uridylate-specific endoribonuclease





4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	86.44Å 140.28Å 426.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.22 – 2.50 49.95 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.7 (43.22-2.50) 96.3 (49.95-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.77 (at 2.51Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, R_{free}	0.241 , 0.285 0.238 , 0.279	Depositor DCC
R_{free} test set	4466 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	37.4	Xtriage
Anisotropy	0.656	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 41.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	16623	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.87	0/2737	0.77	2/3716 (0.1%)
1	B	0.88	0/2737	0.75	1/3716 (0.0%)
1	C	0.80	0/2737	0.79	5/3716 (0.1%)
1	D	0.79	0/2737	0.79	5/3716 (0.1%)
1	E	0.81	1/2737 (0.0%)	0.79	3/3716 (0.1%)
1	F	0.78	0/2737	0.77	4/3716 (0.1%)
All	All	0.82	1/16422 (0.0%)	0.78	20/22296 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2
1	C	0	2
1	D	0	2
1	E	0	1
1	F	0	3
All	All	0	10

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	88	TYR	CD2-CE2	5.41	1.47	1.39

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	137	ALA	C-N-CA	10.82	148.75	121.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	92	ARG	NE-CZ-NH1	-7.87	116.37	120.30
1	D	137	ALA	O-C-N	-7.29	111.03	122.70
1	E	24	VAL	CB-CA-C	-6.85	98.39	111.40
1	F	92	ARG	NE-CZ-NH1	-6.83	116.89	120.30

There are no chirality outliers.

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	141	LYS	Peptide
1	B	146	SER	Peptide
1	C	145	LYS	Peptide
1	C	91	GLU	Peptide
1	D	91	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2679	0	2666	132	0
1	B	2679	0	2667	61	0
1	C	2679	0	2667	70	0
1	D	2679	0	2667	109	1
1	E	2679	0	2667	75	1
1	F	2679	0	2667	70	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	1	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
3	A	63	0	0	15	1
3	B	49	0	0	15	1
3	C	102	0	0	7	0
3	D	109	0	0	7	0
3	E	102	0	0	7	0
3	F	94	0	0	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	16623	0	16001	499	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 499 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:140:VAL:CG1	1:A:147:LEU:HD22	1.20	1.57
1:A:140:VAL:HG11	1:A:147:LEU:CD2	1.34	1.50
1:A:143:GLY:C	1:A:145:LYS:HB3	1.33	1.43
1:D:149:ALA:HB2	1:D:180:TYR:CD1	1.61	1.32
1:A:142:THR:C	1:A:145:LYS:HB2	1.61	1.19

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:82:LYS:NZ	1:E:312:ASP:OD2[1_455]	1.87	0.33
3:A:509:HOH:O	3:B:520:HOH:O[4_555]	1.93	0.27

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	338/349 (97%)	317 (94%)	14 (4%)	7 (2%)	7 11
1	B	338/349 (97%)	319 (94%)	17 (5%)	2 (1%)	25 43
1	C	338/349 (97%)	321 (95%)	14 (4%)	3 (1%)	17 31
1	D	338/349 (97%)	318 (94%)	17 (5%)	3 (1%)	17 31
1	E	338/349 (97%)	323 (96%)	13 (4%)	2 (1%)	25 43

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	338/349 (97%)	320 (95%)	16 (5%)	2 (1%)	25	43
All	All	2028/2094 (97%)	1918 (95%)	91 (4%)	19 (1%)	17	31

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	143	GLY
1	A	147	LEU
1	A	148	PRO
1	D	145	LYS
1	D	148	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/304 (98%)	273 (92%)	25 (8%)	11	21
1	B	298/304 (98%)	273 (92%)	25 (8%)	11	21
1	C	298/304 (98%)	272 (91%)	26 (9%)	10	20
1	D	298/304 (98%)	271 (91%)	27 (9%)	9	18
1	E	298/304 (98%)	273 (92%)	25 (8%)	11	21
1	F	298/304 (98%)	279 (94%)	19 (6%)	17	33
All	All	1788/1824 (98%)	1641 (92%)	147 (8%)	11	22

5 of 147 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	165	VAL
1	F	244	THR
1	E	233	PHE
1	F	47	THR
1	B	320	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	320	HIS
1	F	153	ASN
1	F	216	ASN
1	C	153	ASN
1	C	11	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	E	401	-	4,4,4	0.79	0	6,6,6	0.97	0
2	PO4	B	401	-	4,4,4	0.99	0	6,6,6	1.66	2 (33%)
2	PO4	D	401	-	4,4,4	1.30	0	6,6,6	1.13	1 (16%)
2	PO4	F	401	-	4,4,4	1.87	1 (25%)	6,6,6	1.17	1 (16%)
2	PO4	C	401	-	4,4,4	1.40	1 (25%)	6,6,6	1.25	1 (16%)
2	PO4	A	401	-	4,4,4	0.92	0	6,6,6	0.98	0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	401	PO4	P-O1	3.55	1.59	1.50
2	C	401	PO4	P-O1	2.68	1.57	1.50

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	PO4	O3-P-O1	-2.59	101.40	110.89
2	B	401	PO4	O3-P-O2	2.24	115.15	107.97
2	C	401	PO4	O4-P-O2	-2.21	100.86	107.97
2	F	401	PO4	O3-P-O2	-2.05	101.38	107.97
2	D	401	PO4	O4-P-O2	2.05	114.56	107.97

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	401	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	342/349 (97%)	0.53	31 (9%)	9 9	30, 46, 82, 102	0
1	B	342/349 (97%)	0.65	40 (11%)	4 4	28, 48, 84, 130	0
1	C	342/349 (97%)	0.18	5 (1%)	73 75	28, 43, 80, 101	0
1	D	342/349 (97%)	0.36	14 (4%)	37 40	27, 44, 79, 113	0
1	E	342/349 (97%)	0.19	7 (2%)	65 68	28, 45, 80, 112	0
1	F	342/349 (97%)	0.08	4 (1%)	79 80	28, 45, 81, 109	0
All	All	2052/2094 (97%)	0.33	101 (4%)	29 31	27, 45, 81, 130	0

The worst 5 of 101 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	228	LEU	8.9
1	A	144	GLY	8.7
1	D	144	GLY	8.2
1	D	146	SER	7.0
1	D	142	THR	5.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	F	401	5/5	0.84	0.15	45,47,57,87	0
2	PO4	B	401	5/5	0.87	0.15	56,57,72,107	0
2	PO4	D	401	5/5	0.90	0.12	39,39,63,75	0
2	PO4	A	401	5/5	0.92	0.15	60,63,80,99	0
2	PO4	C	401	5/5	0.92	0.09	42,45,63,86	0
2	PO4	E	401	5/5	0.94	0.12	51,65,75,85	0

6.5 Other polymers [i](#)

There are no such residues in this entry.