



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 18, 2025 – 10:09 pm BST

PDB ID : 9S3A / pdb\_00009s3a  
Title : TaGST-10 in complex with deoxynivalenol-13-glutathione  
Authors : Michlmayr, H.; Papageorgiou, A.C.  
Deposited on : 2025-07-24  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.006 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.45.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	5963 (2.30-2.30)
Clashscore	180529	6698 (2.30-2.30)
Ramachandran outliers	177936	6640 (2.30-2.30)
Sidechain outliers	177891	6640 (2.30-2.30)
RSRZ outliers	164620	5963 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	236	 5% 76% 13% 10%
1	B	236	 6% 78% 10% 12%

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 3600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glutathione S-transferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	212	Total	C	N	O	S	0	0	0
			1587	1022	257	299	9			
1	B	207	Total	C	N	O	S	0	0	0
			1544	996	251	291	6			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP A0A3B5XZG4
A	0	HIS	-	expression tag	UNP A0A3B5XZG4
B	-1	GLY	-	expression tag	UNP A0A3B5XZG4
B	0	HIS	-	expression tag	UNP A0A3B5XZG4

- Molecule 2 is Deoxynivalenol-13-glutathione (CCD ID: A114U) (formula:  $C_{25}H_{37}N_3O_{12}S$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			41	25	3	12	1		
2	B	1	Total	C	N	O	S	0	0
			41	25	3	12	1		

- Molecule 3 is Deoxynivalenol-13-cysteine (CCD ID: A1I4T) (formula:  $C_{18}H_{27}NO_8S$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			28	18	1	8	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is ACETATE ION (CCD ID: ACT) (formula:  $C_2H_3O_2$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	6	Total	Na	0	0
			6	6		
6	B	2	Total	Na	0	0
			2	2		

- Molecule 7 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C<sub>4</sub>H<sub>12</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			8	4	1	3		
7	B	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	179	Total	O	0	0
			179	179		
8	B	138	Total	O	0	0
			138	138		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutathione S-transferase



- Molecule 1: Glutathione S-transferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.41 Å   61.86 Å   168.82 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	49.90 – 2.30 49.90 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.9 (49.90-2.30) 98.9 (49.90-2.30)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.14 (at 2.29 Å)	Xtriage
Refinement program	REFMAC 5.8.0430	Depositor
R, $R_{free}$	0.210   ,   0.258 0.216   ,   0.264	Depositor DCC
$R_{free}$ test set	1281 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.3	Xtriage
Anisotropy	0.986	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 55.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3600	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.40% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, TRS, NA, A1I4T, A1I4U, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.90	1/1620 (0.1%)	1.64	15/2197 (0.7%)
1	B	0.78	0/1577	1.54	9/2143 (0.4%)
All	All	0.84	1/3197 (0.0%)	1.59	24/4340 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	97	ARG	C-O	-5.42	1.17	1.24

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	187	ASP	CA-CB-CG	6.96	119.56	112.60
1	A	95	HIS	CA-CB-CG	-6.75	107.05	113.80
1	B	180	MET	CG-SD-CE	6.30	114.75	100.90
1	B	179	LYS	CB-CA-C	6.26	121.50	110.85
1	A	171	LEU	CA-C-O	-6.20	113.98	120.55
1	A	132	LYS	N-CA-CB	6.16	118.94	110.01
1	A	7	ASP	CA-CB-CG	5.79	118.39	112.60
1	B	211	MET	CG-SD-CE	5.76	113.57	100.90
1	A	66	LYS	CB-CA-C	5.75	116.77	110.15
1	A	172	ALA	CA-C-N	5.69	127.91	120.28

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	172	ALA	C-N-CA	5.69	127.91	120.28
1	A	163	VAL	N-CA-CB	5.65	117.16	110.55
1	B	40	ASP	CA-CB-CG	5.51	118.11	112.60
1	B	187	ASP	CA-CB-CG	5.50	118.10	112.60
1	A	180	MET	CG-SD-CE	5.46	112.91	100.90
1	A	211	MET	CA-C-O	-5.42	115.31	119.32
1	B	218	VAL	N-CA-CB	5.41	116.88	110.55
1	B	7	ASP	CA-CB-CG	5.35	117.95	112.60
1	B	95	HIS	CA-CB-CG	-5.31	108.49	113.80
1	A	199	ASP	CA-CB-CG	5.15	117.75	112.60
1	A	188	PRO	CB-CA-C	-5.11	103.74	112.64
1	A	193	LEU	N-CA-CB	5.07	117.58	110.12
1	B	139	GLU	CB-CG-CD	5.04	121.17	112.60
1	A	180	MET	CB-CA-C	5.00	119.35	110.85

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	97	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1587	0	1594	15	0
1	B	1544	0	1542	9	0
2	A	41	0	0	0	0
2	B	41	0	0	1	0
3	A	28	0	0	0	0
4	A	6	0	8	0	0
5	A	4	0	3	0	0
5	B	8	0	6	0	0
6	A	6	0	0	0	0
6	B	2	0	0	0	0
7	A	8	0	11	0	0
7	B	8	0	12	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	A	179	0	0	13	1
8	B	138	0	0	9	0
All	All	3600	0	3176	25	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:173:TRP:O	8:A:601:HOH:O	1.83	0.96
2:B:501:A1I4U:O7	8:B:601:HOH:O	1.84	0.95
1:B:52:ASN:O	8:B:602:HOH:O	1.91	0.88
1:A:176:ALA:N	8:A:601:HOH:O	2.15	0.80
1:A:173:TRP:C	8:A:601:HOH:O	2.23	0.79
1:A:165:VAL:O	8:A:602:HOH:O	2.07	0.73
1:A:38:GLU:N	8:A:604:HOH:O	2.22	0.72
1:B:56:GLN:N	8:B:602:HOH:O	2.22	0.71
1:A:83:ALA:N	8:A:603:HOH:O	2.27	0.67
1:B:55:LEU:N	8:B:602:HOH:O	2.33	0.61
1:B:58:VAL:HG22	8:B:601:HOH:O	2.05	0.55
1:A:138:VAL:HG12	1:A:191:THR:HG21	1.91	0.53
1:A:79:ASP:O	8:A:603:HOH:O	2.19	0.52
1:B:53:PRO:C	8:B:602:HOH:O	2.54	0.50
1:B:58:VAL:N	8:B:601:HOH:O	2.25	0.48
1:B:172:ALA:HB3	8:B:712:HOH:O	2.12	0.48
1:A:101:ARG:NH1	8:A:610:HOH:O	2.47	0.48
1:A:22:ARG:HA	8:A:602:HOH:O	2.14	0.47
1:A:82:PHE:N	8:A:603:HOH:O	2.47	0.47
1:A:80:GLU:C	8:A:603:HOH:O	2.59	0.46
1:A:21:LEU:HD13	8:A:612:HOH:O	2.14	0.46
1:A:104:ALA:HB1	1:A:166:MET:HE2	1.99	0.43
1:B:138:VAL:HG12	1:B:191:THR:HG21	2.00	0.42
1:B:77:TYR:HB2	8:B:603:HOH:O	2.20	0.41
1:A:25:LEU:HD12	8:A:602:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:607:HOH:O	8:A:637:HOH:O[3_554]	2.00	0.20

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	208/236 (88%)	206 (99%)	2 (1%)	0	100	100
1	B	203/236 (86%)	200 (98%)	2 (1%)	1 (0%)	25	32
All	All	411/472 (87%)	406 (99%)	4 (1%)	1 (0%)	44	55

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	70	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/179 (93%)	163 (98%)	3 (2%)	54	71
1	B	160/179 (89%)	156 (98%)	4 (2%)	42	60
All	All	326/358 (91%)	319 (98%)	7 (2%)	48	66

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	GLU
1	A	54	VAL
1	A	79	ASP
1	B	8	GLU
1	B	50	LYS
1	B	79	ASP
1	B	135	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	GLN
1	B	24	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 8 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	A	503	-	5,5,5	0.16	0	5,5,5	0.38	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	ACT	A	504	-	3,3,3	1.35	0	3,3,3	0.68	0
7	TRS	A	511	6	7,7,7	0.37	0	9,9,9	0.76	0
2	A1I4U	B	501	-	39,44,44	1.09	3 (7%)	38,70,70	1.45	6 (15%)
5	ACT	B	502	-	3,3,3	1.06	0	3,3,3	0.80	0
2	A1I4U	A	501	-	39,44,44	1.07	2 (5%)	38,70,70	1.37	5 (13%)
3	A1I4T	A	502	-	26,31,31	1.29	4 (15%)	20,53,53	1.99	7 (35%)
7	TRS	B	506	-	7,7,7	0.50	0	9,9,9	0.91	0
5	ACT	B	503	-	3,3,3	1.15	0	3,3,3	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	503	-	-	2/4/4/4	-
7	TRS	A	511	6	-	6/9/9/9	-
2	A1I4U	B	501	-	-	1/26/101/101	0/6/4/4
2	A1I4U	A	501	-	-	0/26/101/101	0/6/4/4
3	A1I4T	A	502	-	-	4/8/83/83	0/6/4/4
7	TRS	B	506	-	-	3/9/9/9	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	A1I4T	C15-C4	3.28	1.59	1.54
2	B	501	A1I4U	O1-C4	3.22	1.49	1.45
3	A	502	A1I4T	C17-C18	3.17	1.57	1.54
2	A	501	A1I4U	O1-C4	2.87	1.48	1.45
2	A	501	A1I4U	C22-C4	2.51	1.58	1.54
3	A	502	A1I4T	O1-C4	2.36	1.48	1.45
2	B	501	A1I4U	O5-C18	-2.32	1.22	1.30
2	B	501	A1I4U	C24-C25	2.19	1.56	1.54
3	A	502	A1I4T	C10-C8	2.05	1.60	1.56

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	A1I4U	O3-C10-C11	-4.34	99.39	109.61
2	A	501	A1I4U	O3-C10-C11	-4.01	100.17	109.61

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	A1I4T	O3-C10-C11	-3.78	100.69	109.61
3	A	502	A1I4T	O1-C4-C15	3.33	116.46	112.50
2	A	501	A1I4U	C22-C4-C3	3.14	116.71	113.49
2	A	501	A1I4U	C24-C2-C3	-3.06	116.48	121.14
3	A	502	A1I4T	C13-C12-S1	3.04	124.87	113.74
2	B	501	A1I4U	C24-C2-C3	-2.97	116.62	121.14
3	A	502	A1I4T	C12-C13-C14	2.92	119.54	110.77
3	A	502	A1I4T	C15-C4-C3	2.91	116.47	113.49
3	A	502	A1I4T	C4-C3-C2	-2.79	120.98	124.28
2	B	501	A1I4U	O1-C4-C3	2.75	110.06	105.54
2	A	501	A1I4U	O2-C6-C7	2.48	117.37	111.36
3	A	502	A1I4T	O2-C6-C7	2.45	117.28	111.36
2	A	501	A1I4U	C12-S1-C11	2.43	109.14	102.09
2	B	501	A1I4U	C16-C15-C14	2.32	118.22	113.04
2	B	501	A1I4U	C12-S1-C11	2.17	108.38	102.09
2	B	501	A1I4U	C13-C19-N3	2.11	120.79	116.54

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	A1I4T	S1-C12-C13-C14
3	A	502	A1I4T	S1-C12-C13-N1
7	A	511	TRS	C1-C-C2-O2
7	A	511	TRS	C3-C-C2-O2
7	A	511	TRS	N-C-C2-O2
7	B	506	TRS	C1-C-C3-O3
7	B	506	TRS	C2-C-C3-O3
7	B	506	TRS	N-C-C3-O3
4	A	503	GOL	O1-C1-C2-C3
7	A	511	TRS	C3-C-C1-O1
3	A	502	A1I4T	N1-C13-C14-O4
3	A	502	A1I4T	N1-C13-C14-O5
4	A	503	GOL	O1-C1-C2-O2
7	A	511	TRS	N-C-C1-O1
7	A	511	TRS	C2-C-C1-O1
2	B	501	A1I4U	N2-C17-C18-O5

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	A1I4U	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




## Ligand A1I4U B 501



## Ligand A1I4U A 501





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	212/236 (89%)	0.27	12 (5%) 30 32	22, 33, 67, 95	0
1	B	207/236 (87%)	0.57	14 (6%) 25 26	25, 40, 79, 123	0
All	All	419/472 (88%)	0.42	26 (6%) 28 29	22, 36, 76, 123	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	128	ALA	5.0
1	B	180	MET	3.9
1	B	130	GLY	3.7
1	A	224	MET	3.7
1	A	120	MET	3.6
1	B	131	LYS	3.5
1	B	128	ALA	3.5
1	A	130	GLY	3.5
1	B	183	VAL	3.5
1	B	218	VAL	3.3
1	B	129	GLU	3.2
1	A	127	LYS	3.1
1	B	118	ALA	2.8
1	B	127	LYS	2.8
1	A	118	ALA	2.7
1	A	119	SER	2.6
1	B	64	ASP	2.6
1	B	221	ALA	2.6
1	A	117	GLN	2.6
1	B	150	ASN	2.4
1	B	179	LYS	2.3
1	A	116	SER	2.2
1	B	182	GLY	2.0
1	A	131	LYS	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	216	LYS	2.0
1	A	147	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	NA	A	509	1/1	0.67	0.21	67,67,67,67	0
6	NA	A	508	1/1	0.70	0.24	53,53,53,53	0
5	ACT	A	504	4/4	0.77	0.12	46,46,50,51	0
5	ACT	B	503	4/4	0.77	0.19	59,66,67,67	0
5	ACT	B	502	4/4	0.79	0.20	45,52,57,72	0
7	TRS	A	511	8/8	0.79	0.18	44,55,63,66	0
6	NA	A	507	1/1	0.80	0.18	57,57,57,57	0
4	GOL	A	503	6/6	0.80	0.19	65,67,72,72	0
6	NA	A	510	1/1	0.81	0.16	48,48,48,48	0
7	TRS	B	506	8/8	0.82	0.16	34,44,46,59	0
6	NA	A	506	1/1	0.84	0.27	53,53,53,53	0
3	A1I4T	A	502	28/28	0.86	0.12	37,42,50,51	0
6	NA	B	504	1/1	0.86	0.33	54,54,54,54	0
2	A1I4U	B	501	41/41	0.88	0.12	37,50,58,61	0
2	A1I4U	A	501	41/41	0.89	0.12	38,50,58,61	0
6	NA	B	505	1/1	0.91	0.10	50,50,50,50	0
6	NA	A	505	1/1	0.92	0.11	47,47,47,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around A1I4T A 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around A1I4U B 501:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around A1I4U A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.