



wwPDB EM Validation Summary Report ⓘ

Feb 20, 2025 – 06:09 pm GMT

PDB ID : 8S7V
EMDB ID : EMD-19787
Title : Methyl-coenzyme M reductase activation complex binding to the A2 component
Authors : Ramirez-Amador, F.; Paul, S.; Kumar, A.; Schuller, J.M.
Deposited on : 2024-03-04
Resolution : 2.56 Å(reported)
Based on initial model : .

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : **FAILED**
Mogul : 1.8.4, CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : **FAILED**
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.41


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	260	
1	B	260	
2	D	443	
2	E	443	
3	C	553	
3	F	553	
4	G	183	
5	H	304	
6	I	234	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
7	K	531	
8	J	501	
9	L	93	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GL3	C	448	-	-	X	-

2 Entry composition [i](#)

There are 17 unique types of molecules in this entry. The entry contains 32524 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methyl-coenzyme M reductase subunit gamma.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	259	Total	C	N	O	S	0	0
			2071	1293	369	397	12		
1	B	259	Total	C	N	O	S	0	0
			2071	1293	369	397	12		

- Molecule 2 is a protein called Methyl-coenzyme M reductase subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	E	442	Total	C	N	O	S	0	0
			3263	2066	546	630	21		
2	D	442	Total	C	N	O	S	0	0
			3263	2066	546	630	21		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	173	GLY	SER	conflict	UNP A0A2L1CBB3
D	173	GLY	SER	conflict	UNP A0A2L1CBB3

- Molecule 3 is a protein called Methyl-coenzyme M reductase subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	C	550	Total	C	N	O	S	0	0
			4284	2707	731	826	20		
3	F	520	Total	C	N	O	S	0	0
			4039	2556	684	779	20		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	51	SER	ALA	variant	UNP A0A2L1CBB0
F	51	SER	ALA	variant	UNP A0A2L1CBB0

- Molecule 4 is a protein called Methanogenesis marker protein 17.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	G	114	931	600	152	172	7	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	109	VAL	ILE	variant	UNP G0H411
G	129	ILE	VAL	variant	UNP G0H411
G	167	GLU	GLN	variant	UNP G0H411
G	168	GLU	ASP	variant	UNP G0H411
G	171	ASN	ASP	variant	UNP G0H411

- Molecule 5 is a protein called Methanogenesis marker protein 7.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	H	296	2393	1530	411	440	12	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	115	ASN	SER	variant	UNP G0H350
H	260	GLU	LYS	variant	UNP G0H350

- Molecule 6 is a protein called Methyl-coenzyme M reductase operon protein C.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	I	194	1469	929	261	271	8	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	-35	MET	-	initiating methionine	UNP G0H3B1
I	-34	SER	-	expression tag	UNP G0H3B1
I	-33	ALA	-	expression tag	UNP G0H3B1
I	-32	TRP	-	expression tag	UNP G0H3B1
I	-31	SER	-	expression tag	UNP G0H3B1
I	-30	HIS	-	expression tag	UNP G0H3B1
I	-29	PRO	-	expression tag	UNP G0H3B1
I	-28	GLN	-	expression tag	UNP G0H3B1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
I	-27	PHE	-	expression tag	UNP G0H3B1
I	-26	GLU	-	expression tag	UNP G0H3B1
I	-25	LYS	-	expression tag	UNP G0H3B1
I	-24	GLY	-	expression tag	UNP G0H3B1
I	-23	GLY	-	expression tag	UNP G0H3B1
I	-22	GLY	-	expression tag	UNP G0H3B1
I	-21	SER	-	expression tag	UNP G0H3B1
I	-20	GLY	-	expression tag	UNP G0H3B1
I	-19	GLY	-	expression tag	UNP G0H3B1
I	-18	GLY	-	expression tag	UNP G0H3B1
I	-17	SER	-	expression tag	UNP G0H3B1
I	-16	GLY	-	expression tag	UNP G0H3B1
I	-15	GLY	-	expression tag	UNP G0H3B1
I	-14	SER	-	expression tag	UNP G0H3B1
I	-13	ALA	-	expression tag	UNP G0H3B1
I	-12	TRP	-	expression tag	UNP G0H3B1
I	-11	SER	-	expression tag	UNP G0H3B1
I	-10	HIS	-	expression tag	UNP G0H3B1
I	-9	PRO	-	expression tag	UNP G0H3B1
I	-8	GLN	-	expression tag	UNP G0H3B1
I	-7	PHE	-	expression tag	UNP G0H3B1
I	-6	GLU	-	expression tag	UNP G0H3B1
I	-5	LYS	-	expression tag	UNP G0H3B1
I	-4	SER	-	expression tag	UNP G0H3B1
I	-3	ALA	-	expression tag	UNP G0H3B1
I	-2	GLY	-	expression tag	UNP G0H3B1
I	-1	SER	-	expression tag	UNP G0H3B1
I	0	GLY	-	expression tag	UNP G0H3B1

- Molecule 7 is a protein called Glycine betaine/carnitine/choline transport ATP-binding protein OpuCA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	K	521	4089	2598	693	774	24	0	0

- Molecule 8 is a protein called UPF0288 protein MmarC6_0796.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	J	499	3962	2517	647	790	8	0	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
J	500	SER	LYS	variant	UNP A9A8E0

- Molecule 9 is a protein called DUF2098 domain-containing protein.


Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
9	L	47	390	251	61	78	0	0

- Molecule 10 is O-PHOSPHONO-N-{(2E)-7-[(2-SULFOETHYL)DITHIO]HEPT-2-ENOYL}-L-THREONINE (three-letter code: SHT) (formula: C₁₃H₂₄NO₁₀PS₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf	
			Total	C	N	O	P		S
10	E	1	28	13	1	10	1	3	0

- Molecule 11 is Coenzyme B (three-letter code: TP7) (formula: C₁₁H₂₂NO₇PS).




Mol	Chain	Residues	Atoms					AltConf	
			Total	C	N	O	P		S
11	C	1	21	11	1	7	1	1	0

- Molecule 12 is 1-THIOETHANESULFONIC ACID (three-letter code: COM) (formula: $C_2H_6O_3S_2$).




Mol	Chain	Residues	Atoms			AltConf	
			Total	C	O		S
12	C	1	7	2	3	2	0

- Molecule 13 is FACTOR 430 (three-letter code: F43) (formula: $C_{42}H_{51}N_6NiO_{13}$).



Mol	Chain	Residues	Atoms				AltConf	
13	F	1	Total	C	N	Ni	O	0
			62	42	6	1	13	
13	F	1	Total	C	N	Ni	O	0
			62	42	6	1	13	

- Molecule 14 is FeFe cofactor (three-letter code: S5Q) (formula: CFe_8S_9) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf	
14	H	1	Total	C	Fe	S	0
			18	1	8	9	

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				AltConf
14	I	1	Total	C	Fe	S	0
			18	1	8	9	
14	I	1	Total	C	Fe	S	0
			18	1	8	9	

- Molecule 15 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
15	K	1	Total	Zn	0
			1	1	

- Molecule 16 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					AltConf
16	K	1	Total	C	N	O	P	0
			31	10	5	13	3	
16	K	1	Total	C	N	O	P	0
			31	10	5	13	3	

- Molecule 17 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
17	K	2	Total	Mg	0
			2	2	


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Methyl-coenzyme M reductase subunit gamma




- Molecule 1: Methyl-coenzyme M reductase subunit gamma




- Molecule 2: Methyl-coenzyme M reductase subunit beta




- Molecule 2: Methyl-coenzyme M reductase subunit beta






• Molecule 3: Methyl-coenzyme M reductase subunit alpha




• Molecule 3: Methyl-coenzyme M reductase subunit alpha




• Molecule 4: Methanogenesis marker protein 17







• Molecule 5: Methanogenesis marker protein 7



• Molecule 6: Methyl-coenzyme M reductase operon protein C




• Molecule 7: Glycine betaine/carnitine/choline transport ATP-binding protein OpuCA





● Molecule 8: UPF0288 protein MmarC6_0796



● Molecule 9: DUF2098 domain-containing protein



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	484016	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	60	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	165000	Depositor
Image detector	FEI FALCON IV (4k x 4k)	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, F43, MG, COM, ATP, MGN, S5Q, SMC, GL3, SHT, AGM, MHS, TP7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/2116	0.57	3/2861 (0.1%)
1	B	0.28	0/2116	0.52	0/2861
2	D	0.28	0/3319	0.49	2/4499 (0.0%)
2	E	0.28	0/3319	0.58	5/4499 (0.1%)
3	C	0.29	0/4328	0.47	0/5856
3	F	0.27	0/4078	0.49	1/5522 (0.0%)
4	G	0.25	0/943	0.48	0/1262
5	H	0.27	0/2435	0.52	0/3280
6	I	0.28	0/1490	0.63	3/2006 (0.1%)
7	K	0.31	0/4159	0.73	17/5613 (0.3%)
8	J	0.41	0/4026	0.74	6/5441 (0.1%)
9	L	0.24	0/398	0.51	0/542
All	All	0.30	0/32727	0.58	37/44242 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	1
3	F	0	1
7	K	0	2
8	J	0	6
All	All	0	10

There are no bond length outliers.

The worst 5 of 37 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	J	32	ILE	O-C-N	21.22	156.66	122.70
7	K	88	GLU	CB-CA-C	-18.60	73.20	110.40
8	J	32	ILE	CA-C-N	-18.01	77.57	117.20
2	E	82	ALA	CB-CA-C	-16.20	85.80	110.10
8	J	32	ILE	C-N-CA	-13.52	87.91	121.70

There are no chirality outliers.

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	447	TYR	Mainchain
3	F	274	ARG	Sidechain
8	J	42	GLU	Mainchain
7	K	143	GLY	Peptide
7	K	88	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2071	0	2012	39	0
1	B	2071	0	2012	27	0
2	D	3263	0	3260	59	0
2	E	3263	0	3260	53	0
3	C	4284	0	4147	103	0
3	F	4039	0	3904	57	0
4	G	931	0	970	18	0
5	H	2393	0	2475	55	0
6	I	1469	0	1534	33	0
7	K	4089	0	4162	100	0
8	J	3962	0	3996	186	0
9	L	390	0	378	5	0
10	E	28	0	21	2	0
11	C	21	0	19	1	0
12	C	7	0	5	1	0
13	F	124	0	86	15	0
14	H	18	0	0	1	0
14	I	36	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
15	K	1	0	0	0	0
16	K	62	0	24	2	0
17	K	2	0	0	0	0
All	All	32524	0	32265	664	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 664 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:J:179:TYR:CE1	8:J:295:PHE:HB3	1.36	1.56
3:C:447:TYR:C	3:C:448:GL3:CA	1.86	1.43
3:C:447:TYR:CA	3:C:448:GL3:N	1.84	1.39
8:J:179:TYR:CD1	8:J:295:PHE:HB3	1.61	1.34
8:J:179:TYR:CE1	8:J:295:PHE:CB	2.16	1.28

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/260 (99%)	252 (98%)	5 (2%)	0	100	100
1	B	257/260 (99%)	252 (98%)	5 (2%)	0	100	100
2	D	440/443 (99%)	422 (96%)	18 (4%)	0	100	100
2	E	440/443 (99%)	418 (95%)	21 (5%)	1 (0%)	44	54
3	C	541/553 (98%)	511 (94%)	30 (6%)	0	100	100
3	F	511/553 (92%)	488 (96%)	22 (4%)	1 (0%)	44	54
4	G	110/183 (60%)	106 (96%)	4 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	H	294/304 (97%)	288 (98%)	6 (2%)	0	100	100
6	I	192/234 (82%)	186 (97%)	6 (3%)	0	100	100
7	K	519/531 (98%)	469 (90%)	46 (9%)	4 (1%)	16	23
8	J	493/501 (98%)	435 (88%)	47 (10%)	11 (2%)	5	6
9	L	45/93 (48%)	45 (100%)	0	0	100	100
All	All	4099/4358 (94%)	3872 (94%)	210 (5%)	17 (0%)	32	39

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	E	82	ALA
3	F	153	HIS
7	K	76	GLU
7	K	143	GLY
8	J	6	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/224 (100%)	209 (94%)	14 (6%)	15	20
1	B	223/224 (100%)	217 (97%)	6 (3%)	40	54
2	D	341/342 (100%)	331 (97%)	10 (3%)	37	52
2	E	341/342 (100%)	332 (97%)	9 (3%)	41	56
3	C	437/439 (100%)	416 (95%)	21 (5%)	21	30
3	F	412/439 (94%)	393 (95%)	19 (5%)	23	32
4	G	104/168 (62%)	96 (92%)	8 (8%)	10	14
5	H	264/271 (97%)	244 (92%)	20 (8%)	11	14
6	I	160/186 (86%)	147 (92%)	13 (8%)	9	12
7	K	449/458 (98%)	404 (90%)	45 (10%)	6	7
8	J	452/454 (100%)	377 (83%)	75 (17%)	2	1

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
9	L	43/81 (53%)	40 (93%)	3 (7%)	12	17
All	All	3449/3628 (95%)	3206 (93%)	243 (7%)	15	17

5 of 243 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
6	I	144	MET
8	J	169	GLN
7	K	275	GLU
8	J	164	ILE
8	J	441	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
7	K	228	HIS
8	J	64	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

10 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MGN	F	403	3	6,9,10	0.80	0	5,12,14	0.94	0
3	MGN	C	403	-	6,9,10	0.64	0	5,12,14	0.87	0
3	AGM	F	275	3	10,11,12	0.46	0	6,13,15	0.24	0
3	GL3	C	448	-	2,3,4	1.78	1 (50%)	1,2,4	0.45	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GL3	F	448	3	2,3,4	1.79	1 (50%)	1,2,4	0.10	0
3	AGM	C	275	3	10,11,12	0.44	0	6,13,15	0.22	0
3	SMC	F	455	3	5,6,7	0.68	0	2,6,8	0.96	0
3	MHS	C	261	3	7,11,12	0.75	0	6,14,16	0.82	0
3	MHS	F	261	3	7,11,12	0.76	0	6,14,16	2.83	1 (16%)
3	SMC	C	455	3	5,6,7	0.69	0	2,6,8	1.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MGN	F	403	3	-	0/7/9/12	-
3	MGN	C	403	-	-	2/7/9/12	-
3	AGM	F	275	3	-	5/10/11/13	-
3	GL3	C	448	-	-	1/1/1/2	-
3	GL3	F	448	3	-	1/1/1/2	-
3	AGM	C	275	3	-	5/10/11/13	-
3	SMC	F	455	3	-	1/3/5/7	-
3	MHS	C	261	3	-	2/5/6/8	0/1/1/1
3	MHS	F	261	3	-	3/5/6/8	0/1/1/1
3	SMC	C	455	3	-	3/3/5/7	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	448	GL3	C-S	-2.51	1.72	1.80
3	F	448	GL3	C-S	-2.51	1.72	1.80

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	261	MHS	CB-CA-C	-6.69	98.93	111.47

There are no chirality outliers.

5 of 23 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	261	MHS	O-C-CA-CB

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	C	261	MHS	N-CA-CB-CG
3	C	275	AGM	O-C-CA-CB
3	C	275	AGM	CE2-CD-NE1-CZ
3	C	275	AGM	NH1-CZ-NE1-CD

There are no ring outliers.

4 monomers are involved in 25 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	403	MGN	5	0
3	C	448	GL3	17	0
3	C	261	MHS	1	0
3	C	455	SMC	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 3 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
14	S5Q	H	401	5	18,30,30	2.58	12 (66%)	-		
13	F43	F	602	12,3	61,71,71	2.04	4 (6%)	64,118,118	1.11	4 (6%)
13	F43	F	601	-	61,71,71	1.96	4 (6%)	64,118,118	1.02	5 (7%)
16	ATP	K	603	17	26,33,33	0.60	0	31,52,52	0.74	2 (6%)
14	S5Q	I	201	6	18,30,30	2.58	11 (61%)	-		
16	ATP	K	602	17	26,33,33	0.60	0	31,52,52	0.76	2 (6%)
14	S5Q	I	202	6	18,30,30	2.62	12 (66%)	-		
12	COM	C	602	13	6,6,6	0.42	0	7,8,8	0.68	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
10	SHT	E	601	-	26,27,27	1.28	2 (7%)	30,36,36	0.74	0
11	TP7	C	601	-	19,20,20	1.59	3 (15%)	24,26,26	1.10	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
13	F43	F	602	12,3	-	14/28/185/185	-
13	F43	F	601	-	-	9/28/185/185	-
16	ATP	K	603	17	-	2/18/38/38	0/3/3/3
16	ATP	K	602	17	-	3/18/38/38	0/3/3/3
12	COM	C	602	13	-	4/4/4/4	-
10	SHT	E	601	-	-	18/31/31/31	-
11	TP7	C	601	-	-	4/24/24/24	-

The worst 5 of 48 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
13	F	602	F43	NI-NA	9.77	2.10	1.89
13	F	601	F43	NI-NA	9.36	2.09	1.89
13	F	602	F43	NI-NB	9.19	2.09	1.89
13	F	601	F43	NI-NB	9.12	2.09	1.89
13	F	602	F43	NI-ND	7.18	2.05	1.89

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
13	F	602	F43	C2B-C1B-NB	3.47	107.04	101.84
13	F	601	F43	C2A-C3A-C4A	-3.25	97.39	102.36
13	F	602	F43	C3D-C4D-ND	3.20	107.31	102.34
13	F	601	F43	C3D-C4D-ND	3.11	107.18	102.34
13	F	601	F43	C2B-C3B-C4B	-2.73	98.55	101.63

There are no chirality outliers.

5 of 54 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
10	E	601	SHT	CC-CD-SG2-OS3
10	E	601	SHT	CC-CD-SG2-OS1
10	E	601	SHT	CK1-CK-N-CA
10	E	601	SHT	OK-CK-N-CA
10	E	601	SHT	CB-CA-N-CK


There are no ring outliers.


7 monomers are involved in 21 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
14	H	401	S5Q	1	0
13	F	602	F43	10	0
13	F	601	F43	5	0
16	K	603	ATP	2	0
12	C	602	COM	1	0
10	E	601	SHT	2	0
11	C	601	TP7	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
3	C	4
8	J	2

The worst 5 of 6 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	J	183:THR	C	184:ASN	N	3.53
1	C	402:SER	C	403:MGN	N	2.98
1	C	403:MGN	C	404:ARG	N	2.95
1	J	22:GLU	C	23:TYR	N	2.59
1	C	447:TYR	C	448:GL3	N	0.84