



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 6, 2025 – 01:29 pm BST

PDB ID : 9SDS / pdb\_00009sds  
Title : Structure of native leukocyte myeloperoxidase in complex with a truncated version of the Staphylococcal Peroxidase Inhibitor SPIN and chloride at pH 5.5  
Authors : Leitgeb, U.; Pfanzagl, V.  
Deposited on : 2025-08-14  
Resolution : 2.49 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.46


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	5504 (2.50-2.50)
Clashscore	180529	6282 (2.50-2.50)
Ramachandran outliers	177936	6191 (2.50-2.50)
Sidechain outliers	177891	6193 (2.50-2.50)
RSRZ outliers	164620	5504 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	114	<div> <div>0%</div> <div>74%</div> <div>18%</div> <div>• 8%</div> </div>
1	B	114	<div> <div>3%</div> <div>72%</div> <div>19%</div> <div>• 8%</div> </div>
2	E	60	<div> <div>73%</div> <div>13%</div> <div>5%</div> <div>8%</div> </div>
2	F	60	<div> <div>18%</div> <div>62%</div> <div>23%</div> <div>• • 10%</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	467	<div><div>%</div><div><div></div><div>81%</div><div>16%</div><div>.</div></div></div>
3	D	467	<div><div>%</div><div><div></div><div>81%</div><div>16%</div><div>..</div></div></div>
4	G	6	<div><div><div></div><div>33%</div><div>67%</div></div></div>
4	J	6	<div><div><div></div><div>17%</div><div>67%</div><div>17%</div></div></div>
5	H	2	<div><div><div></div><div>100%</div></div></div>
5	I	2	<div><div><div></div><div>100%</div></div></div>
5	K	2	<div><div><div></div><div>100%</div></div></div>

## 2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 10651 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Myeloperoxidase light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	105	Total	C	N	O	S	0	0	0
			842	532	149	156	5			
1	B	105	Total	C	N	O	S	0	0	0
			842	532	149	156	5			


- Molecule 2 is a protein called Myeloperoxidase inhibitor SPIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	55	Total	C	N	O	0	0	0
			448	284	75	89			
2	F	54	Total	C	N	O	0	0	0
			436	275	74	87			

- Molecule 3 is a protein called Myeloperoxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	465	Total	C	N	O	S	0	0	0
			3726	2348	686	665	27			
3	D	465	Total	C	N	O	S	0	0	0
			3726	2348	686	665	27			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	G	6	Total	C	N	O	0	0	0
			71	40	2	29			

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	J	6	Total	C	N	O	0	0	0
			71	40	2	29			

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	H	2	Total	C	N	O	0	0	0
			28	16	2	10			
5	I	2	Total	C	N	O	0	0	0
			28	16	2	10			
5	K	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 6 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	Fe	N	O	0
			43	34	1	4	4	0
6	B	1	Total	C	Fe	N	O	0
			43	34	1	4	4	0

- Molecule 7 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total I 1 1	0	0
7	B	1	Total I 1 1	0	0


- Molecule 8 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	C	1	Total Ca 1 1	0	0
8	D	1	Total Ca 1 1	0	0

- Molecule 9 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	C	2	Total Cl 2 2	0	0
9	D	1	Total Cl 1 1	0	0

- Molecule 10 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	42	Total	O	0	0
			42	42		
11	E	3	Total	O	0	0
			3	3		
11	C	137	Total	O	0	0
			137	137		
11	B	28	Total	O	0	0
			28	28		
11	D	87	Total	O	0	0
			87	87		
11	F	1	Total	O	0	0
			1	1		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Myeloperoxidase light chain




- Molecule 1: Myeloperoxidase light chain




- Molecule 2: Myeloperoxidase inhibitor SPIN




- Molecule 2: Myeloperoxidase inhibitor SPIN





- Molecule 3: Myeloperoxidase








• Molecule 3: Myeloperoxidase



• Molecule 4: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 4: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




• Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  100%

MAG1  
MAG2

- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  100%

MAG1  
MAG2

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	111.16Å 111.16Å 241.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.74 – 2.49 48.74 – 2.49	Depositor EDS
% Data completeness (in resolution range)	98.0 (48.74-2.49) 98.0 (48.74-2.49)	Depositor EDS
$R_{merge}$	0.70	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.13 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.8.0430	Depositor
R, $R_{free}$	0.188 , 0.234 0.188 , 0.234	Depositor DCC
$R_{free}$ test set	2639 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.3	Xtriage
Anisotropy	0.077	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 25.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	10651	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, BMA, CA, CL, MAN, HEM, IOD, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.77	0/867	1.56	14/1181 (1.2%)
1	B	0.74	0/867	1.55	15/1181 (1.3%)
2	E	0.70	0/455	1.85	11/610 (1.8%)
2	F	0.66	0/442	1.87	9/592 (1.5%)
3	C	0.77	0/3812	1.55	56/5173 (1.1%)
3	D	0.72	0/3812	1.55	50/5173 (1.0%)
All	All	0.74	0/10255	1.58	155/13910 (1.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
2	E	0	1
3	C	0	6
3	D	0	5
All	All	0	13

There are no bond length outliers.

The worst 5 of 155 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	520	GLU	N-CA-CB	12.11	124.68	110.03
3	D	506	GLN	CB-CA-C	11.57	125.53	108.86
3	C	506	GLN	CB-CA-C	11.03	125.49	108.61
3	C	625	LYS	N-CA-CB	10.16	125.07	109.94
3	D	625	LYS	CB-CA-C	10.12	127.59	110.79

There are no chirality outliers.

5 of 13 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	314	ARG	Sidechain
3	C	351	ARG	Sidechain
3	C	436	ARG	Sidechain
3	C	515	ARG	Sidechain
2	E	70	ARG	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	842	0	800	6	0
1	B	842	0	800	7	0
2	E	448	0	436	7	0
2	F	436	0	427	7	0
3	C	3726	0	3720	25	0
3	D	3726	0	3721	28	0
4	G	71	0	61	0	0
4	J	71	0	61	1	0
5	H	28	0	25	0	0
5	I	28	0	25	0	0
5	K	28	0	25	0	0
6	A	43	0	30	2	0
6	B	43	0	30	1	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
8	C	1	0	0	0	0
8	D	1	0	0	0	0
9	C	2	0	0	0	0
9	D	1	0	0	0	0
10	D	14	0	13	0	0
11	A	42	0	0	0	0
11	B	28	0	0	0	0
11	C	137	0	0	0	0
11	D	87	0	0	0	0
11	E	3	0	0	2	0
11	F	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	10651	0	10174	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 76 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:502:HIS:HD1	3:C:587:ASN:HD21	1.33	0.77
3:D:502:HIS:HD1	3:D:587:ASN:HD21	1.33	0.76
2:E:70:ARG:HH12	2:E:81:ILE:HD11	1.50	0.76
3:C:716:ASN:HD22	3:C:718:PHE:H	1.38	0.70
2:E:58:ASP:OD1	2:E:60:ASN:HB2	1.95	0.66

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	103/114 (90%)	99 (96%)	4 (4%)	0	100	100
1	B	103/114 (90%)	97 (94%)	6 (6%)	0	100	100
2	E	53/60 (88%)	53 (100%)	0	0	100	100
2	F	52/60 (87%)	51 (98%)	1 (2%)	0	100	100
3	C	463/467 (99%)	448 (97%)	15 (3%)	0	100	100
3	D	463/467 (99%)	445 (96%)	18 (4%)	0	100	100
All	All	1237/1282 (96%)	1193 (96%)	44 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	90/97 (93%)	89 (99%)	1 (1%)	70	87
1	B	90/97 (93%)	90 (100%)	0	100	100
2	E	49/53 (92%)	49 (100%)	0	100	100
2	F	48/53 (91%)	45 (94%)	3 (6%)	15	30
3	C	411/412 (100%)	397 (97%)	14 (3%)	32	58
3	D	411/412 (100%)	396 (96%)	15 (4%)	30	56
All	All	1099/1124 (98%)	1066 (97%)	33 (3%)	36	63

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	D	713	SER
3	D	743	GLU
2	F	97	LYS
3	C	713	SER
3	C	625	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
3	D	305	ASN
3	D	423	HIS
3	D	575	GLN
3	D	562	GLN
3	C	716	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

18 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	NAG	G	1	4,3	14,14,15	0.48	0	17,19,21	1.30	2 (11%)
4	NAG	G	2	4	14,14,15	0.40	0	17,19,21	1.01	0
4	BMA	G	3	4	11,11,12	0.64	0	15,15,17	1.06	1 (6%)
4	MAN	G	4	4	11,11,12	1.40	2 (18%)	15,15,17	1.17	1 (6%)
4	MAN	G	5	4	11,11,12	0.77	0	15,15,17	1.66	1 (6%)
4	FUC	G	6	4	10,10,11	0.65	0	14,14,16	0.87	0
5	NAG	H	1	5,3	14,14,15	0.52	0	17,19,21	1.84	3 (17%)
5	NAG	H	2	5	14,14,15	0.75	0	17,19,21	1.75	4 (23%)
5	NAG	I	1	5,3	14,14,15	0.41	0	17,19,21	1.15	1 (5%)
5	NAG	I	2	5	14,14,15	0.56	0	17,19,21	1.11	2 (11%)
4	NAG	J	1	4,3	14,14,15	0.42	0	17,19,21	1.05	0
4	NAG	J	2	4	14,14,15	0.39	0	17,19,21	1.25	2 (11%)
4	BMA	J	3	4	11,11,12	0.66	0	15,15,17	1.57	3 (20%)
4	MAN	J	4	4	11,11,12	1.40	1 (9%)	15,15,17	2.40	4 (26%)
4	MAN	J	5	4	11,11,12	0.95	1 (9%)	15,15,17	1.90	2 (13%)
4	FUC	J	6	4	10,10,11	1.01	0	14,14,16	1.05	1 (7%)
5	NAG	K	1	5,3	14,14,15	0.50	0	17,19,21	1.31	1 (5%)
5	NAG	K	2	5	14,14,15	0.65	0	17,19,21	2.32	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	1	4,3	-	0/6/23/26	0/1/1/1
4	NAG	G	2	4	-	0/6/23/26	0/1/1/1
4	BMA	G	3	4	-	0/2/19/22	0/1/1/1
4	MAN	G	4	4	-	1/2/19/22	0/1/1/1
4	MAN	G	5	4	-	1/2/19/22	0/1/1/1
4	FUC	G	6	4	-	-	0/1/1/1
5	NAG	H	1	5,3	-	0/6/23/26	0/1/1/1
5	NAG	H	2	5	-	2/6/23/26	0/1/1/1
5	NAG	I	1	5,3	-	2/6/23/26	0/1/1/1
5	NAG	I	2	5	-	2/6/23/26	0/1/1/1
4	NAG	J	1	4,3	-	0/6/23/26	0/1/1/1
4	NAG	J	2	4	-	0/6/23/26	0/1/1/1
4	BMA	J	3	4	-	0/2/19/22	0/1/1/1
4	MAN	J	4	4	-	2/2/19/22	0/1/1/1
4	MAN	J	5	4	-	0/2/19/22	0/1/1/1
4	FUC	J	6	4	-	-	0/1/1/1
5	NAG	K	1	5,3	-	2/6/23/26	0/1/1/1
5	NAG	K	2	5	-	1/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	J	4	MAN	O5-C5	3.00	1.49	1.43
4	G	4	MAN	C4-C5	2.81	1.58	1.53
4	J	5	MAN	C2-C3	2.32	1.55	1.52
4	G	4	MAN	O5-C5	2.19	1.47	1.43

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	K	2	NAG	C2-N2-C7	6.05	131.52	122.90
4	J	4	MAN	C1-O5-C5	5.59	119.77	112.19
4	G	5	MAN	C1-O5-C5	5.47	119.60	112.19
4	J	5	MAN	O2-C2-C3	5.26	120.67	110.14
5	K	2	NAG	C1-C2-N2	-4.84	102.22	110.49

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	J	4	MAN	O5-C5-C6-O6

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
5	K	1	NAG	O5-C5-C6-O6
5	I	1	NAG	C8-C7-N2-C2
4	J	4	MAN	C4-C5-C6-O6
5	K	1	NAG	C4-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	J	5	MAN	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 7 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
10	NAG	D	801	3	14,14,15	0.83	0	17,19,21	2.38	6 (35%)
6	HEM	A	301	3,1	41,50,50	1.57	8 (19%)	45,82,82	1.90	11 (24%)
6	HEM	B	301	3,1	41,50,50	1.74	8 (19%)	45,82,82	1.44	4 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	NAG	D	801	3	-	2/6/23/26	0/1/1/1
6	HEM	A	301	3,1	-	4/12/54/54	-
6	HEM	B	301	3,1	-	4/12/54/54	-

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	301	HEM	C4D-ND	-5.36	1.31	1.40
6	B	301	HEM	C1D-C2D	-4.71	1.35	1.44
6	A	301	HEM	FE-ND	4.14	2.17	1.96
6	A	301	HEM	C4D-C3D	-4.07	1.38	1.45
6	B	301	HEM	C4D-C3D	-4.04	1.38	1.45

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	301	HEM	C4B-CHC-C1C	6.32	130.90	122.56
10	D	801	NAG	C1-O5-C5	5.47	119.61	112.19
6	B	301	HEM	C4C-CHD-C1D	5.13	129.33	122.56
6	B	301	HEM	C4B-CHC-C1C	4.80	128.90	122.56
10	D	801	NAG	C1-C2-N2	4.51	118.19	110.49

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	D	801	NAG	C8-C7-N2-C2
10	D	801	NAG	O7-C7-N2-C2
6	B	301	HEM	CAD-CBD-CGD-O2D
6	A	301	HEM	CAD-CBD-CGD-O1D
6	A	301	HEM	CAD-CBD-CGD-O2D

There are no ring outliers.

2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	301	HEM	2	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	301	HEM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	105/114 (92%)	-0.64	1 (0%) 79 76	8, 18, 42, 76	0
1	B	105/114 (92%)	-0.43	3 (2%) 54 50	13, 25, 52, 75	0
2	E	55/60 (91%)	-0.18	0 100 100	23, 40, 67, 74	0
2	F	54/60 (90%)	1.30	11 (20%) 3 3	54, 74, 98, 100	0
3	C	465/467 (99%)	-0.71	4 (0%) 81 78	7, 19, 39, 66	0
3	D	465/467 (99%)	-0.43	5 (1%) 77 74	11, 29, 53, 94	0
All	All	1249/1282 (97%)	-0.47	24 (1%) 66 63	7, 24, 61, 100	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	523	PRO	4.0
3	D	715	ASN	3.5
2	F	97	LYS	3.4
3	C	383	HIS	3.2
2	F	44	ASN	3.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	MAN	J	4	11/12	0.77	0.14	51,57,61,61	0
5	NAG	I	2	14/15	0.77	0.16	49,65,76,77	0
5	NAG	K	2	14/15	0.79	0.13	62,65,68,72	0
4	MAN	G	4	11/12	0.82	0.13	49,51,54,58	0
5	NAG	H	2	14/15	0.83	0.13	45,50,54,55	0
5	NAG	K	1	14/15	0.91	0.10	42,49,53,58	0
4	MAN	J	5	11/12	0.91	0.07	26,28,29,30	0
4	FUC	J	6	10/11	0.92	0.10	34,39,43,43	0
5	NAG	I	1	14/15	0.93	0.08	32,37,43,51	0
4	MAN	G	5	11/12	0.94	0.08	30,33,35,39	0
4	FUC	G	6	10/11	0.94	0.07	26,29,31,33	0
4	BMA	G	3	11/12	0.95	0.07	32,33,38,42	0
4	NAG	G	1	14/15	0.95	0.06	22,25,26,27	0
4	NAG	J	2	14/15	0.95	0.07	21,22,26,26	0
4	BMA	J	3	11/12	0.95	0.06	29,29,34,40	0
5	NAG	H	1	14/15	0.96	0.07	26,31,34,40	0
4	NAG	J	1	14/15	0.96	0.06	22,23,25,28	0
4	NAG	G	2	14/15	0.96	0.07	22,26,29,29	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around Chain G:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around Chain J:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain H:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain I:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








## 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
10	NAG	D	801	14/15	0.85	0.12	50,54,57,58	0
9	CL	D	803	1/1	0.93	0.08	46,46,46,46	0
9	CL	C	802	1/1	0.94	0.10	43,43,43,43	0
6	HEM	B	301	43/43	0.97	0.06	15,19,23,30	0
6	HEM	A	301	43/43	0.98	0.05	8,9,12,16	0
9	CL	C	803	1/1	0.98	0.04	24,24,24,24	0
8	CA	C	801	1/1	0.99	0.05	5,5,5,5	0
7	IOD	A	302	1/1	1.00	0.01	11,11,11,11	0
8	CA	D	802	1/1	1.00	0.07	10,10,10,10	0
7	IOD	B	302	1/1	1.00	0.01	15,15,15,15	0



The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.