



Full wwPDB EM Validation Report ⓘ

Oct 7, 2025 – 10:08 pm BST

PDB ID : 9SSR / pdb_00009ssr
EMDB ID : EMD-55192
Title : Human Methionine Synthase With Methyltetrahydrofolate, Hydroxocobalamin, and SAM, N-Half From Full-Length
Authors : Ferreira, D.S.M.; Yue, W.W.; McCorvie, T.J.
Deposited on : 2025-09-26
Resolution : 2.84 Å(reported)
Based on initial model : .

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev129
Mogul : 1.8.4, CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	210492	15764	-
Ramachandran outliers	207382	16835	-
Sidechain outliers	206894	16415	-
Q-score	-	25397	11884 (2.34 - 3.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1265	

2 Entry composition [i](#)


There are 2 unique types of molecules in this entry. The entry contains 4925 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methionine synthase.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	638	4892	3089	832	932	39	0	0

- Molecule 2 is N-[4-({[(6S)-2-AMINO-4-HYDROXY-5-METHYL-5,6,7,8-TETRAHYDRO PTERIDIN-6-YL]METHYL}AMINO)BENZOYL]-L-GLUTAMIC ACID (CCD ID: THH) (formula: C₂₀H₂₅N₇O₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
2	A	1	33	20	7	6	0

GLU	VAL	ARG	ASP	LYS
VAL	GLU	LEU	ARG	ASP
GLU	GLU	ALA	LEU	SER
LYS	ASP	ASP	ALA	SER
TRP	TRP	ILE	GLU	SER
LEU	LEU	GLU	ALA	THR
GLY	GLY	GLN	PHE	GLU
PRO	PRO	SER	ALA	PRO
THR	THR	GLY	GLU	TYR
ASP	ASP	ILE	GLY	CYS
ASP	ASP	LEU	THR	LEU
		ARG	LEU	SER
		THR	GLU	SER
		GLU	ARG	ASP
		VAL	PHE	ASP
		SER	ARG	ILE
		SER	ARG	ALA
		LEU	ARG	ALA
		ALA	GLU	PRO
		MET	LEU	PRO
		ALA	LEU	LEU
		ALA	GLU	GLY
		ALA	GLY	GLY
		VAL	SER	ILE
		VAL	GLY	ARG
		SER	GLU	ASP
		ASN	GLN	TYR
		LEU	GLU	GLY
		LYS	LEU	LEU
		SER	ARG	SER
		LYS	LEU	LYS
		TYR	ARG	VAL
		PHE	TYR	VAL
		ALA	TYR	GLU
		VAL	GLY	GLU
		GLY	ILE	LEU
		LYS	ARG	SER
		ILE	PRO	LYS
		SER	ALA	ALA
		LYS	PRO	GLU
		ASP	GLY	ASP
		GLN	TYR	ASP
		VAL	PRO	GLY
		GLU	SER	ASP
		GLU	GLN	ASP
		TYR	PRO	TYR
		ALA	ASP	SER
		LEU	HIS	SER
		ARG	THR	ILE
		LYS	GLU	MET
		ASN	LYS	VAL
		ILE	THR	LYS
		SER	LEU	ALA
		VAL	THR	ALA
		ALA	MET	GLY
		VAL	TRP	GLY

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	76292	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	65.2	Depositor
Minimum defocus (nm)	600	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	165000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.681	Depositor
Minimum map value	-0.367	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.017	Depositor
Recommended contour level	0.08	Depositor
Map size (Å)	203.20001, 203.20001, 203.20001	wwPDB
Map dimensions	200, 200, 200	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.016, 1.016, 1.016	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: THH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.13	0/4978	0.35	0/6725

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4892	0	4888	73	0
2	A	33	0	23	1	0
All	All	4925	0	4911	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:245:VAL:O	1:A:249:SER:OG	2.05	0.73
1:A:40:MET:HA	1:A:40:MET:HE2	1.74	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:230:ASP:OD1	1:A:231:LYS:N	2.23	0.66
1:A:414:ASP:HA	1:A:447:CYS:HB3	1.80	0.63
1:A:114:ALA:O	1:A:118:ASN:ND2	2.31	0.61
1:A:86:LEU:HB3	1:A:132:VAL:HG11	1.82	0.61
1:A:547:MET:HG2	1:A:549:GLU:HG2	1.86	0.57
1:A:342:LYS:NZ	1:A:343:PRO:O	2.37	0.57
1:A:484:LYS:O	1:A:488:ILE:HG13	2.05	0.56
1:A:584:PHE:HD1	1:A:587:MET:HE1	1.71	0.56
1:A:403:LYS:HG3	1:A:441:ILE:HD11	1.87	0.56
1:A:67:ASN:O	1:A:96:ASN:ND2	2.27	0.55
1:A:564:ILE:HG23	1:A:568:LEU:HD12	1.89	0.54
1:A:136:THR:OG1	1:A:140:ARG:NH1	2.40	0.53
1:A:259:ASN:ND2	1:A:260:CYS:SG	2.80	0.53
1:A:401:VAL:HG12	1:A:405:GLN:NE2	2.23	0.53
1:A:559:HIS:O	1:A:563:VAL:HG23	2.09	0.52
1:A:25:ILE:HA	1:A:28:LYS:HE2	1.91	0.52
1:A:49:GLU:H	1:A:49:GLU:CD	2.17	0.52
1:A:299:GLU:HG3	1:A:303:MET:HE2	1.91	0.52
1:A:48:GLU:HG2	1:A:64:LYS:HB2	1.92	0.51
1:A:387:ALA:O	1:A:391:MET:HG2	2.10	0.51
1:A:401:VAL:HG12	1:A:405:GLN:HE21	1.75	0.51
1:A:161:ARG:HB3	1:A:164:TYR:HB3	1.93	0.50
1:A:262:LEU:HB2	1:A:267:MET:HE3	1.94	0.50
1:A:225:SER:HB3	1:A:283:LEU:HD11	1.92	0.50
1:A:644:GLU:H	1:A:644:GLU:CD	2.20	0.50
1:A:537:ASP:OD2	2:A:1301:THH:NA2	2.42	0.50
1:A:641:GLU:O	1:A:645:LYS:HG3	2.13	0.49
1:A:449:ASP:HA	1:A:470:ASN:HB3	1.95	0.49
1:A:548:GLU:N	1:A:548:GLU:OE2	2.46	0.49
1:A:648:ARG:O	1:A:652:THR:HG23	2.14	0.48
1:A:421:MET:O	1:A:421:MET:HG3	2.13	0.48
1:A:231:LYS:HD2	1:A:231:LYS:HA	1.63	0.47
1:A:301:PRO:HB3	1:A:330:HIS:HA	1.96	0.47
1:A:428:MET:HE3	1:A:448:ILE:HG23	1.96	0.47
1:A:483:GLU:OE2	1:A:483:GLU:HA	2.14	0.47
1:A:435:ILE:HG21	1:A:446:LEU:HD11	1.97	0.47
1:A:66:ASN:HD22	1:A:69:ILE:HB	1.80	0.47
1:A:429:THR:HG23	1:A:462:CYS:SG	2.55	0.47
1:A:413:LEU:O	1:A:447:CYS:N	2.45	0.46
1:A:34:ASP:N	1:A:92:ILE:O	2.44	0.46
1:A:584:PHE:CD2	1:A:594:MET:HE1	2.51	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:256:ILE:HG21	1:A:278:THR:HG21	1.97	0.46
1:A:408:MET:HG2	1:A:615:GLY:HA3	1.98	0.45
1:A:416:ASN:ND2	1:A:449:ASP:OD1	2.26	0.45
1:A:340:ASN:OD1	1:A:340:ASN:N	2.50	0.45
1:A:196:ILE:HG12	1:A:202:ALA:HB2	1.99	0.45
1:A:66:ASN:ND2	1:A:69:ILE:HB	2.32	0.44
1:A:435:ILE:HG23	1:A:441:ILE:HG22	1.99	0.44
1:A:310:ASP:HA	1:A:313:MET:HE2	2.00	0.43
1:A:360:LEU:H	1:A:443:LYS:HA	1.83	0.43
1:A:472:ILE:HG22	1:A:496:VAL:O	2.19	0.43
1:A:374:ILE:HB	1:A:611:ILE:HA	2.00	0.43
1:A:574:SER:HA	1:A:609:MET:O	2.19	0.42
1:A:599:LEU:HD23	1:A:599:LEU:HA	1.87	0.42
1:A:119:MET:HE3	1:A:119:MET:HB3	1.77	0.42
1:A:297:TYR:HB2	1:A:325:GLY:HA3	2.02	0.42
1:A:482:LEU:HD11	1:A:526:LYS:HD2	2.01	0.42
1:A:274:ILE:O	1:A:278:THR:HG22	2.19	0.42
1:A:403:LYS:NZ	1:A:407:GLU:OE2	2.50	0.42
1:A:31:MET:HE2	1:A:31:MET:HB3	1.92	0.41
1:A:195:THR:HA	1:A:259:ASN:HD21	1.86	0.41
1:A:256:ILE:O	1:A:282:VAL:HA	2.19	0.41
1:A:69:ILE:HD12	1:A:69:ILE:HA	1.90	0.41
1:A:45:LYS:HE2	1:A:45:LYS:HB2	1.92	0.41
1:A:372:VAL:HB	1:A:609:MET:HB3	2.02	0.41
1:A:257:GLY:HA3	1:A:283:LEU:O	2.21	0.41
1:A:419:ASP:OD1	1:A:420:GLY:N	2.54	0.41
1:A:495:MET:HE3	1:A:497:VAL:HG22	2.04	0.41
1:A:267:MET:HE3	1:A:267:MET:HB2	1.83	0.40
1:A:554:ALA:HB2	1:A:580:LEU:HD11	2.03	0.40
1:A:649:TYR:O	1:A:652:THR:OG1	2.33	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	636/1265 (50%)	613 (96%)	23 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	525/1059 (50%)	517 (98%)	8 (2%)	60	81

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	70	LEU
1	A	76	ASP
1	A	93	ILE
1	A	119	MET
1	A	426	SER
1	A	567	THR
1	A	587	MET
1	A	594	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	HIS
1	A	307	HIS
1	A	318	ASN
1	A	556	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	THH	A	1301	-	32,35,35	2.09	8 (25%)	34,49,49	1.31	2 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	THH	A	1301	-	-	3/22/35/35	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1301	THH	C2-NA2	6.20	1.46	1.33
2	A	1301	THH	C-N	5.73	1.46	1.34
2	A	1301	THH	C4A-N5	3.54	1.50	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1301	THH	C15-N10	3.30	1.48	1.38
2	A	1301	THH	C7-N8	-2.75	1.40	1.44
2	A	1301	THH	C12-C	2.51	1.55	1.50
2	A	1301	THH	O4-C4	2.44	1.40	1.29
2	A	1301	THH	O-C	-2.27	1.18	1.23

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1301	THH	N3-C2-N1	-3.33	120.20	125.42
2	A	1301	THH	C2-N1-C8A	2.44	120.02	114.54

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1301	THH	OE1-CD-CG-CB
2	A	1301	THH	OE2-CD-CG-CB
2	A	1301	THH	CA-CB-CG-CD

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1301	THH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-55192. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 100




Y Index: 100




Z Index: 100


6.2.2 Raw map



X Index: 100



Y Index: 100




Z Index: 100


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 104




Y Index: 97




Z Index: 114


6.3.2 Raw map



X Index: 104



Y Index: 97




Z Index: 84


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.08. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 29 nm^3 ; this corresponds to an approximate mass of 26 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.352 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.352 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.84	-	-
Author-provided FSC curve	2.84	3.22	2.88
Unmasked-calculated*	3.66	4.15	3.73

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.66 differs from the reported value 2.84 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-55192 and PDB model 9SSR. Per-residue inclusion information can be found in [section 3](#) on [page 4](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.08 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.08).

9.4 Atom inclusion [i](#)



At the recommended contour level, 87% of all backbone atoms, 83% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.08) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div><div></div></div> 0.8320	<div><div></div></div> 0.5550
A	<div><div></div></div> 0.8320	<div><div></div></div> 0.5550

