



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2024 – 03:23 PM EDT

PDB ID : 6SW5  
Title : Crystal structure of the human S-adenosylmethionine synthetase 1 (ligand-free form)  
Authors : Panmanee, J.; Antoyuk, S.V.; Hasnain, S.S.  
Deposited on : 2019-09-19  
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	395	<div> <div>83%</div> <div>8%</div> <div>9%</div> </div>
1	B	395	<div> <div>88%</div> <div>8%</div> </div>
1	C	395	<div> <div>2%</div> <div>88%</div> <div>8%</div> </div>
1	D	395	<div> <div>85%</div> <div>10%</div> </div>

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 11679 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called S-adenosylmethionine synthase isoform type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	360	Total	C	N	O	S	0	3	0
			2795	1771	484	522	18			
1	B	365	Total	C	N	O	S	0	0	0
			2831	1791	490	533	17			
1	C	362	Total	C	N	O	S	0	1	0
			2811	1776	490	527	18			
1	D	355	Total	C	N	O	S	0	2	0
			2772	1756	483	516	17			

- Molecule 2 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	4	3		
2	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	105	Total 107	O 107	0	2
4	B	111	Total 113	O 113	0	2
4	C	89	Total 90	O 90	0	1
4	D	94	Total 98	O 98	0	4


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: S-adenosylmethionine synthase isoform type-1




- Molecule 1: S-adenosylmethionine synthase isoform type-1




- Molecule 1: S-adenosylmethionine synthase isoform type-1



- Molecule 1: S-adenosylmethionine synthase isoform type-1





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	218.68Å 61.00Å 119.76Å 90.00° 90.55° 90.00°	Depositor
Resolution (Å)	58.76 – 2.35 58.76 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.9 (58.76-2.35) 99.9 (58.76-2.35)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.96 (at 2.34Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, $R_{free}$	0.188 , 0.239 0.193 , 0.239	Depositor DCC
$R_{free}$ test set	3386 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.9	Xtriage
Anisotropy	0.356	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 51.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.007 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	11679	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.53	0/2857	0.73	1/3864 (0.0%)
1	B	0.54	0/2885	0.74	0/3906
1	C	0.51	0/2867	0.71	0/3879
1	D	0.51	0/2827	0.73	1/3825 (0.0%)
All	All	0.52	0/11436	0.73	2/15474 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	362	ARG	NE-CZ-NH2	5.26	122.93	120.30
1	A	382	ARG	NE-CZ-NH2	5.05	122.83	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2795	0	2777	19	0
1	B	2831	0	2808	10	0
1	C	2811	0	2802	8	0
1	D	2772	0	2763	8	0
2	A	7	0	10	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	7	0	10	1	0
3	A	24	0	36	1	0
3	B	8	0	12	2	0
3	C	12	0	18	0	0
3	D	4	0	6	0	0
4	A	107	0	0	1	0
4	B	113	0	0	0	0
4	C	90	0	0	0	0
4	D	98	0	0	0	0
All	All	11679	0	11242	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:261:VAL:HG11	1:D:261:VAL:HG11	1.55	0.86
1:B:266:ILE:HD11	1:B:281:ALA:HB2	1.62	0.79
1:A:261:VAL:HG21	1:B:261:VAL:HG21	1.67	0.74
1:B:266:ILE:HD11	1:B:281:ALA:CB	2.21	0.69
1:A:26:GLY:HA2	1:A:161[B]:ASN:HD21	1.66	0.59
1:A:376[B]:CYS:SG	4:A:577:HOH:O	2.45	0.58
1:A:354:ASP:HA	3:A:405:EDO:H12	1.87	0.57
1:A:261:VAL:HG11	1:B:261:VAL:HG11	1.89	0.55
1:A:176:LEU:HD22	1:A:207:VAL:HG21	1.90	0.54
1:C:45:HIS:CE1	1:C:71:ILE:HG21	2.44	0.52
1:C:261:VAL:HG21	1:D:261:VAL:HG21	1.92	0.52
1:B:223:LYS:O	1:B:228:ARG:HG2	2.10	0.52
1:D:56:CYS:SG	1:D:67:LEU:HD21	2.50	0.52
1:B:139:PHE:H	3:B:1002:EDO:H21	1.76	0.50
1:A:109:ALA:HB2	1:C:107:LEU:CD2	2.43	0.49
1:A:26:GLY:HA2	1:A:161[B]:ASN:ND2	2.27	0.48
1:A:109:ALA:HB2	1:C:107:LEU:HD23	1.96	0.47
1:B:143:THR:O	1:B:149:CYS:HA	2.14	0.47
1:A:321:ALA:HB2	1:A:328:LEU:HD21	1.97	0.46
1:C:176:LEU:HD22	1:C:207:VAL:HG21	1.97	0.46
1:A:105:ASN:HB3	1:A:107:LEU:HD21	1.97	0.46
1:C:65:VAL:HG21	1:C:87:ILE:HD11	1.98	0.45
1:A:276:ALA:HB2	2:B:1001:PEG:H22	1.98	0.45
1:C:143:THR:O	1:C:149:CYS:HA	2.17	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:143:THR:O	1:A:149:CYS:HA	2.17	0.44
1:B:139:PHE:H	3:B:1002:EDO:C2	2.30	0.44
1:D:321:ALA:HB2	1:D:328:LEU:HD11	1.99	0.44
1:A:29:HIS:CE1	1:A:265:LYS:HE2	2.53	0.44
1:D:71:ILE:HD12	1:D:110:LEU:HD13	2.00	0.43
1:D:289[B]:LYS:HD3	1:D:289[B]:LYS:HA	1.77	0.43
1:A:307:LYS:HG3	1:A:390:PRO:HG3	2.00	0.43
1:B:148:GLU:HG3	1:B:155:ILE:HD12	2.01	0.43
1:A:138:MET:HG2	1:A:318:VAL:HG13	2.01	0.42
1:B:59:VAL:HG22	1:B:261:VAL:HG12	2.01	0.42
1:A:196:ILE:HD12	1:A:196:ILE:N	2.34	0.41
1:D:180:SER:HB3	1:D:207:VAL:HG23	2.02	0.41
1:A:299:ARG:HG2	1:A:380:PHE:CD1	2.56	0.41
1:D:357:PRO:O	1:D:361:VAL:HG13	2.21	0.40
1:A:191:ASP:HB2	1:A:196:ILE:HD13	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	355/395 (90%)	343 (97%)	11 (3%)	1 (0%)	41	47
1	B	359/395 (91%)	347 (97%)	11 (3%)	1 (0%)	41	47
1	C	357/395 (90%)	348 (98%)	9 (2%)	0	100	100
1	D	349/395 (88%)	340 (97%)	9 (3%)	0	100	100
All	All	1420/1580 (90%)	1378 (97%)	40 (3%)	2 (0%)	51	63

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	62	THR
1	B	62	THR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/332 (90%)	292 (97%)	8 (3%)	44	55
1	B	304/332 (92%)	298 (98%)	6 (2%)	55	66
1	C	303/332 (91%)	299 (99%)	4 (1%)	69	80
1	D	298/332 (90%)	288 (97%)	10 (3%)	37	46
All	All	1205/1328 (91%)	1177 (98%)	28 (2%)	50	61

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	MET
1	A	60	CYS
1	A	61	LYS
1	A	99	PHE
1	A	107	LEU
1	A	138	MET
1	A	172	LEU
1	A	215	LEU
1	B	134	ASP
1	B	225	GLN
1	B	252	ILE
1	B	266	ILE
1	B	290	VAL
1	B	347	ASP
1	C	19	MET
1	C	114	SER
1	C	206	SER
1	C	361	VAL
1	D	19	MET
1	D	60	CYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	67	LEU
1	D	137	LEU
1	D	225	GLN
1	D	289[A]	LYS
1	D	289[B]	LYS
1	D	318	VAL
1	D	361	VAL
1	D	384	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	105	ASN
1	C	126	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PEG	B	1001	-	6,6,6	0.83	0	5,5,5	0.83	0
3	EDO	A	406	-	3,3,3	0.38	0	2,2,2	0.44	0
2	PEG	A	401	-	6,6,6	0.56	0	5,5,5	0.52	0
3	EDO	C	401	-	3,3,3	0.50	0	2,2,2	0.36	0
3	EDO	D	401	-	3,3,3	0.29	0	2,2,2	0.75	0
3	EDO	A	403	-	3,3,3	0.39	0	2,2,2	0.48	0
3	EDO	B	1003	-	3,3,3	0.26	0	2,2,2	0.82	0
3	EDO	B	1002	-	3,3,3	0.48	0	2,2,2	0.51	0
3	EDO	C	403	-	3,3,3	0.25	0	2,2,2	0.76	0
3	EDO	A	404	-	3,3,3	0.57	0	2,2,2	0.11	0
3	EDO	A	407	-	3,3,3	0.28	0	2,2,2	0.59	0
3	EDO	A	405	-	3,3,3	0.44	0	2,2,2	0.33	0
3	EDO	C	402	-	3,3,3	0.42	0	2,2,2	0.26	0
3	EDO	A	402	-	3,3,3	0.55	0	2,2,2	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	B	1001	-	-	2/4/4/4	-
3	EDO	A	406	-	-	1/1/1/1	-
2	PEG	A	401	-	-	4/4/4/4	-
3	EDO	C	401	-	-	0/1/1/1	-
3	EDO	D	401	-	-	0/1/1/1	-
3	EDO	A	403	-	-	0/1/1/1	-
3	EDO	B	1003	-	-	1/1/1/1	-
3	EDO	B	1002	-	-	1/1/1/1	-
3	EDO	C	403	-	-	1/1/1/1	-
3	EDO	A	404	-	-	1/1/1/1	-
3	EDO	A	407	-	-	1/1/1/1	-
3	EDO	A	405	-	-	1/1/1/1	-
3	EDO	C	402	-	-	1/1/1/1	-
3	EDO	A	402	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1001	PEG	O2-C3-C4-O4
3	A	404	EDO	O1-C1-C2-O2
3	C	402	EDO	O1-C1-C2-O2
3	B	1002	EDO	O1-C1-C2-O2
3	C	403	EDO	O1-C1-C2-O2
2	A	401	PEG	C4-C3-O2-C2
3	A	402	EDO	O1-C1-C2-O2
2	A	401	PEG	O2-C3-C4-O4
3	A	406	EDO	O1-C1-C2-O2
2	A	401	PEG	C1-C2-O2-C3
3	A	405	EDO	O1-C1-C2-O2
3	A	407	EDO	O1-C1-C2-O2
2	B	1001	PEG	C1-C2-O2-C3
2	A	401	PEG	O1-C1-C2-O2
3	B	1003	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1001	PEG	1	0
3	B	1002	EDO	2	0
3	A	405	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	360/395 (91%)	-0.17	5 (1%) 75 83	20, 31, 65, 106	0
1	B	365/395 (92%)	-0.15	3 (0%) 86 91	18, 30, 60, 89	0
1	C	362/395 (91%)	-0.12	6 (1%) 70 78	21, 35, 64, 87	0
1	D	355/395 (89%)	-0.06	5 (1%) 75 83	22, 35, 68, 113	0
All	All	1442/1580 (91%)	-0.13	19 (1%) 77 84	18, 33, 66, 113	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	96	ALA	4.7
1	D	250	PHE	3.7
1	A	98	GLY	3.7
1	D	251	VAL	3.0
1	C	339	GLN	2.9
1	B	95	SER	2.8
1	A	211	GLU	2.8
1	B	94	ASP	2.7
1	D	215	LEU	2.6
1	A	212	ASP	2.6
1	C	192	ASN	2.5
1	C	98	GLY	2.4
1	C	395	PHE	2.1
1	D	104	CYS	2.1
1	C	228	ARG	2.1
1	A	99	PHE	2.1
1	C	95	SER	2.1
1	A	250	PHE	2.1
1	D	93	ASP	2.0



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	EDO	A	406	4/4	0.75	0.22	58,61,64,66	0
3	EDO	A	405	4/4	0.81	0.26	47,57,59,62	0
3	EDO	A	404	4/4	0.82	0.17	58,59,62,62	0
2	PEG	B	1001	7/7	0.85	0.23	35,40,42,45	0
2	PEG	A	401	7/7	0.85	0.26	56,60,63,63	0
3	EDO	C	401	4/4	0.88	0.14	59,59,59,59	0
3	EDO	B	1003	4/4	0.91	0.17	38,38,39,41	0
3	EDO	B	1002	4/4	0.91	0.18	37,38,38,44	0
3	EDO	A	407	4/4	0.93	0.19	39,40,41,44	0
3	EDO	C	403	4/4	0.93	0.24	44,44,45,47	0
3	EDO	C	402	4/4	0.94	0.16	38,39,39,44	0
3	EDO	A	403	4/4	0.94	0.17	38,41,42,44	0
3	EDO	A	402	4/4	0.95	0.18	27,28,30,30	0
3	EDO	D	401	4/4	0.97	0.10	28,31,33,34	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.