



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 24, 2024 – 12:04 AM EDT

PDB ID : 1TJW
Title : Crystal Structure of T161D Duck Delta 2 Crystallin Mutant with bound argininosuccinate
Authors : Sampaleanu, L.M.; Coddling, P.W.; Lobsanov, Y.D.; Tsai, M.; Smith, G.D.; Horvatin, C.; Howell, P.L.
Deposited on : 2004-06-07
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.002 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.38.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	474	
1	B	474	
1	C	474	
1	D	474	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	AS1	A	1004	X	-	-	-
2	AS1	B	1003	X	-	-	-
2	AS1	C	1001	X	-	-	-
2	AS1	D	1002	X	-	-	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 14819 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Delta crystallin II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	449	Total	C	N	O	S	0	0	0
			3464	2195	588	669	12			
1	B	448	Total	C	N	O	S	0	0	0
			3441	2179	581	669	12			
1	C	449	Total	C	N	O	S	0	0	0
			3456	2190	585	669	12			
1	D	449	Total	C	N	O	S	0	0	0
			3468	2194	586	676	12			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	161	ASP	THR	engineered mutation	UNP P24058
B	161	ASP	THR	engineered mutation	UNP P24058
C	161	ASP	THR	engineered mutation	UNP P24058
D	161	ASP	THR	engineered mutation	UNP P24058
A	469	HIS	-	expression tag	UNP P24058
A	470	HIS	-	expression tag	UNP P24058
A	471	HIS	-	expression tag	UNP P24058
A	472	HIS	-	expression tag	UNP P24058
A	473	HIS	-	expression tag	UNP P24058
A	474	HIS	-	expression tag	UNP P24058
B	469	HIS	-	expression tag	UNP P24058
B	470	HIS	-	expression tag	UNP P24058
B	471	HIS	-	expression tag	UNP P24058
B	472	HIS	-	expression tag	UNP P24058
B	473	HIS	-	expression tag	UNP P24058
B	474	HIS	-	expression tag	UNP P24058
C	469	HIS	-	expression tag	UNP P24058
C	470	HIS	-	expression tag	UNP P24058
C	471	HIS	-	expression tag	UNP P24058
C	472	HIS	-	expression tag	UNP P24058
C	473	HIS	-	expression tag	UNP P24058

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	474	HIS	-	expression tag	UNP P24058
D	469	HIS	-	expression tag	UNP P24058
D	470	HIS	-	expression tag	UNP P24058
D	471	HIS	-	expression tag	UNP P24058
D	472	HIS	-	expression tag	UNP P24058
D	473	HIS	-	expression tag	UNP P24058
D	474	HIS	-	expression tag	UNP P24058

- Molecule 2 is ARGININOSUCCINATE (three-letter code: AS1) (formula: C₁₀H₁₈N₄O₆).



Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	258	Total	O	0	0
			258	258		
3	D	203	Total	O	0	0
			203	203		



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	94.00Å 98.81Å 106.39Å 90.00° 101.34° 90.00°	Depositor
Resolution (Å)	41.76 – 2.00 41.76 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.5 (41.76-2.00) 99.6 (41.76-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.37 (at 2.00Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.190 , 0.225 0.182 , 0.217	Depositor DCC
R_{free} test set	12767 reflections (9.95%)	wwPDB-VP
Wilson B-factor (Å ²)	19.4	Xtriage
Anisotropy	0.479	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 56.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	14819	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AS1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/3509	0.62	0/4733
1	B	0.41	0/3486	0.60	0/4708
1	C	0.44	0/3501	0.62	0/4727
1	D	0.43	0/3513	0.61	0/4743
All	All	0.43	0/14009	0.61	0/18911

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	2	0
1	D	1	0
All	All	3	0

There are no bond length outliers.

There are no bond angle outliers.

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	B	105	ILE	CB
1	B	405	ILE	CB
1	D	105	ILE	CB

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3464	0	3570	64	0
1	B	3441	0	3512	76	0
1	C	3456	0	3545	72	0
1	D	3468	0	3547	75	0
2	A	20	0	14	1	0
2	B	20	0	14	0	0
2	C	20	0	14	1	0
2	D	20	0	14	1	0
3	A	253	0	0	8	0
3	B	196	0	0	5	0
3	C	258	0	0	8	0
3	D	203	0	0	5	0
All	All	14819	0	14230	276	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 276 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:102:ILE:HD11	1:B:106:ALA:HB2	1.42	1.00
1:A:110:HIS:HD2	1:A:113:ARG:HE	1.16	0.93
1:A:30:ILE:O	1:A:34:GLN:HG3	1.69	0.93
1:C:50:ALA:HB1	1:C:213:ILE:HD11	1.50	0.92
1:D:110:HIS:HD2	1:D:113:ARG:HE	1.17	0.92

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	447/474 (94%)	440 (98%)	6 (1%)	1 (0%)	44	42
1	B	446/474 (94%)	438 (98%)	5 (1%)	3 (1%)	19	14
1	C	447/474 (94%)	439 (98%)	7 (2%)	1 (0%)	44	42
1	D	447/474 (94%)	437 (98%)	9 (2%)	1 (0%)	44	42
All	All	1787/1896 (94%)	1754 (98%)	27 (2%)	6 (0%)	37	35

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	206	LEU
1	B	206	LEU
1	C	206	LEU
1	B	466	GLU
1	D	206	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	386/411 (94%)	374 (97%)	12 (3%)	35	36
1	B	380/411 (92%)	368 (97%)	12 (3%)	34	35
1	C	383/411 (93%)	371 (97%)	12 (3%)	35	36
1	D	386/411 (94%)	372 (96%)	14 (4%)	30	30
All	All	1535/1644 (93%)	1485 (97%)	50 (3%)	33	33

5 of 50 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	201	LEU
1	C	457	LEU
1	D	457	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	213	ILE
1	C	264	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	390	HIS
1	C	209	ASN
1	D	230	ASN
1	C	173	GLN
1	C	230	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	AS1	B	1003	-	18,19,19	1.02	0	19,24,24	1.34	2 (10%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	AS1	C	1001	-	18,19,19	1.02	1 (5%)	19,24,24	1.37	2 (10%)
2	AS1	D	1002	-	18,19,19	1.08	0	19,24,24	1.25	2 (10%)
2	AS1	A	1004	-	18,19,19	1.05	0	19,24,24	1.57	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AS1	B	1003	-	1/1/6/7	10/21/23/23	-
2	AS1	C	1001	-	1/1/6/7	10/21/23/23	-
2	AS1	D	1002	-	1/1/6/7	10/21/23/23	-
2	AS1	A	1004	-	1/1/6/7	12/21/23/23	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1001	AS1	C-N2	2.01	1.33	1.28

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1004	AS1	CG-CA-N1	4.74	121.56	110.57
2	B	1003	AS1	CG-CA-N1	3.63	119.00	110.57
2	C	1001	AS1	CG-CA-N1	3.35	118.35	110.57
2	A	1004	AS1	N1-C-N2	-3.09	117.50	125.00
2	B	1003	AS1	N1-C-N2	-2.87	118.06	125.00

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	1004	AS1	CA
2	B	1003	AS1	CA
2	C	1001	AS1	CA
2	D	1002	AS1	CA

5 of 42 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1004	AS1	N3-C-N1-CA
2	A	1004	AS1	CG-CA-N1-C
2	A	1004	AS1	N1-CA-CB-CD
2	B	1003	AS1	N3-C-N1-CA
2	B	1003	AS1	CG-CA-N1-C



There are no ring outliers.


3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1001	AS1	1	0
2	D	1002	AS1	1	0
2	A	1004	AS1	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	449/474 (94%)	-0.28	2 (0%) 89 88	10, 19, 41, 55	0
1	B	448/474 (94%)	-0.06	5 (1%) 77 76	10, 24, 40, 56	0
1	C	449/474 (94%)	-0.28	3 (0%) 84 83	10, 19, 36, 57	0
1	D	449/474 (94%)	-0.17	3 (0%) 84 83	11, 21, 37, 50	0
All	All	1795/1896 (94%)	-0.20	13 (0%) 84 83	10, 21, 39, 57	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	468	ALA	7.3
1	D	467	GLN	3.2
1	B	407	ILE	3.1
1	B	467	GLN	3.0
1	A	467	GLN	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	AS1	D	1002	20/20	0.88	0.11	24,31,38,38	0
2	AS1	B	1003	20/20	0.90	0.11	15,23,35,37	0
2	AS1	C	1001	20/20	0.91	0.09	18,30,36,36	0
2	AS1	A	1004	20/20	0.92	0.08	18,24,30,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around AS1 B 1003:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around AS1 C 1001:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.