



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 12, 2024 – 05:33 PM EST

PDB ID : 3UDL
Title : 3-heterocyclyl quinolone bound to HCV NS5B
Authors : Somoza, J.R.
Deposited on : 2011-10-28
Resolution : 2.17 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	8336 (2.20-2.16)
Clashscore	180529	9404 (2.20-2.16)
Ramachandran outliers	177936	9297 (2.20-2.16)
Sidechain outliers	177891	9297 (2.20-2.16)
RSRZ outliers	164620	8337 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	576	
1	B	576	
1	C	576	
1	D	576	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 18412 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HCV NS5B polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	555	Total	C	N	O	S	0	0	0
			4319	2725	761	802	31			
1	B	555	Total	C	N	O	S	0	0	0
			4319	2725	761	802	31			
1	C	555	Total	C	N	O	S	0	0	0
			4319	2725	761	802	31			
1	D	555	Total	C	N	O	S	0	0	0
			4319	2725	761	802	31			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	expression tag	UNP Q99AU2
A	-4	HIS	-	expression tag	UNP Q99AU2
A	-3	HIS	-	expression tag	UNP Q99AU2
A	-2	HIS	-	expression tag	UNP Q99AU2
A	-1	HIS	-	expression tag	UNP Q99AU2
A	0	HIS	-	expression tag	UNP Q99AU2
B	-5	HIS	-	expression tag	UNP Q99AU2
B	-4	HIS	-	expression tag	UNP Q99AU2
B	-3	HIS	-	expression tag	UNP Q99AU2
B	-2	HIS	-	expression tag	UNP Q99AU2
B	-1	HIS	-	expression tag	UNP Q99AU2
B	0	HIS	-	expression tag	UNP Q99AU2
C	-5	HIS	-	expression tag	UNP Q99AU2
C	-4	HIS	-	expression tag	UNP Q99AU2
C	-3	HIS	-	expression tag	UNP Q99AU2
C	-2	HIS	-	expression tag	UNP Q99AU2
C	-1	HIS	-	expression tag	UNP Q99AU2
C	0	HIS	-	expression tag	UNP Q99AU2
D	-5	HIS	-	expression tag	UNP Q99AU2
D	-4	HIS	-	expression tag	UNP Q99AU2
D	-3	HIS	-	expression tag	UNP Q99AU2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-2	HIS	-	expression tag	UNP Q99AU2
D	-1	HIS	-	expression tag	UNP Q99AU2
D	0	HIS	-	expression tag	UNP Q99AU2

- Molecule 2 is 3-(5-benzyl-1,2,4-oxadiazol-3-yl)-6-fluoro-1-[2-fluoro-4-(trifluoromethyl)benzyl]-7-(4-methylpiperazin-1-yl)quinolin-4(1H)-one (three-letter code: KLI) (formula: $C_{31}H_{26}F_5N_5O_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			43	31	5	5	2		
2	B	1	Total	C	F	N	O	0	0
			43	31	5	5	2		
2	C	1	Total	C	F	N	O	0	0
			43	31	5	5	2		
2	D	1	Total	C	F	N	O	0	0
			43	31	5	5	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	274	Total	O	0	0
			274	274		
3	B	248	Total	O	0	0
			248	248		
3	C	231	Total	O	0	0
			231	231		


Continued on next page...

Continued from previous page...



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	211	Total 211	O 211	0	0

- Molecule 1: HCV NS5B polymerase





• Molecule 1: HCV NS5B polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	101.75Å 101.86Å 250.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.91 – 2.17 49.91 – 2.17	Depositor EDS
% Data completeness (in resolution range)	92.9 (49.91-2.17) 92.4 (49.91-2.17)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 2.18Å)	Xtriage
Refinement program	PHENIX 1.7 _650	Depositor
R, R_{free}	0.191 , 0.243 0.186 , 0.235	Depositor DCC
R_{free} test set	13245 reflections (9.95%)	wwPDB-VP
Wilson B-factor (Å ²)	30.4	Xtriage
Anisotropy	0.367	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 30.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.087 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	18412	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.64 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.3176e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: KLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.22	0/4412	0.41	0/5985
1	B	0.23	0/4412	0.41	0/5985
1	C	0.23	0/4412	0.40	0/5985
1	D	0.22	0/4412	0.41	0/5985
All	All	0.23	0/17648	0.41	0/23940

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4319	0	4334	39	0
1	B	4319	0	4336	63	0
1	C	4319	0	4334	57	0
1	D	4319	0	4334	60	0
2	A	43	0	26	0	0
2	B	43	0	26	2	0
2	C	43	0	26	0	0
2	D	43	0	26	1	0
3	A	274	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	248	0	0	2	0
3	C	231	0	0	1	0
3	D	211	0	0	1	0
All	All	18412	0	17442	220	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 220 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:71:MET:HE1	1:D:297:LEU:HD13	1.49	0.94
1:D:268:ASN:HD21	1:D:272:GLN:HE21	1.18	0.91
1:B:187:MET:HE3	1:B:293:LEU:HA	1.60	0.83
1:B:187:MET:HE3	1:B:293:LEU:HD23	1.60	0.83
1:C:527:ASN:HD21	1:C:534:LEU:H	1.23	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	549/576 (95%)	533 (97%)	15 (3%)	1 (0%)	44	49
1	B	549/576 (95%)	538 (98%)	11 (2%)	0	100	100
1	C	549/576 (95%)	534 (97%)	15 (3%)	0	100	100
1	D	549/576 (95%)	531 (97%)	18 (3%)	0	100	100
All	All	2196/2304 (95%)	2136 (97%)	59 (3%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	351	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	474/491 (96%)	453 (96%)	21 (4%)	24	28
1	B	474/491 (96%)	449 (95%)	25 (5%)	19	21
1	C	474/491 (96%)	459 (97%)	15 (3%)	34	42
1	D	474/491 (96%)	454 (96%)	20 (4%)	25	30
All	All	1896/1964 (96%)	1815 (96%)	81 (4%)	25	29

5 of 81 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	469	LEU
1	D	372	VAL
1	C	547	LEU
1	D	182	LEU
1	D	431	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 40 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	514	GLN
1	D	436	GLN
1	C	527	ASN
1	D	251	GLN
1	D	475	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	KLI	B	571	-	42,48,48	1.29	2 (4%)	58,71,71	1.19	6 (10%)
2	KLI	D	571	-	42,48,48	1.27	2 (4%)	58,71,71	1.17	8 (13%)
2	KLI	C	571	-	42,48,48	1.25	2 (4%)	58,71,71	1.14	6 (10%)
2	KLI	A	571	-	42,48,48	1.26	2 (4%)	58,71,71	1.21	5 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	KLI	B	571	-	-	0/17/32/32	0/6/6/6
2	KLI	D	571	-	-	3/17/32/32	0/6/6/6
2	KLI	C	571	-	-	3/17/32/32	0/6/6/6
2	KLI	A	571	-	-	1/17/32/32	0/6/6/6

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	571	KLI	C12-N01	3.95	1.39	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	571	KLI	C12-N01	3.83	1.39	1.34
2	D	571	KLI	C12-N01	3.82	1.39	1.34
2	C	571	KLI	C12-N01	3.66	1.39	1.34
2	C	571	KLI	C18-C23	2.29	1.42	1.38

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	571	KLI	C18-C17-N01	-3.78	107.34	113.08
2	B	571	KLI	C11-C06-C07	-3.16	120.56	123.35
2	B	571	KLI	F43-C06-C07	3.14	121.36	118.37
2	D	571	KLI	C18-C17-N01	-3.06	108.44	113.08
2	D	571	KLI	C12-C13-C14	-3.05	117.06	120.46

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	571	KLI	C06-C07-N04-C28
2	D	571	KLI	C06-C07-N04-C28
2	C	571	KLI	C06-C07-N04-C25
2	D	571	KLI	C06-C07-N04-C25
2	C	571	KLI	C08-C07-N04-C28


There are no ring outliers.

2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	571	KLI	2	0
2	D	571	KLI	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient


equivalents in the CSD to analyse the geometry.



Ligand KLI C 571



Ligand KLI A 571



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	555/576 (96%)	-0.09	12 (2%) 62 60	21, 33, 57, 82	0
1	B	555/576 (96%)	-0.00	12 (2%) 62 60	21, 33, 59, 80	0
1	C	555/576 (96%)	0.00	11 (1%) 64 63	21, 34, 59, 85	0
1	D	555/576 (96%)	0.06	16 (2%) 54 53	20, 36, 61, 85	0
All	All	2220/2304 (96%)	-0.01	51 (2%) 61 59	20, 34, 59, 85	0

The worst 5 of 51 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	563	SER	4.6
1	C	563	SER	4.5
1	B	148	GLN	4.1
1	C	26	LEU	3.8
1	B	563	SER	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	KLI	B	571	43/43	0.88	0.11	30,46,55,66	0
2	KLI	A	571	43/43	0.89	0.10	27,41,49,55	0
2	KLI	C	571	43/43	0.90	0.10	30,42,53,61	0
2	KLI	D	571	43/43	0.92	0.09	27,41,52,61	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around KLI B 571:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around KLI A 571:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around KLI C 571:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around KLI D 571:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.