



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 8, 2024 – 04:00 AM EDT

PDB ID : 3V4N
Title : The Biochemical and Structural Basis for Inhibition of Enterococcus faecalis HMG-CoA Synthetase, mvaS, by Hymeglusin
Authors : Skaff, D.A.; Ramyar, K.X.; McWhorter, W.J.; Geisbrecht, B.V.; Mizioro, H.M.
Deposited on : 2011-12-15
Resolution : 1.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	4274 (1.60-1.60)
Clashscore	180529	4682 (1.60-1.60)
Ramachandran outliers	177936	4583 (1.60-1.60)
Sidechain outliers	177891	4582 (1.60-1.60)
RSRZ outliers	164620	4272 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	<div> <div>91%</div> <div>8%</div> </div>
1	B	388	<div> <div>93%</div> <div>6%</div> </div>
1	C	388	<div> <div>90%</div> <div>9%</div> </div>
1	D	388	<div> <div>91%</div> <div>8%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 13483 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HMG-CoA synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	387	Total	C	N	O	S	0	7	0
			3031	1928	498	593	12			
1	B	384	Total	C	N	O	S	0	3	0
			2990	1897	493	589	11			
1	C	383	Total	C	N	O	S	0	2	0
			2979	1892	492	584	11			
1	D	383	Total	C	N	O	S	0	1	0
			2975	1888	492	584	11			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP Q9FD71
A	-3	SER	-	expression tag	UNP Q9FD71
A	-2	THR	-	expression tag	UNP Q9FD71
A	-1	GLY	-	expression tag	UNP Q9FD71
A	0	SER	-	expression tag	UNP Q9FD71
A	163	SER	ALA	engineered mutation	UNP Q9FD71
B	-4	GLY	-	expression tag	UNP Q9FD71
B	-3	SER	-	expression tag	UNP Q9FD71
B	-2	THR	-	expression tag	UNP Q9FD71
B	-1	GLY	-	expression tag	UNP Q9FD71
B	0	SER	-	expression tag	UNP Q9FD71
B	163	SER	ALA	engineered mutation	UNP Q9FD71
C	-4	GLY	-	expression tag	UNP Q9FD71
C	-3	SER	-	expression tag	UNP Q9FD71
C	-2	THR	-	expression tag	UNP Q9FD71
C	-1	GLY	-	expression tag	UNP Q9FD71
C	0	SER	-	expression tag	UNP Q9FD71
C	163	SER	ALA	engineered mutation	UNP Q9FD71
D	-4	GLY	-	expression tag	UNP Q9FD71
D	-3	SER	-	expression tag	UNP Q9FD71
D	-2	THR	-	expression tag	UNP Q9FD71

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-1	GLY	-	expression tag	UNP Q9FD71
D	0	SER	-	expression tag	UNP Q9FD71
D	163	SER	ALA	engineered mutation	UNP Q9FD71

- Molecule 2 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (three-letter code: BTB) (formula: C₈H₁₉NO₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	536	Total	O	0	0
			536	536		
3	B	483	Total	O	0	0
			483	483		
3	C	222	Total	O	0	0
			222	222		
3	D	253	Total	O	0	0
			253	253		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HMG-CoA synthase




- Molecule 1: HMG-CoA synthase



- Molecule 1: HMG-CoA synthase



- Molecule 1: HMG-CoA synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	109.53Å 56.28Å 123.89Å 90.00° 100.02° 90.00°	Depositor
Resolution (Å)	26.96 – 1.60 26.96 – 1.60	Depositor EDS
% Data completeness (in resolution range)	98.3 (26.96-1.60) 94.8 (26.96-1.60)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.01 (at 1.60Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.2_869)	Depositor
R, R_{free}	0.141 , 0.194 0.145 , 0.198	Depositor DCC
R_{free} test set	19347 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	16.8	Xtriage
Anisotropy	0.602	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 44.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	13483	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BTB, SNC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	0/3103	0.43	0/4204
1	B	0.24	0/3047	0.43	0/4131
1	C	0.22	0/3033	0.41	0/4112
1	D	0.23	0/3026	0.42	0/4102
All	All	0.23	0/12209	0.42	0/16549

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3031	0	3019	29	0
1	B	2990	0	2948	24	0
1	C	2979	0	2942	17	0
1	D	2975	0	2933	25	0
2	B	14	0	19	3	0
3	A	536	0	0	7	1
3	B	483	0	0	6	0
3	C	222	0	0	1	0
3	D	253	0	0	8	0
All	All	13483	0	11861	95	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 95 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:324:GLN:O	3:D:571:HOH:O	1.99	0.79
1:B:341:GLU:OE2	3:B:974:HOH:O	2.03	0.77
1:A:9[A]:SER:OG	3:A:899:HOH:O	2.07	0.73
1:B:0:SER:N	3:B:949:HOH:O	2.19	0.71
1:A:116:ALA:O	1:A:120:LEU:HG	1.93	0.67

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:855:HOH:O	3:A:856:HOH:O[2_645]	2.14	0.06

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/388 (101%)	381 (97%)	11 (3%)	0	100	100
1	B	384/388 (99%)	376 (98%)	8 (2%)	0	100	100
1	C	382/388 (98%)	368 (96%)	14 (4%)	0	100	100
1	D	381/388 (98%)	374 (98%)	7 (2%)	0	100	100
All	All	1539/1552 (99%)	1499 (97%)	40 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	323/315 (102%)	320 (99%)	3 (1%)	75	62
1	B	316/315 (100%)	314 (99%)	2 (1%)	84	74
1	C	314/315 (100%)	309 (98%)	5 (2%)	58	37
1	D	313/315 (99%)	310 (99%)	3 (1%)	73	57
All	All	1266/1260 (100%)	1253 (99%)	13 (1%)	73	57

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	177	MET
1	C	257	GLU
1	D	383	ASN
1	D	109	GLU
1	D	276	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	191	HIS
1	D	326	HIS
1	A	211	GLN
1	B	326	HIS
1	C	326	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SNC	C	111	1	4,7,8	1.36	0	2,7,9	1.42	0
1	SNC	D	111	1	4,7,8	1.37	0	2,7,9	1.54	1 (50%)
1	SNC	B	111	1	4,7,8	1.27	0	2,7,9	1.24	0
1	SNC	A	111	1	4,7,8	1.41	0	2,7,9	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SNC	C	111	1	-	0/0/6/8	-
1	SNC	D	111	1	-	0/0/6/8	-
1	SNC	B	111	1	-	0/0/6/8	-
1	SNC	A	111	1	-	0/0/6/8	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	111	SNC	CA-CB-SG	2.09	117.27	112.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BTB	B	401	-	13,13,13	1.66	4 (30%)	7,16,16	2.12	4 (57%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BTB	B	401	-	-	7/21/21/21	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	BTB	C4-C2	-3.20	1.49	1.53
2	B	401	BTB	C1-C2	-3.07	1.49	1.53
2	B	401	BTB	C3-C2	-2.13	1.50	1.53
2	B	401	BTB	C5-N	-2.03	1.45	1.48

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	BTB	O3-C3-C2	3.13	118.76	111.40
2	B	401	BTB	O1-C1-C2	2.60	117.51	111.40
2	B	401	BTB	C6-C5-N	2.30	120.56	111.59
2	B	401	BTB	O4-C4-C2	2.16	116.49	111.40

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	BTB	C1-C2-C3-O3
2	B	401	BTB	C4-C2-C3-O3
2	B	401	BTB	N-C2-C3-O3
2	B	401	BTB	C1-C2-C4-O4
2	B	401	BTB	C3-C2-C4-O4

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	BTB	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	386/388 (99%)	-0.67	1 (0%) 90 91	7, 17, 36, 63	7 (1%)
1	B	383/388 (98%)	-0.71	0 100 100	8, 18, 31, 52	3 (0%)
1	C	382/388 (98%)	-0.00	1 (0%) 90 91	20, 36, 53, 63	2 (0%)
1	D	382/388 (98%)	-0.09	1 (0%) 90 91	16, 33, 51, 69	1 (0%)
All	All	1533/1552 (98%)	-0.37	3 (0%) 92 92	7, 26, 48, 69	13 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	98	ILE	2.5
1	C	254	ALA	2.0
1	A	-3	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SNC	A	111	8/9	0.97	0.07	7,12,29,42	0
1	SNC	B	111	8/9	0.97	0.07	10,13,34,47	0
1	SNC	C	111	8/9	0.97	0.06	22,29,40,41	0
1	SNC	D	111	8/9	0.98	0.05	19,25,31,46	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	BTB	B	401	14/14	0.82	0.16	31,41,63,72	0

6.5 Other polymers [i](#)

There are no such residues in this entry.