



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 19, 2025 – 08:11 PM JST

PDB ID : 9VMJ / pdb_00009vmj
Title : Crystal structure of TeGH116 from Thermosynechococcus elongatus
Authors : Beagbandee, C.; Charoenwattanasatien, R.; Pengthaisong, S.; Kurisu, G.; Ketudat Cairns, J.R.
Deposited on : 2025-06-28
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46

i

X-RAY DIFFRACTION

A.

the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	809	<div><div></div><div>88%</div><div>9%</div><div></div></div>
1	B	809	<div><div></div><div>88%</div><div>8%</div><div></div></div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 13185 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tlr0193 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	782	Total	C	N	O	S	0	0	0
			6213	3999	1055	1133	26			
1	B	785	Total	C	N	O	S	0	2	0
			6255	4026	1064	1139	26			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ALA	-	expression tag	UNP Q8DMC7
A	3	MET	-	expression tag	UNP Q8DMC7
A	4	ALA	-	expression tag	UNP Q8DMC7
B	2	ALA	-	expression tag	UNP Q8DMC7
B	3	MET	-	expression tag	UNP Q8DMC7
B	4	ALA	-	expression tag	UNP Q8DMC7

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Ca 1 1	0	0
3	B	1	Total Ca 1 1	0	0

- Molecule 4 is L(+)-TARTARIC ACID (CCD ID: TLA) (formula: C₄H₆O₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 10 4 6	0	0
4	B	1	Total C O 10 4 6	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	329	Total 329	O 329	0	0
5	B	342	Total 342	O 342	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tlr0193 protein



• Molecule 1: Tlr0193 protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	266.85Å 55.30Å 175.92Å 90.00° 129.72° 90.00°	Depositor
Resolution (Å)	26.66 – 2.20 26.66 – 2.20	Depositor EDS
% Data completeness (in resolution range)	96.8 (26.66-2.20) 96.8 (26.66-2.20)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.82 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.147 , 0.174 0.160 , 0.184	Depositor DCC
R_{free} test set	4966 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	21.9	Xtriage
Anisotropy	0.135	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 37.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.016 for -h-2*k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13185	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, TLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	1/6422 (0.0%)	1.22	13/8780 (0.1%)
1	B	0.88	2/6465 (0.0%)	1.24	20/8839 (0.2%)
All	All	0.87	3/12887 (0.0%)	1.23	33/17619 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	258	ASN	C-O	-5.75	1.20	1.25
1	B	765	ARG	NE-CZ	5.66	1.39	1.33
1	A	245	SER	CA-CB	-5.17	1.45	1.53

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	218	ASP	CA-CB-CG	8.03	120.63	112.60
1	A	709	ASN	CB-CA-C	-7.95	98.95	111.51
1	B	670	ASP	CA-CB-CG	7.59	120.19	112.60
1	B	648	GLN	CB-CA-C	-7.11	98.99	110.79
1	A	657	ARG	NE-CZ-NH1	-6.47	115.03	121.50
1	B	632	VAL	N-CA-CB	-6.42	102.44	110.47
1	A	584	ASP	CA-CB-CG	6.24	118.84	112.60
1	B	219	ASP	CA-CB-CG	6.17	118.77	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	685	ASP	CA-CB-CG	5.97	118.57	112.60
1	B	362	PHE	CA-CB-CG	-5.96	107.84	113.80
1	A	67	PHE	CA-CB-CG	5.91	119.70	113.80
1	B	476	ARG	NE-CZ-NH1	-5.91	115.59	121.50
1	B	253	ARG	NE-CZ-NH1	-5.84	115.66	121.50
1	B	648	GLN	N-CA-CB	5.83	118.69	110.12
1	A	701	ASP	CA-CB-CG	5.77	118.37	112.60
1	A	109	THR	CA-CB-OG1	-5.76	100.97	109.60
1	B	234	ASP	CB-CA-C	-5.75	100.93	109.90
1	B	497	GLU	CB-CG-CD	5.59	122.10	112.60
1	B	316	GLU	CB-CG-CD	5.53	122.00	112.60
1	B	794	PRO	CB-CA-C	-5.48	103.62	111.62
1	B	709	ASN	CA-CB-CG	-5.45	107.15	112.60
1	A	791	TYR	CA-C-N	5.44	128.01	120.29
1	A	791	TYR	C-N-CA	5.44	128.01	120.29
1	B	538	ASP	CA-CB-CG	5.41	118.01	112.60
1	A	344	PHE	CA-CB-CG	-5.36	108.44	113.80
1	A	94	THR	CA-CB-OG1	-5.34	101.59	109.60
1	A	641	GLN	N-CA-CB	5.30	117.98	110.13
1	B	645	ARG	CG-CD-NE	-5.21	100.54	112.00
1	B	765	ARG	CG-CD-NE	-5.18	100.61	112.00
1	B	557	THR	CA-CB-OG1	-5.14	101.89	109.60
1	B	253	ARG	NH1-CZ-NH2	5.05	125.87	119.30
1	A	372	THR	CA-CB-OG1	-5.03	102.05	109.60
1	B	145	PHE	CA-CB-CG	5.02	118.82	113.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	486	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6213	0	5862	31	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	6255	0	5908	24	0
2	A	18	0	24	0	0
2	B	6	0	8	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	10	0	4	1	0
4	B	10	0	4	1	0
5	A	329	0	0	0	0
5	B	342	0	0	0	0
All	All	13185	0	11810	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:724:PRO:HB3	1:B:733:LEU:HD21	1.64	0.80
1:A:47:ARG:NH2	4:A:905:TLA:O41	2.24	0.71
1:A:238:ASN:HB2	1:A:282:SER:O	2.02	0.58
1:A:206:CYS:O	1:A:228:PRO:HA	2.04	0.57
1:B:269:ALA:HA	1:B:339:VAL:O	2.04	0.57
1:B:238:ASN:HB2	1:B:282:SER:O	2.07	0.55
1:A:60:TYR:CE1	1:A:775:ARG:HD3	2.43	0.54
1:A:102:LEU:O	1:A:111:ALA:HA	2.07	0.54
1:B:47:ARG:NH2	4:B:903:TLA:O11	2.39	0.53
1:B:443:TYR:HB3	1:B:445:TRP:CZ2	2.43	0.53
1:B:206:CYS:O	1:B:228:PRO:HA	2.09	0.53
1:A:510:VAL:HG21	1:A:540:ALA:HA	1.91	0.52
1:B:47:ARG:O	1:B:785:THR:HA	2.10	0.52
1:B:409:LEU:HD11	1:B:801:ALA:HB2	1.94	0.50
1:B:618:GLU:OE2	1:B:643:ARG:NH1	2.45	0.49
1:A:269:ALA:HA	1:A:339:VAL:O	2.13	0.49
1:A:666:VAL:CG2	1:A:725:GLU:HB2	2.43	0.49
1:A:724:PRO:HB3	1:A:733:LEU:HD21	1.94	0.48
1:B:454:TYR:HB3	1:B:796:ALA:HB2	1.95	0.48
1:A:806:GLY:O	1:A:807:SER:C	2.55	0.48
1:A:793:ARG:N	1:A:794:PRO:HD2	2.29	0.48
1:A:575:HIS:O	1:A:578:GLN:HG2	2.13	0.47
1:A:47:ARG:O	1:A:785:THR:HA	2.14	0.47
1:B:194:LEU:O	1:B:323:ALA:HA	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:443:TYR:HB3	1:A:445:TRP:CZ2	2.49	0.47
1:B:664:SER:OG	1:B:666:VAL:HG12	2.15	0.46
1:A:48:VAL:HB	1:A:55:ASP:HB2	1.98	0.46
1:A:405:PHE:O	1:A:409:LEU:HB2	2.15	0.46
1:B:388:ILE:HA	1:B:391:TRP:CD1	2.52	0.45
1:A:537:LYS:HE3	1:A:579:PHE:CG	2.51	0.45
1:B:208:THR:OG1	1:B:227:THR:HB	2.16	0.45
1:B:592:ALA:O	1:B:593:PRO:C	2.61	0.44
1:B:512:HIS:ND1	1:B:528:THR:HG21	2.33	0.43
1:A:96:PRO:HB2	1:A:123:LEU:HD12	2.00	0.43
1:A:69:ALA:HB1	1:A:174:PRO:HB3	2.01	0.43
1:B:254:MET:HE3	1:B:254:MET:HB3	1.70	0.43
1:A:454:TYR:HB3	1:A:796:ALA:HB2	2.01	0.43
1:B:537:LYS:HE3	1:B:579:PHE:CG	2.54	0.43
1:A:726:ASN:OD1	1:A:726:ASN:C	2.62	0.42
1:A:667:ILE:O	1:A:717:GLY:HA2	2.19	0.42
1:A:807:SER:O	1:A:807:SER:OG	2.31	0.42
1:B:716:ASN:HD21	1:B:736:TRP:CD1	2.37	0.42
1:B:34:TRP:O	1:B:137:ALA:HA	2.20	0.42
1:A:46:TYR:CD1	1:A:770:ASN:HB3	2.54	0.41
1:A:467:GLU:OE2	1:A:550:ASP:OD2	2.38	0.41
1:A:447:GLU:O	1:A:448:SER:C	2.63	0.41
1:A:793:ARG:N	1:A:794:PRO:CD	2.83	0.41
1:B:765:ARG:NH1	1:B:765:ARG:HG2	2.35	0.41
1:A:324:LEU:HD23	1:A:336:ILE:HD13	2.02	0.41
1:B:792:LEU:HD12	1:B:795:MET:CE	2.51	0.41
1:A:592:ALA:O	1:A:593:PRO:C	2.63	0.41
1:B:767:ILE:HD13	1:B:776:THR:HG23	2.03	0.41
1:B:129:TYR:HA	1:B:130:PRO:HD3	1.94	0.41
1:A:401:TRP:O	1:A:406:LYS:HE3	2.21	0.40
1:A:733:LEU:HD23	1:A:733:LEU:HA	1.88	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	780/809 (96%)	753 (96%)	26 (3%)	1 (0%)	48	57
1	B	785/809 (97%)	752 (96%)	32 (4%)	1 (0%)	48	57
All	All	1565/1618 (97%)	1505 (96%)	58 (4%)	2 (0%)	48	57

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	806	GLY
1	B	775	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	632/650 (97%)	625 (99%)	7 (1%)	70	82
1	B	637/650 (98%)	628 (99%)	9 (1%)	62	77
All	All	1269/1300 (98%)	1253 (99%)	16 (1%)	65	78

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	116	SER
1	A	185	PRO
1	A	232	GLN
1	A	334	LYS
1	A	637	GLN
1	A	702	THR
1	A	807	SER
1	B	72	ILE
1	B	148	GLU
1	B	159	GLN
1	B	188	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	486	ARG
1	B	626	LEU
1	B	677	TYR
1	B	709	ASN
1	B	743	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	154	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	TLA	A	905	-	9,9,9	1.16	1 (11%)	12,12,12	1.30	1 (8%)
2	GOL	A	903	-	5,5,5	0.56	0	5,5,5	1.15	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	TLA	B	903	-	9,9,9	1.47	3 (33%)	12,12,12	1.12	0
2	GOL	B	901	-	5,5,5	0.28	0	5,5,5	0.61	0
2	GOL	A	901	-	5,5,5	0.16	0	5,5,5	0.46	0
2	GOL	A	902	-	5,5,5	0.25	0	5,5,5	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TLA	A	905	-	-	2/12/12/12	-
2	GOL	A	903	-	-	3/4/4/4	-
4	TLA	B	903	-	-	2/12/12/12	-
2	GOL	B	901	-	-	4/4/4/4	-
2	GOL	A	901	-	-	2/4/4/4	-
2	GOL	A	902	-	-	2/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	903	TLA	O41-C4	-2.51	1.22	1.30
4	A	905	TLA	O41-C4	-2.41	1.22	1.30
4	B	903	TLA	C2-C1	2.38	1.55	1.52
4	B	903	TLA	O11-C1	-2.02	1.23	1.30

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	905	TLA	O1-C1-C2	-2.14	116.00	121.63

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	901	GOL	C1-C2-C3-O3
2	A	902	GOL	C1-C2-C3-O3
2	A	903	GOL	O1-C1-C2-C3
2	B	901	GOL	C1-C2-C3-O3
2	A	902	GOL	O2-C2-C3-O3

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
2	A	903	GOL	O1-C1-C2-O2
2	A	903	GOL	C1-C2-C3-O3
2	B	901	GOL	O1-C1-C2-C3
2	B	901	GOL	O1-C1-C2-O2
2	B	901	GOL	O2-C2-C3-O3
2	A	901	GOL	O2-C2-C3-O3
4	B	903	TLA	O11-C1-C2-C3
4	A	905	TLA	C2-C3-C4-O41
4	B	903	TLA	O1-C1-C2-C3
4	A	905	TLA	C2-C3-C4-O4



There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	905	TLA	1	0
4	B	903	TLA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	782/809 (96%)	-0.68	2 (0%) 90 89	12, 22, 41, 76	0
1	B	785/809 (97%)	-0.70	3 (0%) 89 87	8, 21, 41, 74	2 (0%)
All	All	1567/1618 (96%)	-0.69	5 (0%) 90 89	8, 21, 41, 76	2 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	807	SER	2.7
1	B	214	ILE	2.2
1	B	809	LEU	2.1
1	B	810	PRO	2.1
1	A	213	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	A	901	6/6	0.89	0.10	30,31,34,40	0
2	GOL	A	902	6/6	0.89	0.12	39,42,44,48	0
4	TLA	A	905	10/10	0.90	0.10	33,39,52,59	0
2	GOL	A	903	6/6	0.91	0.11	27,35,41,43	0
4	TLA	B	903	10/10	0.91	0.09	39,45,52,54	0
2	GOL	B	901	6/6	0.92	0.12	29,38,41,46	0
3	CA	A	904	1/1	0.99	0.03	25,25,25,25	0
3	CA	B	902	1/1	0.99	0.04	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around GOL A 902:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around GOL A 903:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.