



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 26, 2024 – 09:04 AM EDT

PDB ID : 5VU8
Title : TNA polymerase, closed ternary complex
Authors : Chim, N.; Chaput, J.C.
Deposited on : 2017-05-18
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1370 (3.20-3.20)
Clashscore	180529	1497 (3.20-3.20)
Ramachandran outliers	177936	1479 (3.20-3.20)
Sidechain outliers	177891	1478 (3.20-3.20)
RSRZ outliers	164620	1371 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	774	 3% 71% 25% ..
2	T	16	 6% 50% 38% 12%
3	P	12	 67% 33%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6681 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	757	Total	C	N	O	S	0	0	0
			6120	3948	1026	1130	16			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	141	ALA	ASP	engineered mutation	UNP D0VWU9
A	143	ALA	GLU	engineered mutation	UNP D0VWU9
A	147	HIS	GLU	engineered mutation	UNP D0VWU9
A	485	ARG	ALA	engineered mutation	UNP D0VWU9
A	584	LYS	GLU	engineered mutation	UNP D0VWU9
A	664	ILE	GLU	engineered mutation	UNP D0VWU9


- Molecule 2 is a DNA chain called DNA template.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	T	14	Total	C	N	O	P	0	0	0
			286	136	50	86	14			

- Molecule 3 is a DNA chain called DNA/TNA hybrid primer.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	12	Total	C	N	O	P	0	0	0
			242	115	49	67	11			

- Molecule 4 is [(3 {S},4 {R},5 {R})-5-(6-aminopurin-9-yl)-4-oxidanyl-oxolan-3-yl] [oxidanyl(phosphonooxy)phosphoryl] hydrogen phosphate (three-letter code: 9O7) (formula: C₉H₁₄N₅O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			29	9	5	12	3		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	2	Total Mg 2 2	0	0
5	P	1	Total Mg 1 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total O 1 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase



- Molecule 2: DNA template



- Molecule 3: DNA/TNA hybrid primer



C1	C2	C3	C4	C10	C11	90412
----	----	----	----	-----	-----	-------

4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2 ₁	Depositor
Cell constants a, b, c, α , β , γ	72.29Å 107.36Å 147.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.42 – 3.20 43.42 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.6 (43.42-3.20) 87.6 (43.42-3.20)	Depositor EDS
R_{merge}	0.28	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.11 (at 2.96Å)	Xtriage
Refinement program	PHENIX (1.11.1 _2575: ???)	Depositor
R, R_{free}	0.258 , 0.295 0.260 , 0.297	Depositor DCC
R_{free} test set	17995 reflections (8.23%)	wwPDB-VP
Wilson B-factor (Å ²)	66.6	Xtriage
Anisotropy	0.618	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 26.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	6681	wwPDB-VP
Average B, all atoms (Å ²)	101.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 9O4, MG, 9O7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/6256	0.42	0/8452
2	T	0.52	0/319	0.92	0/490
3	P	0.49	0/250	0.77	0/384
All	All	0.27	0/6825	0.48	0/9326

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6120	0	6083	116	0
2	T	286	0	159	6	0
3	P	242	0	123	2	0
4	A	29	0	0	1	0
5	A	2	0	0	0	0
5	P	1	0	0	0	0
6	A	1	0	0	0	0
All	All	6681	0	6365	120	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (120) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:714:ALA:HA	1:A:715:ILE:HB	1.69	0.74
1:A:701:TYR:HA	1:A:716:PRO:HA	1.69	0.74
1:A:234:ARG:HH21	1:A:255:ARG:HH12	1.39	0.71
1:A:185:SER:HB2	1:A:189:GLU:HG3	1.72	0.70
1:A:685:ARG:NH2	1:A:714:ALA:O	2.24	0.70
1:A:592:LYS:NZ	2:T:7:DG:N3	2.42	0.68
1:A:428:CYS:HB2	1:A:431:TYR:HE2	1.60	0.66
1:A:472:ASP:HB3	1:A:475:GLU:HB3	1.77	0.66
1:A:756:ARG:HB3	1:A:757:TYR:HA	1.77	0.65
4:A:801:9O7:N6	2:T:4:DT:O4	2.29	0.65
1:A:564:LEU:O	1:A:568:ASN:ND2	2.30	0.64
1:A:661:VAL:HG22	1:A:702:ILE:HD11	1.80	0.63
1:A:241:ILE:HG12	1:A:250:VAL:HG22	1.81	0.63
1:A:670:LEU:HG	1:A:671:LYS:HG3	1.81	0.63
1:A:169:ARG:NH2	1:A:182:ASP:OD1	2.32	0.62
2:T:15:DC:H1'	2:T:16:DG:H8	1.66	0.60
1:A:426:GLU:OE2	1:A:440:ARG:NH2	2.37	0.58
1:A:730:GLU:O	1:A:732:TYR:N	2.37	0.58
1:A:15:VAL:HG22	1:A:32:ARG:HG2	1.86	0.57
1:A:616:SER:HB2	1:A:660:LEU:HA	1.87	0.57
1:A:709:ARG:HB3	1:A:711:GLY:N	2.20	0.56
1:A:48:ILE:HD11	1:A:68:VAL:HG11	1.87	0.56
1:A:153:ALA:O	1:A:222:ARG:NH1	2.38	0.56
1:A:202:ASP:HA	1:A:234:ARG:HH22	1.71	0.55
1:A:437:VAL:HG11	1:A:511:GLU:HG2	1.89	0.55
1:A:517:GLY:H	1:A:520:TYR:HD1	1.55	0.55
1:A:680:VAL:O	1:A:684:LYS:N	2.32	0.55
1:A:110:TYR:CG	1:A:367:LEU:HD21	2.42	0.54
1:A:554:GLU:OE1	1:A:554:GLU:N	2.40	0.54
1:A:6:ASP:OD1	1:A:7:TYR:N	2.39	0.54
1:A:435:PRO:O	1:A:437:VAL:N	2.40	0.54
1:A:122:ILE:HG23	1:A:359:ARG:HA	1.89	0.54
1:A:72:GLN:HG3	1:A:81:GLU:HG2	1.90	0.54
1:A:274:THR:OG1	1:A:276:GLU:OE1	2.26	0.53
1:A:126:LEU:O	1:A:359:ARG:NH2	2.41	0.53
1:A:401:VAL:HG22	1:A:582:PHE:HD1	1.75	0.52
1:A:14:PRO:HG3	1:A:90:PRO:HD3	1.92	0.52
1:A:202:ASP:OD1	1:A:234:ARG:NH1	2.39	0.52
1:A:649:LYS:HG3	1:A:652:LYS:HE3	1.91	0.52
1:A:669:ASP:O	1:A:671:LYS:N	2.43	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21:LYS:HE2	1:A:24:GLY:HA2	1.91	0.51
1:A:738:LEU:HB3	1:A:739:PRO:HD3	1.93	0.51
1:A:678:PRO:HA	1:A:711:GLY:HA2	1.94	0.50
1:A:260:LEU:HD21	1:A:323:GLY:HA2	1.92	0.50
1:A:340:SER:OG	1:A:343:ASP:OD2	2.29	0.50
1:A:184:VAL:HG21	1:A:190:MET:HB3	1.93	0.50
1:A:369:PRO:HG2	1:A:503:ARG:HB3	1.94	0.50
1:A:660:LEU:HD12	1:A:702:ILE:HD12	1.94	0.50
1:A:709:ARG:HB3	1:A:712:ASP:H	1.76	0.50
1:A:288:GLU:OE1	1:A:310:ARG:NH2	2.35	0.49
1:A:679:HIS:O	1:A:681:ALA:N	2.40	0.49
1:A:709:ARG:CB	1:A:712:ASP:H	2.26	0.49
1:A:21:LYS:HD2	1:A:204:ASP:OD2	2.13	0.48
1:A:188:ARG:HG3	1:A:228:ILE:HD11	1.95	0.48
1:A:384:TYR:HB2	1:A:511:GLU:HG3	1.95	0.48
1:A:392:PRO:HB3	1:A:538:TYR:HB2	1.95	0.48
1:A:267:THR:HG21	1:A:326:PHE:HE2	1.78	0.48
1:A:202:ASP:HA	1:A:234:ARG:NH2	2.28	0.48
1:A:721:ASP:OD2	1:A:724:LYS:N	2.39	0.48
1:A:137:MET:HB3	1:A:205:VAL:HB	1.95	0.47
1:A:208:THR:OG1	1:A:257:HIS:NE2	2.47	0.47
1:A:398:GLU:OE2	1:A:585:ARG:NH1	2.48	0.47
1:A:123:ASP:O	1:A:125:GLY:N	2.44	0.47
1:A:480:ASP:OD1	1:A:483:GLN:NE2	2.45	0.47
1:A:743:ARG:NH2	2:T:11:DT:OP1	2.48	0.47
1:A:400:ILE:HD11	1:A:596:VAL:HG23	1.96	0.47
1:A:564:LEU:HD11	1:A:579:TYR:HB2	1.97	0.47
1:A:597:ILE:HG22	1:A:603:ILE:HG12	1.95	0.47
1:A:597:ILE:HD13	1:A:631:LEU:HB3	1.96	0.47
2:T:15:DC:H1'	2:T:16:DG:C8	2.48	0.47
1:A:4:ASP:OD1	1:A:5:THR:N	2.43	0.46
1:A:52:LYS:HG2	1:A:68:VAL:HG21	1.97	0.46
1:A:334:SER:OG	1:A:339:GLN:O	2.30	0.46
1:A:525:ILE:HG23	1:A:536:VAL:HG21	1.97	0.46
1:A:618:ILE:HB	1:A:659:LYS:HD3	1.98	0.46
1:A:274:THR:HG23	1:A:277:ALA:H	1.80	0.46
1:A:93:VAL:HB	1:A:94:PRO:HD3	1.98	0.46
1:A:207:ILE:HD13	1:A:323:GLY:HA3	1.99	0.45
1:A:499:TYR:HB3	1:A:502:ALA:HB2	1.98	0.45
1:A:709:ARG:HA	1:A:710:ILE:CB	2.46	0.45
1:A:59:HIS:O	1:A:59:HIS:ND1	2.48	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:186:THR:OG1	1:A:187:GLU:N	2.50	0.45
1:A:384:TYR:HE1	1:A:514:THR:HG21	1.82	0.45
1:A:188:ARG:NH1	1:A:226:LEU:O	2.50	0.45
1:A:626:VAL:HG11	1:A:643:VAL:HG12	1.98	0.45
1:A:146:TYR:HB3	1:A:292:ALA:HB1	1.99	0.45
1:A:299:TRP:HA	1:A:305:LEU:HD21	1.99	0.45
1:A:751:ARG:O	1:A:752:LYS:HG2	2.17	0.45
1:A:70:LYS:HB2	1:A:83:TRP:CZ3	2.52	0.45
1:A:214:PHE:O	1:A:218:TYR:N	2.49	0.44
1:A:448:PHE:O	1:A:449:ILE:HB	2.17	0.44
1:A:419:SER:OG	1:A:421:ASP:OD2	2.35	0.44
1:A:551:ALA:HB1	1:A:555:THR:HB	2.00	0.44
1:A:220:LYS:O	1:A:224:GLU:HG2	2.18	0.44
1:A:640:VAL:O	1:A:643:VAL:HG22	2.17	0.44
3:P:3:DC:H2''	3:P:4:DG:C8	2.53	0.44
1:A:234:ARG:HB2	1:A:254:GLY:HA3	2.00	0.44
1:A:142:ILE:HG22	1:A:160:ILE:HG12	1.99	0.44
1:A:281:ALA:O	1:A:477:LYS:NZ	2.33	0.44
1:A:579:TYR:OH	1:A:582:PHE:HB2	2.18	0.44
1:A:207:ILE:HG21	1:A:323:GLY:HA3	1.99	0.43
1:A:145:LEU:HB2	1:A:158:LEU:HD21	2.01	0.43
1:A:123:ASP:C	1:A:125:GLY:H	2.21	0.42
1:A:733:ILE:HG23	1:A:734:GLU:HG3	2.02	0.42
1:A:361:ALA:O	1:A:366:GLU:N	2.51	0.42
1:A:108:ASP:OD1	1:A:109:ILE:N	2.51	0.42
1:A:662:ILE:HD11	1:A:701:TYR:CE2	2.55	0.42
1:A:212:ASP:HB3	1:A:249:ALA:HA	2.02	0.42
1:A:261:TYR:CE2	1:A:265:ARG:HD2	2.55	0.42
1:A:103:HIS:HB3	1:A:106:VAL:HG13	2.03	0.41
1:A:171:ILE:HG22	1:A:190:MET:HB2	2.01	0.41
1:A:612:ARG:HD2	3:P:10:DC:H5'	2.02	0.41
1:A:162:TYR:CZ	1:A:197:VAL:HG11	2.56	0.41
1:A:738:LEU:H	1:A:741:VAL:HG23	1.85	0.41
1:A:743:ARG:HD3	2:T:10:DG:OP1	2.21	0.41
1:A:261:TYR:HB3	1:A:262:PRO:HD3	2.03	0.40
1:A:16:ILE:HB	1:A:31:ASP:HB3	2.03	0.40
1:A:609:GLU:OE1	1:A:623:GLN:NE2	2.41	0.40
1:A:656:PRO:HA	1:A:657:PRO:HD3	1.83	0.40
1:A:97:ARG:HD2	1:A:112:TYR:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	751/774 (97%)	684 (91%)	56 (8%)	11 (2%)	8	38

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	449	ILE
1	A	670	LEU
1	A	738	LEU
1	A	567	ILE
1	A	710	ILE
1	A	716	PRO
1	A	755	LEU
1	A	66	LYS
1	A	245	GLY
1	A	436	GLN
1	A	715	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	636/673 (94%)	620 (98%)	16 (2%)	42	71

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	TYR
1	A	48	ILE
1	A	219	LEU
1	A	230	PHE
1	A	248	PHE
1	A	354	GLU
1	A	417	ASN
1	A	494	TYR
1	A	499	TYR
1	A	542	ASP
1	A	630	LEU
1	A	662	ILE
1	A	691	VAL
1	A	741	VAL
1	A	742	GLU
1	A	745	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	9O4	P	12	3,5	15,21,22	5.93	9 (60%)	15,30,33	2.98	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	9O4	P	12	3,5	-	1/1/17/18	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	12	9O4	C2-N3	14.41	1.54	1.32
3	P	12	9O4	C4-N3	10.92	1.50	1.35
3	P	12	9O4	C6-C5	-7.12	1.16	1.43
3	P	12	9O4	C8-N7	-7.06	1.21	1.34
3	P	12	9O4	C6-N1	-5.73	1.14	1.36
3	P	12	9O4	C2'-C3'	-4.11	1.44	1.52
3	P	12	9O4	O4'-C4'	3.99	1.52	1.43
3	P	12	9O4	O4'-C1'	3.57	1.48	1.41
3	P	12	9O4	C6-N6	3.44	1.46	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	12	9O4	C4-C5-N7	-7.59	101.32	109.34
3	P	12	9O4	C2'-C3'-C4'	4.84	109.27	102.76
3	P	12	9O4	C6-C5-C4	4.68	127.00	117.90
3	P	12	9O4	N3-C2-N1	-3.96	123.30	128.67

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	P	12	9O4	C2'-C3'-O3'-P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	9O7	A	801	5	26,31,31	3.26	14 (53%)	29,49,49	1.50	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	9O7	A	801	5	-	4/17/34/34	0/3/3/3

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	801	9O7	O2'-C2'	-9.69	1.19	1.43
4	A	801	9O7	PB-O3A	5.27	1.65	1.59
4	A	801	9O7	PA-O3A	5.22	1.65	1.59
4	A	801	9O7	PB-O3B	5.14	1.65	1.59
4	A	801	9O7	O4'-C1'	4.02	1.47	1.42
4	A	801	9O7	C5-N7	3.49	1.52	1.39
4	A	801	9O7	C6-N6	3.21	1.45	1.34
4	A	801	9O7	C2'-C3'	-2.80	1.45	1.53
4	A	801	9O7	C4-N3	-2.78	1.31	1.35
4	A	801	9O7	O4'-C4'	2.48	1.49	1.43
4	A	801	9O7	PG-O3G	-2.27	1.46	1.54
4	A	801	9O7	PG-O1G	-2.23	1.46	1.54
4	A	801	9O7	C1'-N9	-2.20	1.44	1.49
4	A	801	9O7	C2-N3	2.00	1.35	1.32

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	801	9O7	O1G-PG-O3B	2.86	114.24	104.64
4	A	801	9O7	C3'-C2'-C1'	2.85	106.19	99.92
4	A	801	9O7	O3G-PG-O3B	2.73	113.78	104.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	801	9O7	C1'-N9-C4	-2.70	121.90	126.64
4	A	801	9O7	O2A-PA-O1A	-2.40	101.30	112.44
4	A	801	9O7	O1B-PB-O2B	-2.36	101.45	112.44

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	801	9O7	C3'-O3'-PA-O2A
4	A	801	9O7	C3'-O3'-PA-O3A
4	A	801	9O7	PB-O3A-PA-O3'
4	A	801	9O7	C3'-O3'-PA-O1A

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	801	9O7	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	757/774 (97%)	0.15	23 (3%) 52 37	67, 92, 154, 271	0
2	T	14/16 (87%)	0.64	1 (7%) 23 16	83, 113, 171, 194	0
3	P	11/12 (91%)	0.23	0 100 100	113, 128, 139, 152	0
All	All	782/802 (97%)	0.16	24 (3%) 51 36	67, 92, 155, 271	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	614	ASP	5.9
1	A	657	PRO	3.7
1	A	715	ILE	3.6
1	A	688	ALA	3.5
1	A	110	TYR	3.3
1	A	727	TYR	3.1
1	A	80	VAL	3.1
1	A	75	PHE	3.0
1	A	728	ASP	3.0
1	A	740	ALA	2.7
1	A	490	ALA	2.5
1	A	567	ILE	2.5
1	A	662	ILE	2.3
1	A	753	GLU	2.3
2	T	16	DG	2.3
1	A	177	ASP	2.3
1	A	367	LEU	2.2
1	A	686	LEU	2.2
1	A	738	LEU	2.2
1	A	122	ILE	2.1
1	A	655	VAL	2.1
1	A	2	ILE	2.1
1	A	734	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	731	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	9O4	P	12	19/20	0.81	0.11	77,87,109,116	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MG	A	802	1/1	0.62	0.13	77,77,77,77	0
5	MG	A	803	1/1	0.62	0.20	61,61,61,61	1
5	MG	P	101	1/1	0.68	0.21	81,81,81,81	0
4	9O7	A	801	29/29	0.76	0.19	70,80,97,107	29

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 9O7 A 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.