wwPDB EM Validation Summary Report

Nov 5, 2025 – 12:52 PM JST

PDB ID	:	9VVR / pdb_00009vvr
EMDB ID	:	EMD-65388
Title	:	Structure of the bacteriophage E1004 tail
Authors	:	Sun, B.N.; Liu, H.R.
Deposited on	:	2025-07-16
Resolution	:	3.11 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev129
MolProbity	:	4-5-2 with Phenix2.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.46

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	210492	15764	-
Ramachandran outliers	207382	16835	-
Sidechain outliers	206894	16415	-
Q-score	-	25397	14465 (2.61 - 3.61)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	843	12% 84%
1	B	843	15% 83%
1	C	843	12% 86%
1	D	843	12% 84%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	843	14% 83%
1	F	843	13% 86%
1	G	843	12% 84%
1	H	843	12% 84%
1	I	843	12% 84%
1	J	843	14% 83%
1	K	843	16% 83%
1	L	843	15% 83%
1	M	843	12% 86%
1	N	843	12% 86%
1	O	843	12% 86%
1	P	843	12% 84%
1	Q	843	15% 83%
1	R	843	13% 86%
2	S	785	90% 8%
2	V	785	93% 7%
2	X	785	93% 7%
2	Z	785	91% 9%
2	b	785	91% 9%
2	d	785	90% 10%
3	T	188	89% 11%
3	U	188	90% 10%
3	W	188	90% 10%
3	Y	188	93% 6%
3	a	188	89% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	c	188	91% 9%
3	e	188	91% 9%
3	f	188	88% 11%
3	g	188	89% 11%
3	h	188	91% 8%
3	i	188	90% 9%
3	j	188	92% 7%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 72911 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tail spike protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	131	Total	C	N	O	S	0	0
			1017	629	178	208	2		
1	B	140	Total	C	N	O	S	0	0
			1092	672	193	225	2		
1	C	121	Total	C	N	O	S	0	0
			956	596	166	192	2		
1	D	131	Total	C	N	O	S	0	0
			1017	629	178	208	2		
1	E	140	Total	C	N	O	S	0	0
			1092	672	193	225	2		
1	F	121	Total	C	N	O	S	0	0
			956	596	166	192	2		
1	G	131	Total	C	N	O	S	0	0
			1017	629	178	208	2		
1	H	131	Total	C	N	O	S	0	0
			1017	629	178	208	2		
1	I	131	Total	C	N	O	S	0	0
			1017	629	178	208	2		
1	J	140	Total	C	N	O	S	0	0
			1092	672	193	225	2		
1	K	140	Total	C	N	O	S	0	0
			1092	672	193	225	2		
1	L	140	Total	C	N	O	S	0	0
			1092	672	193	225	2		
1	M	121	Total	C	N	O	S	0	0
			956	596	166	192	2		
1	N	121	Total	C	N	O	S	0	0
			956	596	166	192	2		
1	O	121	Total	C	N	O	S	0	0
			956	596	166	192	2		
1	P	131	Total	C	N	O	S	0	0
			1017	629	178	208	2		
1	Q	140	Total	C	N	O	S	0	0
			1092	672	193	225	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
1	R	121	Total	C	N	O	S	0	0
			956	596	166	192	2		

- Molecule 2 is a protein called Nozzle protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	S	776	Total	C	N	O	S	0	0
			6048	3831	1027	1177	13		
2	V	784	Total	C	N	O	S	0	0
			6101	3862	1036	1190	13		
2	X	784	Total	C	N	O	S	0	0
			6101	3862	1036	1190	13		
2	Z	784	Total	C	N	O	S	0	0
			6101	3862	1036	1190	13		
2	b	784	Total	C	N	O	S	0	0
			6101	3862	1036	1190	13		
2	d	780	Total	C	N	O	S	0	0
			6069	3842	1031	1183	13		

- Molecule 3 is a protein called Adaptor protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	T	187	Total	C	N	O	S	0	0
			1496	939	250	298	9		
3	U	188	Total	C	N	O	S	0	0
			1504	944	251	299	10		
3	W	188	Total	C	N	O	S	0	0
			1504	944	251	299	10		
3	Y	187	Total	C	N	O	S	0	0
			1496	939	250	298	9		
3	a	188	Total	C	N	O	S	0	0
			1504	944	251	299	10		
3	c	188	Total	C	N	O	S	0	0
			1504	944	251	299	10		
3	e	188	Total	C	N	O	S	0	0
			1504	944	251	299	10		
3	f	187	Total	C	N	O	S	0	0
			1496	939	250	298	9		
3	g	188	Total	C	N	O	S	0	0
			1504	944	251	299	10		
3	h	187	Total	C	N	O	S	0	0
			1496	939	250	298	9		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
3	i	187	Total	C	N	O	S	0	0
			1496	939	250	298	9		
3	j	187	Total	C	N	O	S	0	0
			1496	939	250	298	9		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tail spike protein

Chain A:



• Molecule 1: Tail spike protein

Chain B:



• Molecule 1: Tail spike protein

Chain C:





• Molecule 1: Tail spike protein

Chain D:





• Molecule 1: Tail spike protein

Chain E:





• Molecule 1: Tail spike protein

Chain F:



• Molecule 1: Tail spike protein

Chain G:





- Molecule 1: Tail spike protein

Chain H:







• Molecule 1: Tail spike protein

Chain I:



• Molecule 1: Tail spike protein

Chain J:



- Molecule 1: Tail spike protein

Chain K:





- Molecule 1: Tail spike protein

Chain L:







• Molecule 1: Tail spike protein

Chain M:



• Molecule 1: Tail spike protein

Chain N:



• Molecule 1: Tail spike protein



• Molecule 1: Tail spike protein

Chain P:





• Molecule 1: Tail spike protein

Chain Q:



• Molecule 1: Tail spike protein

Chain R:



• Molecule 2: Nozzle protein





• Molecule 2: Nozzle protein

Chain V:



• Molecule 2: Nozzle protein

Chain X:



• Molecule 2: Nozzle protein

Chain Z:



• Molecule 2: Nozzle protein

Chain b:



• Molecule 2: Nozzle protein

Chain d:



• Molecule 3: Adaptor protein

Chain T:



• Molecule 3: Adaptor protein

Chain U:



• Molecule 3: Adaptor protein

Chain W:



• Molecule 3: Adaptor protein

Chain Y:



• Molecule 3: Adaptor protein

Chain a:



• Molecule 3: Adaptor protein

Chain c:



• Molecule 3: Adaptor protein

Chain e:





• Molecule 3: Adaptor protein

Chain f:



• Molecule 3: Adaptor protein

Chain g:



• Molecule 3: Adaptor protein

Chain h:



• Molecule 3: Adaptor protein

Chain i:

• Molecule 3: Adaptor protein

Chain j:


92%
7%
MET
A2
L7
Q30
L31
D32
E33
T68
P69
D73
N74
T91
F111
E155
L156
Y157
G177
M182
D188

4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	50105	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	32	Depositor
Minimum defocus (nm)	1600	Depositor
Maximum defocus (nm)	2200	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	1.439	Depositor
Minimum map value	-0.778	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.057	Depositor
Recommended contour level	0.18	Depositor
Map size (Å)	484.0, 484.0, 484.0	wwPDB
Map dimensions	440, 440, 440	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.1, 1.1, 1.1	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	A	0.34	0/1031	0.56	0/1400
1	B	0.18	0/1106	0.40	0/1500
1	C	0.14	0/970	0.31	0/1319
1	D	0.28	0/1031	0.51	0/1400
1	E	0.30	0/1106	0.53	0/1500
1	F	0.15	0/970	0.34	0/1319
1	G	0.19	0/1031	0.46	0/1400
1	H	0.35	0/1031	0.60	0/1400
1	I	0.19	0/1031	0.48	0/1400
1	J	0.17	0/1106	0.37	0/1500
1	K	0.16	0/1106	0.36	0/1500
1	L	0.16	0/1106	0.39	0/1500
1	M	0.15	0/970	0.36	0/1319
1	N	0.16	0/970	0.38	0/1319
1	O	0.17	0/970	0.42	0/1319
1	P	0.19	0/1031	0.46	0/1400
1	Q	0.17	0/1106	0.39	0/1500
1	R	0.15	0/970	0.35	0/1319
2	S	0.16	0/6184	0.37	0/8396
2	V	0.15	0/6239	0.36	0/8474
2	X	0.15	0/6239	0.36	0/8474
2	Z	0.19	0/6239	0.41	0/8474
2	b	0.15	0/6239	0.36	0/8474
2	d	0.18	0/6204	0.41	0/8424
3	T	0.14	0/1525	0.36	0/2069
3	U	0.15	0/1533	0.41	0/2079
3	W	0.15	0/1533	0.39	0/2079
3	Y	0.15	0/1525	0.35	0/2069
3	a	0.16	0/1533	0.39	0/2079
3	c	0.15	0/1533	0.40	0/2079
3	e	0.15	0/1533	0.38	0/2079
3	f	0.15	0/1525	0.36	0/2069
3	g	0.15	0/1533	0.38	0/2079
3	h	0.15	0/1525	0.37	0/2069

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
3	i	0.16	0/1525	0.37	0/2069
3	j	0.15	0/1525	0.37	0/2069
All	All	0.17	0/74334	0.39	0/100918

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1017	0	986	23	0
1	B	1092	0	1055	11	0
1	C	956	0	932	11	0
1	D	1017	0	986	18	0
1	E	1092	0	1055	17	0
1	F	956	0	932	10	0
1	G	1017	0	986	20	0
1	H	1017	0	986	24	0
1	I	1017	0	986	26	0
1	J	1092	0	1055	14	0
1	K	1092	0	1055	6	0
1	L	1092	0	1055	10	0
1	M	956	0	932	13	0
1	N	956	0	932	11	0
1	O	956	0	932	15	0
1	P	1017	0	986	21	0
1	Q	1092	0	1055	11	0
1	R	956	0	932	10	0
2	S	6048	0	5900	41	0
2	V	6101	0	5949	32	0
2	X	6101	0	5949	31	0
2	Z	6101	0	5949	39	0
2	b	6101	0	5949	42	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	d	6069	0	5926	51	0
3	T	1496	0	1430	13	0
3	U	1504	0	1442	12	0
3	W	1504	0	1442	13	0
3	Y	1496	0	1430	15	0
3	a	1504	0	1442	15	0
3	c	1504	0	1442	12	0
3	e	1504	0	1442	12	0
3	f	1496	0	1430	19	0
3	g	1504	0	1442	15	0
3	h	1496	0	1430	12	0
3	i	1496	0	1430	11	0
3	j	1496	0	1430	10	0
All	All	72911	0	70692	613	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 613 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:d:630:LEU:HD11	2:d:661:VAL:HG11	1.16	1.09
2:d:630:LEU:CD1	2:d:661:VAL:HG11	1.85	1.05
2:d:661:VAL:HG12	2:d:662:PHE:N	1.87	0.87
2:d:661:VAL:HG12	2:d:662:PHE:H	1.42	0.83
1:A:32:ASN:HD22	1:A:38:LEU:HD23	1.44	0.81

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	129/843 (15%)	123 (95%)	5 (4%)	1 (1%)	16	46
1	B	138/843 (16%)	135 (98%)	3 (2%)	0	100	100
1	C	119/843 (14%)	118 (99%)	1 (1%)	0	100	100
1	D	129/843 (15%)	120 (93%)	8 (6%)	1 (1%)	16	46
1	E	138/843 (16%)	130 (94%)	8 (6%)	0	100	100
1	F	119/843 (14%)	117 (98%)	1 (1%)	1 (1%)	16	46
1	G	129/843 (15%)	124 (96%)	5 (4%)	0	100	100
1	H	129/843 (15%)	120 (93%)	9 (7%)	0	100	100
1	I	129/843 (15%)	125 (97%)	4 (3%)	0	100	100
1	J	138/843 (16%)	137 (99%)	1 (1%)	0	100	100
1	K	138/843 (16%)	133 (96%)	5 (4%)	0	100	100
1	L	138/843 (16%)	137 (99%)	1 (1%)	0	100	100
1	M	119/843 (14%)	117 (98%)	2 (2%)	0	100	100
1	N	119/843 (14%)	117 (98%)	2 (2%)	0	100	100
1	O	119/843 (14%)	114 (96%)	5 (4%)	0	100	100
1	P	129/843 (15%)	125 (97%)	4 (3%)	0	100	100
1	Q	138/843 (16%)	136 (99%)	2 (1%)	0	100	100
1	R	119/843 (14%)	118 (99%)	1 (1%)	0	100	100
2	S	772/785 (98%)	717 (93%)	51 (7%)	4 (0%)	25	56
2	V	782/785 (100%)	726 (93%)	53 (7%)	3 (0%)	30	61
2	X	782/785 (100%)	728 (93%)	51 (6%)	3 (0%)	30	61
2	Z	782/785 (100%)	726 (93%)	54 (7%)	2 (0%)	37	67
2	b	782/785 (100%)	725 (93%)	54 (7%)	3 (0%)	30	61
2	d	776/785 (99%)	716 (92%)	58 (8%)	2 (0%)	37	67
3	T	185/188 (98%)	175 (95%)	10 (5%)	0	100	100
3	U	186/188 (99%)	171 (92%)	15 (8%)	0	100	100
3	W	186/188 (99%)	176 (95%)	10 (5%)	0	100	100
3	Y	185/188 (98%)	176 (95%)	9 (5%)	0	100	100
3	a	186/188 (99%)	176 (95%)	10 (5%)	0	100	100
3	c	186/188 (99%)	170 (91%)	16 (9%)	0	100	100
3	e	186/188 (99%)	171 (92%)	15 (8%)	0	100	100
3	f	185/188 (98%)	170 (92%)	15 (8%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	g	186/188 (99%)	174 (94%)	12 (6%)	0	100	100
3	h	185/188 (98%)	175 (95%)	10 (5%)	0	100	100
3	i	185/188 (98%)	174 (94%)	11 (6%)	0	100	100
3	j	185/188 (98%)	174 (94%)	11 (6%)	0	100	100
All	All	9218/22140 (42%)	8666 (94%)	532 (6%)	20 (0%)	45	73

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	V	603	TYR
2	d	603	TYR
1	A	40	ARG
1	D	103	ILE
1	F	63	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/685 (16%)	112 (99%)	1 (1%)	75	87
1	B	120/685 (18%)	120 (100%)	0	100	100
1	C	107/685 (16%)	107 (100%)	0	100	100
1	D	113/685 (16%)	113 (100%)	0	100	100
1	E	120/685 (18%)	119 (99%)	1 (1%)	79	89
1	F	107/685 (16%)	107 (100%)	0	100	100
1	G	113/685 (16%)	113 (100%)	0	100	100
1	H	113/685 (16%)	112 (99%)	1 (1%)	75	87
1	I	113/685 (16%)	113 (100%)	0	100	100
1	J	120/685 (18%)	120 (100%)	0	100	100
1	K	120/685 (18%)	120 (100%)	0	100	100
1	L	120/685 (18%)	120 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	M	107/685 (16%)	107 (100%)	0	100	100
1	N	107/685 (16%)	107 (100%)	0	100	100
1	O	107/685 (16%)	107 (100%)	0	100	100
1	P	113/685 (16%)	113 (100%)	0	100	100
1	Q	120/685 (18%)	120 (100%)	0	100	100
1	R	107/685 (16%)	107 (100%)	0	100	100
2	S	663/670 (99%)	663 (100%)	0	100	100
2	V	669/670 (100%)	669 (100%)	0	100	100
2	X	669/670 (100%)	669 (100%)	0	100	100
2	Z	669/670 (100%)	668 (100%)	1 (0%)	92	97
2	b	669/670 (100%)	669 (100%)	0	100	100
2	d	666/670 (99%)	663 (100%)	3 (0%)	86	92
3	T	162/163 (99%)	162 (100%)	0	100	100
3	U	163/163 (100%)	163 (100%)	0	100	100
3	W	163/163 (100%)	163 (100%)	0	100	100
3	Y	162/163 (99%)	162 (100%)	0	100	100
3	a	163/163 (100%)	163 (100%)	0	100	100
3	c	163/163 (100%)	163 (100%)	0	100	100
3	e	163/163 (100%)	163 (100%)	0	100	100
3	f	162/163 (99%)	162 (100%)	0	100	100
3	g	163/163 (100%)	163 (100%)	0	100	100
3	h	162/163 (99%)	162 (100%)	0	100	100
3	i	162/163 (99%)	162 (100%)	0	100	100
3	j	162/163 (99%)	162 (100%)	0	100	100
All	All	7995/18306 (44%)	7988 (100%)	7 (0%)	92	97

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	Z	117	VAL
2	d	651	THR
2	d	665	ARG
2	d	660	THR
1	H	11	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 75 such sidechains are listed below:

Mol	Chain	Res	Type
2	d	334	ASN
3	i	60	ASN
2	d	634	GLN
3	a	145	ASN
2	S	311	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Map visualisation

This section contains visualisations of the EMDB entry EMD-65388. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections

6.1.1 Primary map



X



Y



Z

6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map



X Index: 220



Y Index: 220



Z Index: 220

6.2.2 Raw map



X Index: 220



Y Index: 220



Z Index: 220

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 206



Y Index: 196



Z Index: 184

6.3.2 Raw map



X Index: 235



Y Index: 196



Z Index: 184

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



X



Y



Z

6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map

X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.18. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency

A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

6.6.1 emd_65388_msk_1.map (i)

X



Y



Z

7 Map analysis

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate (i)



The volume at the recommended contour level is 1184 nm^3; this corresponds to an approximate mass of 1069 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.322 $Å^{-1}$

8 Fourier-Shell correlation

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC



*Reported resolution corresponds to spatial frequency of 0.322 Å^{-1}

8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.11	-	-
Author-provided FSC curve	3.11	3.62	3.16
Unmasked-calculated*	4.09	6.87	4.21

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.09 differs from the reported value 3.11 by more than 10 %

9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-65388 and PDB model 9VVR. Per-residue inclusion information can be found in section 3 on page 8.

9.1 Map-model overlay



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.18 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.18).

9.4 Atom inclusion (i)



At the recommended contour level, 90% of all backbone atoms, 84% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.18) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8420	0.5190
A	0.7240	0.4060
B	0.7920	0.4770
C	0.8190	0.4950
D	0.7340	0.4020
E	0.7870	0.4870
F	0.8260	0.4940
G	0.7290	0.4070
H	0.7240	0.4100
I	0.7110	0.3940
J	0.7940	0.4820
K	0.7920	0.4840
L	0.7840	0.4820
M	0.8230	0.5020
N	0.8190	0.5010
O	0.8260	0.4940
P	0.7270	0.4100
Q	0.7890	0.4780
R	0.8220	0.4990
S	0.8660	0.5380
T	0.8630	0.5410
U	0.8500	0.5370
V	0.8660	0.5380
W	0.8540	0.5380
X	0.8670	0.5370
Y	0.8640	0.5420
Z	0.8680	0.5390
a	0.8500	0.5370
b	0.8670	0.5360
c	0.8450	0.5340
d	0.8690	0.5380
e	0.8520	0.5330
f	0.8610	0.5440
g	0.8540	0.5370
h	0.8620	0.5420

Continued on next page...

Continued from previous page...

Chain	Atom inclusion	Q-score
i	0.8630	0.5410
j	0.8670	0.5400