



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 15, 2024 – 11:40 PM EST

PDB ID : 5W1O
Title : Crystal Structure of HPV16 L1 Pentamer Bound to Heparin Oligosaccharides
Authors : Dasgupta, J.; Chen, X.S.
Deposited on : 2017-06-04
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.21
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3657 (2.80-2.80)
Clashscore	180529	4123 (2.80-2.80)
Ramachandran outliers	177936	4071 (2.80-2.80)
Sidechain outliers	177891	4073 (2.80-2.80)
RSRZ outliers	164620	3659 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	427	<div> <div>4%</div> <div>69%</div> <div>25%</div> <div>5%</div> </div>
1	B	427	<div> <div>4%</div> <div>71%</div> <div>24%</div> <div>• •</div> </div>
1	C	427	<div> <div>4%</div> <div>68%</div> <div>26%</div> <div>5%</div> </div>
1	D	427	<div> <div>5%</div> <div>71%</div> <div>25%</div> <div>•</div> </div>
1	E	427	<div> <div>5%</div> <div>70%</div> <div>26%</div> <div>•</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	2	 50% 50%
2	H	2	 100%
2	I	2	 100%
2	J	2	 100%
2	K	2	 50% 50%
2	L	2	 50% 50%
2	M	2	 50% 50%
2	N	2	 50% 50%
2	O	2	 100%
2	P	2	 100%
3	G	6	 33% 67%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	JHM	F	1	-	-	X	-
2	JHM	H	1	-	-	X	-
2	JHM	I	1	-	-	X	-
2	IDS	I	2	-	-	X	-
2	JHM	J	1	-	-	X	-
2	IDS	J	2	-	-	X	-
2	JHM	M	1	-	-	X	-
2	JHM	P	1	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 17224 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major capsid protein L1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	427	Total	C	N	O	S	0	0	0
			3344	2125	562	637	20			
1	B	427	Total	C	N	O	S	0	0	0
			3344	2125	562	637	20			
1	C	427	Total	C	N	O	S	0	0	0
			3344	2125	562	637	20			
1	D	425	Total	C	N	O	S	0	0	0
			3328	2114	560	634	20			
1	E	425	Total	C	N	O	S	0	0	0
			3327	2115	559	633	20			

There are 35 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	20	ALA	-	expression tag	UNP Q81007
A	432	GLY	-	linker	UNP Q81007
A	433	GLY	-	linker	UNP Q81007
A	434	GLY	-	linker	UNP Q81007
A	435	SER	-	linker	UNP Q81007
A	436	GLY	-	linker	UNP Q81007
A	437	ALA	-	linker	UNP Q81007
B	20	ALA	-	expression tag	UNP Q81007
B	432	GLY	-	linker	UNP Q81007
B	433	GLY	-	linker	UNP Q81007
B	434	GLY	-	linker	UNP Q81007
B	435	SER	-	linker	UNP Q81007
B	436	GLY	-	linker	UNP Q81007
B	437	ALA	-	linker	UNP Q81007
C	20	ALA	-	expression tag	UNP Q81007
C	432	GLY	-	linker	UNP Q81007
C	433	GLY	-	linker	UNP Q81007
C	434	GLY	-	linker	UNP Q81007
C	435	SER	-	linker	UNP Q81007

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	436	GLY	-	linker	UNP Q81007
C	437	ALA	-	linker	UNP Q81007
D	20	ALA	-	expression tag	UNP Q81007
D	432	GLY	-	linker	UNP Q81007
D	433	GLY	-	linker	UNP Q81007
D	434	GLY	-	linker	UNP Q81007
D	435	SER	-	linker	UNP Q81007
D	436	GLY	-	linker	UNP Q81007
D	437	ALA	-	linker	UNP Q81007
E	20	ALA	-	expression tag	UNP Q81007
E	432	GLY	-	linker	UNP Q81007
E	433	GLY	-	linker	UNP Q81007
E	434	GLY	-	linker	UNP Q81007
E	435	SER	-	linker	UNP Q81007
E	436	GLY	-	linker	UNP Q81007
E	437	ALA	-	linker	UNP Q81007

- Molecule 2 is an oligosaccharide called 2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	F	2	Total	C	O	S	0	0	0
			30	12	16	2			
2	H	2	Total	C	O	S	0	0	0
			30	12	16	2			
2	I	2	Total	C	O	S	0	0	0
			30	12	16	2			
2	J	2	Total	C	O	S	0	0	0
			30	12	16	2			
2	K	2	Total	C	O	S	0	0	0
			30	12	16	2			
2	L	2	Total	C	O	S	0	0	0
			30	12	16	2			
2	M	2	Total	C	O	S	0	0	0
			30	12	16	2			
2	N	2	Total	C	O	S	0	0	0
			30	12	16	2			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	O	2	Total	C	O	S	0	0	0
			30	12	16	2			
2	P	2	Total	C	O	S	0	0	0
			30	12	16	2			

- Molecule 3 is an oligosaccharide called 2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	G	6	Total	C	O	S	0	0	0
			90	36	48	6			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	32	Total	O	0	0
			32	32		
4	B	26	Total	O	0	0
			26	26		
4	C	34	Total	O	0	0
			34	34		
4	D	23	Total	O	0	0
			23	23		
4	E	32	Total	O	0	0
			32	32		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Major capsid protein L1




- Molecule 1: Major capsid protein L1




- Molecule 1: Major capsid protein L1



• Molecule 1: Major capsid protein L1



• Molecule 1: Major capsid protein L1





- Molecule 2: 2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-alpha-D-glucopyranose

Chain F: 50%



- Molecule 2: 2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-alpha-D-glucopyranose

Chain H: 100%



- Molecule 2: 2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-alpha-D-glucopyranose

Chain I: 100%



- Molecule 2: 2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-alpha-D-glucopyranose

Chain J: 100%



- Molecule 2: 2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-alpha-D-glucopyranose

Chain K: 50%



- Molecule 2: 2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-alpha-D-glucopyranose

Chain L: 50%




- Molecule 2: 2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-alpha-D-glucopyranose

Chain M:  50% 50%

JHM1
IDS2

- Molecule 2: 2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-alpha-D-glucopyranose

Chain N:  50% 50%

JHM1
IDS2

- Molecule 2: 2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-alpha-D-glucopyranose

Chain O:  100%

JHM1
IDS2

- Molecule 2: 2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-alpha-D-glucopyranose

Chain P:  100%

JHM1
IDS2

- Molecule 3: 2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-alpha-D-glucopyranose

Chain G:  33% 67%

JHM1
IDS2
JHM3
IDS4
JHM5
IDS6

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	96.62Å 101.25Å 128.18Å 90.00° 90.63° 90.00°	Depositor
Resolution (Å)	47.43 – 2.80 47.43 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.1 (47.43-2.80) 93.7 (47.43-2.80)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.65 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, R_{free}	0.224 , 0.270 0.222 , 0.267	Depositor DCC
R_{free} test set	3025 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	44.0	Xtriage
Anisotropy	0.242	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 34.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.011 for -k,-h,-l 0.012 for k,h,-l 0.020 for h,-k,-l	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	17224	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: JHM, IDS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	1/3430 (0.0%)	0.62	1/4664 (0.0%)
1	B	0.48	1/3430 (0.0%)	0.63	2/4664 (0.0%)
1	C	0.45	0/3430	0.62	1/4664 (0.0%)
1	D	0.47	1/3412 (0.0%)	0.63	2/4638 (0.0%)
1	E	0.47	0/3412	0.64	1/4639 (0.0%)
All	All	0.47	3/17114 (0.0%)	0.63	7/23269 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	225	CYS	CB-SG	-6.74	1.70	1.82
1	A	225	CYS	CB-SG	-6.62	1.71	1.82
1	D	225	CYS	CB-SG	-5.21	1.73	1.81

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	68	LEU	CA-CB-CG	-6.95	99.31	115.30
1	E	68	LEU	CA-CB-CG	-6.34	100.73	115.30
1	A	246	LEU	CA-CB-CG	5.54	128.05	115.30
1	B	68	LEU	CA-CB-CG	-5.51	102.62	115.30
1	C	246	LEU	CA-CB-CG	5.46	127.86	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3344	0	3240	90	6
1	B	3344	0	3240	89	1
1	C	3344	0	3240	103	7
1	D	3328	0	3220	71	0
1	E	3327	0	3222	74	0
2	F	30	0	14	22	0
2	H	30	0	14	13	0
2	I	30	0	14	16	0
2	J	30	0	13	12	10
2	K	30	0	13	5	0
2	L	30	0	14	5	0
2	M	30	0	14	6	0
2	N	30	0	14	4	1
2	O	30	0	14	8	0
2	P	30	0	14	12	0
3	G	90	0	39	11	1
4	A	32	0	0	5	0
4	B	26	0	0	4	0
4	C	34	0	0	10	0
4	D	23	0	0	2	0
4	E	32	0	0	6	0
All	All	17224	0	16339	412	13

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 412 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:266:THR:HG21	2:F:1:JHM:C6	1.21	1.63
1:B:266:THR:CG2	2:F:1:JHM:H6A	1.19	1.62
1:A:358:THR:CB	2:F:1:JHM:H1	1.30	1.56
1:A:358:THR:CG2	2:F:1:JHM:C1	1.82	1.55
1:E:452:LYS:NZ	2:P:1:JHM:C6	1.72	1.49

The worst 5 of 13 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:278:LYS:CE	2:J:2:IDS:O6B[2_545]	1.60	0.60
1:C:278:LYS:CE	2:J:2:IDS:O6A[2_545]	1.65	0.55
1:A:353:THR:CG2	2:J:2:IDS:O2S[2_545]	1.81	0.39
1:A:353:THR:OG1	2:J:2:IDS:O2[2_545]	1.81	0.39
1:C:278:LYS:CE	2:J:2:IDS:C6[2_545]	1.86	0.34

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	425/427 (100%)	391 (92%)	22 (5%)	12 (3%)	4	14
1	B	425/427 (100%)	384 (90%)	27 (6%)	14 (3%)	3	11
1	C	425/427 (100%)	390 (92%)	24 (6%)	11 (3%)	4	16
1	D	421/427 (99%)	388 (92%)	23 (6%)	10 (2%)	5	18
1	E	421/427 (99%)	386 (92%)	20 (5%)	15 (4%)	3	10
All	All	2117/2135 (99%)	1939 (92%)	116 (6%)	62 (3%)	3	13

5 of 62 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	137	ALA
1	A	182	PRO
1	A	435	SER
1	A	437	ALA
1	A	460	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	368/368 (100%)	326 (89%)	42 (11%)	4	15
1	B	368/368 (100%)	337 (92%)	31 (8%)	9	28
1	C	368/368 (100%)	330 (90%)	38 (10%)	6	19
1	D	366/368 (100%)	334 (91%)	32 (9%)	8	26
1	E	366/368 (100%)	329 (90%)	37 (10%)	6	20
All	All	1836/1840 (100%)	1656 (90%)	180 (10%)	6	21

5 of 180 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	91	TYR
1	E	43	LEU
1	D	117	ILE
1	D	251	ARG
1	E	96	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	192	ASN
1	E	319	HIS
1	E	366	HIS
1	E	341	ASN
1	C	196	GLN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

26 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	JHM	F	1	2	15,15,15	1.03	0	19,22,22	1.21	2 (10%)
2	IDS	F	2	2	15,15,17	2.01	1 (6%)	14,22,26	1.51	3 (21%)
3	JHM	G	1	3	15,15,15	1.04	1 (6%)	19,22,22	1.20	2 (10%)
3	IDS	G	2	3	16,16,17	1.99	3 (18%)	16,24,26	1.17	1 (6%)
3	JHM	G	3	3	14,14,15	0.96	1 (7%)	19,20,22	1.40	3 (15%)
3	IDS	G	4	3	16,16,17	1.98	3 (18%)	16,24,26	4.11	4 (25%)
3	JHM	G	5	3	14,14,15	0.96	1 (7%)	19,20,22	1.39	3 (15%)
3	IDS	G	6	3	15,15,17	2.02	2 (13%)	14,22,26	1.56	3 (21%)
2	JHM	H	1	2	15,15,15	1.04	1 (6%)	19,22,22	1.22	2 (10%)
2	IDS	H	2	2	15,15,17	2.02	2 (13%)	14,22,26	1.55	3 (21%)
2	JHM	I	1	2	15,15,15	1.05	1 (6%)	19,22,22	1.23	2 (10%)
2	IDS	I	2	2	15,15,17	1.99	1 (6%)	14,22,26	1.51	3 (21%)
2	JHM	J	1	2	15,15,15	1.04	1 (6%)	19,22,22	1.21	2 (10%)
2	IDS	J	2	2	15,15,17	2.00	1 (6%)	14,22,26	1.51	3 (21%)
2	JHM	K	1	2	15,15,15	1.04	1 (6%)	19,22,22	1.21	2 (10%)
2	IDS	K	2	2	15,15,17	1.99	1 (6%)	14,22,26	1.52	3 (21%)
2	JHM	L	1	2	15,15,15	1.04	1 (6%)	19,22,22	1.22	2 (10%)
2	IDS	L	2	2	15,15,17	2.03	2 (13%)	14,22,26	1.54	3 (21%)
2	JHM	M	1	2	15,15,15	1.02	0	19,22,22	1.22	2 (10%)
2	IDS	M	2	2	15,15,17	2.01	2 (13%)	14,22,26	1.54	3 (21%)
2	JHM	N	1	2	15,15,15	1.02	1 (6%)	19,22,22	1.21	2 (10%)
2	IDS	N	2	2	15,15,17	2.00	1 (6%)	14,22,26	1.52	3 (21%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	JHM	O	1	2	15,15,15	1.03	1 (6%)	19,22,22	1.19	2 (10%)
2	IDS	O	2	2	15,15,17	2.00	1 (6%)	14,22,26	1.51	3 (21%)
2	JHM	P	1	2	15,15,15	1.04	1 (6%)	19,22,22	1.21	2 (10%)
2	IDS	P	2	2	15,15,17	1.99	1 (6%)	14,22,26	1.49	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	JHM	F	1	2	-	1/6/22/22	0/1/1/1
2	IDS	F	2	2	-	0/9/22/29	0/1/1/1
3	JHM	G	1	3	-	1/6/22/22	0/1/1/1
3	IDS	G	2	3	-	0/9/26/29	0/1/1/1
3	JHM	G	3	3	-	0/6/20/22	0/1/1/1
3	IDS	G	4	3	-	0/9/26/29	0/1/1/1
3	JHM	G	5	3	-	0/6/20/22	0/1/1/1
3	IDS	G	6	3	-	0/9/22/29	0/1/1/1
2	JHM	H	1	2	-	0/6/22/22	0/1/1/1
2	IDS	H	2	2	-	0/9/22/29	0/1/1/1
2	JHM	I	1	2	-	0/6/22/22	0/1/1/1
2	IDS	I	2	2	-	0/9/22/29	0/1/1/1
2	JHM	J	1	2	-	1/6/22/22	0/1/1/1
2	IDS	J	2	2	-	0/9/22/29	0/1/1/1
2	JHM	K	1	2	-	1/6/22/22	0/1/1/1
2	IDS	K	2	2	-	0/9/22/29	0/1/1/1
2	JHM	L	1	2	-	0/6/22/22	0/1/1/1
2	IDS	L	2	2	-	0/9/22/29	0/1/1/1
2	JHM	M	1	2	-	0/6/22/22	0/1/1/1
2	IDS	M	2	2	-	0/9/22/29	0/1/1/1
2	JHM	N	1	2	-	1/6/22/22	0/1/1/1
2	IDS	N	2	2	-	0/9/22/29	0/1/1/1
2	JHM	O	1	2	-	1/6/22/22	0/1/1/1
2	IDS	O	2	2	-	0/9/22/29	0/1/1/1
2	JHM	P	1	2	-	0/6/22/22	0/1/1/1
2	IDS	P	2	2	-	0/9/22/29	0/1/1/1

The worst 5 of 32 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	2	IDS	O2-C2	-6.78	1.37	1.47
2	N	2	IDS	O2-C2	-6.77	1.37	1.47
2	L	2	IDS	O2-C2	-6.75	1.37	1.47
2	J	2	IDS	O2-C2	-6.74	1.37	1.47
2	O	2	IDS	O2-C2	-6.73	1.37	1.47

The worst 5 of 66 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	4	IDS	O4-C4-C5	15.22	144.52	109.76
3	G	4	IDS	O4-C4-C3	-4.28	100.30	110.38
3	G	3	JHM	C2-C3-C4	-3.51	107.14	111.16
3	G	5	JHM	C2-C3-C4	-3.46	107.20	111.16
2	J	1	JHM	C3-C4-C5	-3.36	106.62	109.99

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	1	JHM	C6-O6-S-O8
2	J	1	JHM	C6-O6-S-O8
2	K	1	JHM	C6-O6-S-O8
2	N	1	JHM	C6-O6-S-O8
2	O	1	JHM	C6-O6-S-O8

There are no ring outliers.

19 monomers are involved in 106 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	H	2	IDS	4	0
3	G	2	IDS	5	0
2	J	1	JHM	10	0
2	I	1	JHM	6	0
3	G	1	JHM	4	0
2	N	1	JHM	4	1
2	P	2	IDS	4	0
3	G	6	IDS	0	1
2	J	2	IDS	2	10
2	O	2	IDS	4	0
2	H	1	JHM	9	0
2	K	1	JHM	5	0
2	F	1	JHM	22	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	P	1	JHM	8	0
2	O	1	JHM	4	0
3	G	3	JHM	2	0
2	M	1	JHM	6	0
2	L	2	IDS	5	0
2	I	2	IDS	10	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

















Oligosaccharide Chain M




Bond lengths



Bond angles




Torsions




Rings


Oligosaccharide Chain N




Bond lengths




Bond angles





Torsions



Rings







5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	427/427 (100%)	-0.05	19 (4%) 39 32	19, 38, 86, 131	0
1	B	427/427 (100%)	-0.04	16 (3%) 45 37	20, 36, 88, 133	0
1	C	427/427 (100%)	0.01	16 (3%) 45 37	20, 39, 95, 144	0
1	D	425/427 (99%)	0.04	21 (4%) 36 28	18, 39, 94, 141	0
1	E	425/427 (99%)	-0.04	21 (4%) 36 28	19, 34, 81, 115	0
All	All	2131/2135 (99%)	-0.01	93 (4%) 39 32	18, 37, 91, 144	0

The worst 5 of 93 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	437	ALA	5.2
1	A	353	THR	5.1
1	E	435	SER	5.1
1	B	403	ASN	4.5
1	E	403	ASN	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	JHM	L	1	15/15	-0.10	0.24	46,66,88,89	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	JHM	O	1	15/15	0.01	0.23	46,66,88,89	0
3	JHM	G	1	15/15	0.04	0.22	46,66,88,89	0
3	JHM	G	5	14/15	0.05	0.25	46,66,88,89	0
2	JHM	J	1	15/15	0.08	0.27	46,66,88,89	0
3	JHM	G	3	14/15	0.11	0.23	46,66,88,89	0
2	JHM	I	1	15/15	0.11	0.21	46,66,88,89	0
2	JHM	K	1	15/15	0.14	0.23	46,66,88,89	0
2	JHM	N	1	15/15	0.14	0.21	46,66,88,89	0
2	JHM	P	1	15/15	0.15	0.22	46,66,88,89	0
2	JHM	M	1	15/15	0.18	0.20	46,66,88,89	0
2	JHM	F	1	15/15	0.19	0.21	46,66,88,89	0
2	JHM	H	1	15/15	0.20	0.23	46,66,88,89	0
2	IDS	K	2	15/17	0.37	0.24	30,31,38,42	0
2	IDS	P	2	15/17	0.40	0.22	30,31,38,42	0
3	IDS	G	2	16/17	0.42	0.24	30,32,42,69	0
3	IDS	G	6	15/17	0.45	0.22	30,31,38,42	0
2	IDS	O	2	15/17	0.46	0.20	30,31,38,42	0
2	IDS	F	2	15/17	0.49	0.19	30,31,38,42	0
2	IDS	J	2	15/17	0.50	0.23	30,31,38,42	0
2	IDS	M	2	15/17	0.51	0.21	30,31,38,42	0
2	IDS	H	2	15/17	0.51	0.24	30,31,38,42	0
2	IDS	N	2	15/17	0.51	0.20	30,31,38,42	0
3	IDS	G	4	16/17	0.52	0.19	30,32,42,69	0
2	IDS	I	2	15/17	0.56	0.19	30,31,38,42	0
2	IDS	L	2	15/17	0.58	0.19	30,31,38,42	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain F:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain H:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain J:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain K:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain L:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain M:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain N:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain O:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain P:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.