



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2024 – 06:40 AM EDT

PDB ID : 5W84
Title : CRYSTAL STRUCTURE OF IRAK-4 WITH A 4,6-DIAMINONICOTINAMIDE INHIBITOR (COMPOUND NUMBER 4)
Authors : Sack, J.S.
Deposited on : 2017-06-21
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	305	 5% 80% 12% • 8%
1	B	305	 10% 76% 11% 13%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4324 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Interleukin-1 receptor-associated kinase 4.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	282	Total	C	N	O	P	S	0	0	0
			2197	1376	366	438	3	14			
1	B	264	Total	C	N	O	P	S	0	0	0
			2044	1277	344	406	3	14			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	156	GLY	-	expression tag	UNP Q9NWZ3
A	157	ALA	-	expression tag	UNP Q9NWZ3
A	158	MET	-	expression tag	UNP Q9NWZ3
A	159	GLY	-	expression tag	UNP Q9NWZ3
B	156	GLY	-	expression tag	UNP Q9NWZ3
B	157	ALA	-	expression tag	UNP Q9NWZ3
B	158	MET	-	expression tag	UNP Q9NWZ3
B	159	GLY	-	expression tag	UNP Q9NWZ3

- Molecule 2 is 4-(benzylamino)-6-[(2-oxo-2,3-dihydro-1H-indol-5-yl)amino]pyridine-3-carboxamide (three-letter code: 9YY) (formula: C₂₁H₁₉N₅O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			28	21	5	2		
2	B	1	Total	C	N	O	0	0
			28	21	5	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	18	Total	O	0	0
			18	18		
3	B	9	Total	O	0	0
			9	9		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Interleukin-1 receptor-associated kinase 4



- Molecule 1: Interleukin-1 receptor-associated kinase 4



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	88.19Å 125.20Å 141.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	93.82 – 2.90 93.82 – 2.90	Depositor EDS
% Data completeness (in resolution range)	95.5 (93.82-2.90) 95.5 (93.82-2.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 2.91Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.200 , 0.248 0.208 , 0.266	Depositor DCC
R_{free} test set	813 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	67.6	Xtriage
Anisotropy	0.625	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 78.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4324	wwPDB-VP
Average B, all atoms (Å ²)	83.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.68 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.0417e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, SEP, 9YY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/2200	0.68	0/2970
1	B	0.50	0/2043	0.67	0/2758
All	All	0.51	0/4243	0.68	0/5728

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2197	0	2122	12	0
1	B	2044	0	1973	12	0
2	A	28	0	0	0	0
2	B	28	0	0	0	0
3	A	18	0	0	0	0
3	B	9	0	0	0	0
All	All	4324	0	4095	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:284:SER:H	1:B:287:MET:HE3	1.51	0.75
1:A:352:THR:O	1:A:353:ALA:HB3	1.89	0.71
1:A:266:PRO:HG3	1:A:321:GLU:HG3	1.77	0.67
1:B:176:VAL:HG22	1:B:204:TYR:H	1.60	0.66
1:B:266:PRO:HG3	1:B:321:GLU:HG3	1.79	0.63
1:A:234:ILE:HD11	1:A:251:PHE:HB3	1.80	0.63
1:A:265:MET:HG3	1:A:318:LEU:HB3	1.82	0.62
1:A:352:THR:O	1:A:353:ALA:CB	2.51	0.58
1:A:237:MET:O	1:A:238:ALA:HB3	2.05	0.56
1:B:212:VAL:HG22	1:B:261:VAL:HG22	1.92	0.52
1:B:452:GLN:HA	1:B:455:GLN:HE21	1.76	0.49
1:A:439:GLU:H	1:A:439:GLU:CD	2.16	0.49
1:A:300:ASN:HA	1:A:447:ILE:HG21	1.94	0.49
1:B:300:ASN:HA	1:B:447:ILE:HG21	1.94	0.49
1:B:176:VAL:HG11	1:B:205:VAL:HG22	1.94	0.48
1:A:356:ALA:HA	1:A:371:TYR:CD2	2.51	0.45
1:A:232:GLN:O	1:A:236:VAL:HG23	2.16	0.45
1:B:240:CYS:HB3	1:B:301:PHE:HE1	1.81	0.45
1:B:388:ASP:HB3	1:B:391:ARG:HB3	1.98	0.45
1:B:356:ALA:HA	1:B:371:TYR:CD2	2.52	0.45
1:A:168:PHE:HE1	1:A:205:VAL:HG11	1.82	0.44
1:B:176:VAL:CG2	1:B:204:TYR:H	2.29	0.42
1:A:237:MET:O	1:A:238:ALA:CB	2.68	0.42
1:B:265:MET:SD	1:B:326:LYS:HG3	2.60	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	272/305 (89%)	248 (91%)	21 (8%)	3 (1%)	14 42

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	254/305 (83%)	240 (94%)	11 (4%)	3 (1%)	13	40
All	All	526/610 (86%)	488 (93%)	32 (6%)	6 (1%)	14	42

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	166	HIS
1	A	228	GLN
1	B	169	SER
1	B	406	GLU
1	A	406	GLU
1	B	441	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/260 (91%)	221 (94%)	15 (6%)	17	45
1	B	216/260 (83%)	210 (97%)	6 (3%)	43	76
All	All	452/520 (87%)	431 (95%)	21 (5%)	27	60

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	167	SER
1	A	169	SER
1	A	186	SER
1	A	233	GLU
1	A	239	LYS
1	A	257	ASP
1	A	277	LEU
1	A	318	LEU
1	A	329	ASP
1	A	344	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	365	THR
1	A	401	GLU
1	A	434	SER
1	A	438	HIS
1	A	439	GLU
1	B	233	GLU
1	B	259	CYS
1	B	277	LEU
1	B	329	ASP
1	B	401	GLU
1	B	434	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	178	ASN
1	B	455	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	SEP	A	346	1	8,9,10	0.81	0	8,12,14	3.26	2 (25%)
1	TPO	A	342	1	8,10,11	1.01	0	10,14,16	1.54	2 (20%)
1	TPO	B	345	1	8,10,11	1.23	2 (25%)	10,14,16	1.22	1 (10%)
1	SEP	B	346	1	8,9,10	0.78	0	8,12,14	2.62	4 (50%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	TPO	A	345	1	8,10,11	1.33	2 (25%)	10,14,16	0.99	0
1	TPO	B	342	1	8,10,11	0.91	0	10,14,16	1.67	3 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	346	1	-	5/5/8/10	-
1	TPO	A	342	1	-	1/9/11/13	-
1	TPO	B	345	1	-	5/9/11/13	-
1	SEP	B	346	1	-	1/5/8/10	-
1	TPO	A	345	1	-	4/9/11/13	-
1	TPO	B	342	1	-	2/9/11/13	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	345	TPO	CB-CA	2.50	1.59	1.53
1	A	345	TPO	CG2-CB	2.25	1.57	1.51
1	B	345	TPO	CB-CA	2.06	1.58	1.53
1	B	345	TPO	P-OG1	-2.00	1.55	1.59

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	346	SEP	OG-CB-CA	8.59	116.51	108.14
1	B	346	SEP	OG-CB-CA	6.00	113.98	108.14
1	B	342	TPO	P-OG1-CB	-3.22	113.47	123.21
1	A	342	TPO	P-OG1-CB	-3.18	113.61	123.21
1	B	342	TPO	O2P-P-OG1	2.90	118.99	105.99
1	A	346	SEP	O3P-P-OG	2.78	114.13	106.73
1	A	342	TPO	O2P-P-OG1	2.54	117.36	105.99
1	B	346	SEP	OG-P-O1P	2.48	113.42	106.47
1	B	345	TPO	O2P-P-OG1	2.45	116.98	105.99
1	B	346	SEP	P-OG-CB	-2.24	112.13	118.30
1	B	346	SEP	O2P-P-O1P	-2.23	101.95	110.68
1	B	342	TPO	CG2-CB-CA	2.19	117.48	113.16

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	345	TPO	N-CA-CB-OG1
1	A	346	SEP	CA-CB-OG-P
1	A	346	SEP	CB-OG-P-O1P
1	B	342	TPO	CB-OG1-P-O1P
1	B	345	TPO	N-CA-CB-OG1
1	B	346	SEP	N-CA-CB-OG
1	A	342	TPO	CB-OG1-P-O1P
1	A	346	SEP	CB-OG-P-O2P
1	A	346	SEP	N-CA-CB-OG
1	A	345	TPO	CB-OG1-P-O1P
1	A	346	SEP	CB-OG-P-O3P
1	B	345	TPO	CB-OG1-P-O1P
1	A	345	TPO	CB-OG1-P-O3P
1	B	342	TPO	CB-OG1-P-O3P
1	B	345	TPO	CB-OG1-P-O2P
1	B	345	TPO	CB-OG1-P-O3P
1	A	345	TPO	O-C-CA-CB
1	B	345	TPO	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	9YY	B	4000	-	31,31,31	0.45	0	41,43,43	0.48	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	9YY	A	4000	-	31,31,31	0.43	0	41,43,43	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	9YY	B	4000	-	-	0/13/21/21	0/4/4/4
2	9YY	A	4000	-	-	0/13/21/21	0/4/4/4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	279/305 (91%)	0.69	16 (5%) 23 19	42, 77, 112, 162	0
1	B	261/305 (85%)	0.83	29 (11%) 5 4	46, 77, 145, 167	0
All	All	540/610 (88%)	0.76	45 (8%) 11 8	42, 77, 134, 167	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	196	GLY	4.6
1	B	185	ILE	4.5
1	B	343	VAL	4.1
1	B	173	LEU	3.7
1	A	195	GLY	3.6
1	B	207	ASN	3.6
1	A	187	VAL	3.5
1	B	186	SER	3.4
1	B	192	MET	3.4
1	A	168	PHE	3.3
1	B	205	VAL	3.2
1	B	336	SER	3.1
1	A	251	PHE	3.1
1	A	236	VAL	3.0
1	B	250	GLY	2.9
1	B	194	GLU	2.9
1	A	399	ILE	2.9
1	A	258	LEU	2.9
1	B	171	TYR	2.8
1	B	188	GLY	2.8
1	B	206	ASN	2.7
1	B	197	PHE	2.7
1	B	251	PHE	2.6
1	A	230	PHE	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	190	ASN	2.6
1	B	240	CYS	2.5
1	B	212	VAL	2.5
1	B	201	TYR	2.4
1	A	343	VAL	2.4
1	B	198	GLY	2.4
1	B	263	VAL	2.3
1	A	336	SER	2.3
1	A	199	VAL	2.2
1	B	204	TYR	2.2
1	B	334	ARG	2.2
1	B	264	TYR	2.2
1	A	165	PHE	2.1
1	B	202	LYS	2.1
1	B	179	ASN	2.1
1	A	433	ALA	2.1
1	A	413	TYR	2.1
1	B	330	PHE	2.0
1	B	191	LYS	2.0
1	B	395	LEU	2.0
1	A	253	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TPO	B	342	11/12	0.76	0.24	118,120,125,125	0
1	TPO	A	342	11/12	0.82	0.24	110,113,122,122	0
1	SEP	A	346	10/11	0.89	0.14	111,116,123,124	0
1	SEP	B	346	10/11	0.91	0.11	116,119,125,125	0
1	TPO	B	345	11/12	0.92	0.16	111,112,118,118	0
1	TPO	A	345	11/12	0.96	0.14	104,108,110,111	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	9YY	B	4000	28/28	0.82	0.44	101,106,118,118	0
2	9YY	A	4000	28/28	0.90	0.39	84,90,96,96	0

6.5 Other polymers [i](#)

There are no such residues in this entry.