



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 08:38 PM EDT

PDB ID : 3WSF
Title : Oxidized HcgD from Methanocaldococcus jannaschii with citrate
Authors : Fujishiro, T.; Ermler, U.; Shima, S.
Deposited on : 2014-03-13
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	252	 92% 7%
1	B	252	 94%
1	C	252	 89% 8%
1	D	252	 94%
1	E	252	 96%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	252	<div><div></div><div>89%</div><div>8% •</div></div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13050 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative GTP cyclohydrolase 1 type 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	248	Total	C	N	O	S	0	0	0
			1964	1274	312	375	3			
1	B	244	Total	C	N	O	S	0	0	0
			1933	1255	307	368	3			
1	C	245	Total	C	N	O	S	0	0	0
			1941	1259	309	370	3			
1	D	247	Total	C	N	O	S	0	0	0
			1955	1269	311	372	3			
1	E	246	Total	C	N	O	S	0	0	0
			1948	1264	310	371	3			
1	F	246	Total	C	N	O	S	0	2	0
			1961	1273	312	373	3			

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Fe	0	0
			1	1		
2	B	1	Total	Fe	0	0
			1	1		
2	C	1	Total	Fe	0	0
			1	1		
2	D	1	Total	Fe	0	0
			1	1		
2	E	1	Total	Fe	0	0
			1	1		
2	F	1	Total	Fe	0	0
			1	1		

- Molecule 3 is CITRIC ACID (three-letter code: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		
3	B	1	Total	C	O	0	0
			13	6	7		
3	C	1	Total	C	O	0	0
			13	6	7		
3	D	1	Total	C	O	0	0
			13	6	7		
3	E	1	Total	C	O	0	0
			13	6	7		
3	F	1	Total	C	O	0	0
			13	6	7		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	231	Total	O	0	0
			231	231		
4	B	212	Total	O	0	0
			212	212		
4	C	224	Total	O	0	0
			224	224		
4	D	190	Total	O	0	0
			190	190		
4	E	186	Total	O	0	0
			186	186		
4	F	221	Total	O	0	0
			221	221		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative GTP cyclohydrolase 1 type 2

Chain A: 





- Molecule 1: Putative GTP cyclohydrolase 1 type 2

Chain B: 



- Molecule 1: Putative GTP cyclohydrolase 1 type 2

Chain C: 



- Molecule 1: Putative GTP cyclohydrolase 1 type 2

Chain D: 



- Molecule 1: Putative GTP cyclohydrolase 1 type 2

Chain E: 




- Molecule 1: Putative GTP cyclohydrolase 1 type 2

Chain F:

89%

8%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	95.19Å 137.25Å 160.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.78 – 2.00 45.78 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.5 (45.78-2.00) 99.5 (45.78-2.00)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, R_{free}	0.178 , 0.214 0.182 , 0.216	Depositor DCC
R_{free} test set	7116 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	24.4	Xtriage
Anisotropy	0.221	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13050	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CIT, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/1999	0.55	0/2694
1	B	0.42	0/1968	0.55	0/2651
1	C	0.42	0/1976	0.55	1/2662 (0.0%)
1	D	0.38	0/1990	0.55	0/2682
1	E	0.38	0/1983	0.55	0/2672
1	F	0.42	0/2003	0.56	0/2700
All	All	0.41	0/11919	0.55	1/16061 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	45	LEU	CA-CB-CG	5.04	126.90	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1964	0	2008	13	0
1	B	1933	0	1978	4	0
1	C	1941	0	1984	13	0
1	D	1955	0	2002	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1948	0	1993	3	0
1	F	1961	0	2006	14	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	13	0	5	2	0
3	B	13	0	5	1	0
3	C	13	0	5	1	0
3	D	13	0	5	1	0
3	E	13	0	5	1	0
3	F	13	0	5	1	0
4	A	231	0	0	1	0
4	B	212	0	0	0	0
4	C	224	0	0	1	0
4	D	190	0	0	1	0
4	E	186	0	0	0	0
4	F	221	0	0	1	0
All	All	13050	0	12001	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:179:TYR:HB3	3:F:302:CIT:H41	1.63	0.79
1:D:179:TYR:HB3	3:D:302:CIT:H41	1.68	0.75
1:C:77:PRO:HB2	1:D:77:PRO:HB2	1.78	0.66
1:F:4:VAL:HG23	1:F:5:ASN:H	1.61	0.65
1:A:77:PRO:HB2	1:B:77:PRO:HB2	1.76	0.65
1:B:179:TYR:HB3	3:B:302:CIT:H41	1.80	0.63
1:A:179:TYR:HB3	3:A:302:CIT:H41	1.83	0.60
1:E:77:PRO:HB2	1:F:77:PRO:HB2	1.89	0.54
1:A:187:LYS:CG	1:A:208:LEU:HD21	2.39	0.53
1:C:204:HIS:NE2	3:E:302:CIT:O1	2.38	0.53
1:F:49:LEU:HD11	1:F:225:GLU:OE1	2.09	0.52
1:D:183:GLN:NE2	4:D:572:HOH:O	2.42	0.52
1:C:45:LEU:HD12	1:C:244:ILE:HG23	1.92	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:49:LEU:HD11	1:C:225:GLU:OE2	2.12	0.50
1:D:49:LEU:HD11	1:D:225:GLU:OE1	2.12	0.50
1:C:179:TYR:HB3	3:C:302:CIT:H41	1.95	0.48
3:A:302:CIT:O3	1:F:204:HIS:NE2	2.46	0.47
1:A:2:GLU:O	1:A:241:ASP:HB3	2.15	0.47
1:B:123:LEU:HD21	1:B:173:LEU:HD23	1.97	0.47
1:C:6:MET:HA	1:C:10:GLU:OE2	2.15	0.47
1:F:16:GLU:OE2	1:F:21:LYS:HD2	2.15	0.46
1:A:182:SER:O	1:A:186:ILE:HG12	2.16	0.46
1:F:123:LEU:HD21	1:F:173:LEU:HD23	1.98	0.45
1:F:36:ASP:HB3	4:F:573:HOH:O	2.16	0.45
1:B:49:LEU:HD11	1:B:225:GLU:OE1	2.16	0.45
1:F:40:LYS:HE3	1:F:40:LYS:HB2	1.73	0.44
1:D:123:LEU:HD21	1:D:173:LEU:HD23	1.99	0.44
1:F:140:PHE:O	1:F:170:ASN:HA	2.18	0.44
1:A:187:LYS:HE2	1:A:208:LEU:HD21	2.00	0.44
1:A:187:LYS:HG2	1:A:208:LEU:HD21	2.01	0.43
1:C:176:LEU:HD22	1:C:180:GLY:HA3	2.02	0.42
1:D:50:ASP:OD1	1:D:71:HIS:ND1	2.36	0.42
1:F:42:ILE:HG23	1:F:65:ASP:OD1	2.19	0.42
1:A:123:LEU:HD21	1:A:173:LEU:HD23	2.02	0.42
1:C:7:LYS:O	1:C:10:GLU:HG2	2.20	0.42
1:F:6:MET:HB2	1:F:10:GLU:HG3	2.01	0.42
1:A:49:LEU:HD11	1:A:225:GLU:OE1	2.20	0.42
1:A:140:PHE:O	1:A:170:ASN:HA	2.20	0.42
1:C:94:GLU:HB2	4:C:545:HOH:O	2.19	0.42
1:C:152:LYS:HA	1:C:156:HIS:O	2.20	0.42
1:F:21:LYS:HG2	1:F:30:ILE:HD13	2.01	0.41
1:A:139:ILE:HD13	4:A:574:HOH:O	2.20	0.41
1:C:140:PHE:O	1:C:170:ASN:HA	2.20	0.41
1:C:231:LYS:O	1:C:234:GLU:HG2	2.20	0.41
1:C:42:ILE:HG23	1:C:65:ASP:OD1	2.21	0.41
1:E:123:LEU:HD21	1:E:173:LEU:HD23	2.02	0.41
1:E:86:TYR:CE2	1:F:93:MET:HG2	2.56	0.41
1:A:47:ILE:HD13	1:A:68:PHE:HB3	2.03	0.40
1:A:187:LYS:HG3	1:A:208:LEU:HD21	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	246/252 (98%)	241 (98%)	5 (2%)	0	100	100
1	B	242/252 (96%)	237 (98%)	5 (2%)	0	100	100
1	C	243/252 (96%)	238 (98%)	5 (2%)	0	100	100
1	D	245/252 (97%)	240 (98%)	5 (2%)	0	100	100
1	E	244/252 (97%)	239 (98%)	5 (2%)	0	100	100
1	F	246/252 (98%)	241 (98%)	5 (2%)	0	100	100
All	All	1466/1512 (97%)	1436 (98%)	30 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/222 (99%)	219 (100%)	0	100	100
1	B	215/222 (97%)	215 (100%)	0	100	100
1	C	216/222 (97%)	216 (100%)	0	100	100
1	D	218/222 (98%)	217 (100%)	1 (0%)	88	92
1	E	217/222 (98%)	216 (100%)	1 (0%)	88	92
1	F	219/222 (99%)	219 (100%)	0	100	100
All	All	1304/1332 (98%)	1302 (100%)	2 (0%)	93	95

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	75	LYS
1	E	26	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	CIT	A	302	2	12,12,12	1.09	0	17,17,17	1.38	4 (23%)
3	CIT	F	302	2	12,12,12	1.18	0	17,17,17	1.77	5 (29%)
3	CIT	D	302	2	12,12,12	1.09	0	17,17,17	1.59	4 (23%)
3	CIT	E	302	2	12,12,12	1.05	0	17,17,17	1.45	4 (23%)
3	CIT	C	302	2	12,12,12	1.13	0	17,17,17	1.30	3 (17%)
3	CIT	B	302	2	12,12,12	0.98	0	17,17,17	1.60	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CIT	A	302	2	-	0/16/16/16	-
3	CIT	F	302	2	-	5/16/16/16	-
3	CIT	D	302	2	-	5/16/16/16	-
3	CIT	E	302	2	-	4/16/16/16	-
3	CIT	C	302	2	-	0/16/16/16	-
3	CIT	B	302	2	-	9/16/16/16	-

There are no bond length outliers.

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	302	CIT	O6-C6-C3	4.12	121.05	113.14
3	B	302	CIT	O6-C6-C3	3.63	120.11	113.14
3	F	302	CIT	O2-C1-C2	3.50	125.44	114.35
3	A	302	CIT	O6-C6-C3	2.88	118.67	113.14
3	F	302	CIT	O2-C1-O1	-2.81	116.11	123.33
3	B	302	CIT	O2-C1-C2	2.78	123.15	114.35
3	E	302	CIT	O6-C6-C3	2.76	118.43	113.14
3	F	302	CIT	C3-C2-C1	2.51	120.77	113.92
3	C	302	CIT	O6-C6-C3	2.50	117.94	113.14
3	D	302	CIT	O2-C1-O1	-2.44	117.06	123.33
3	F	302	CIT	O6-C6-C3	2.41	117.75	113.14
3	E	302	CIT	O4-C5-C4	2.39	121.91	114.35
3	A	302	CIT	O4-C5-C4	2.31	121.66	114.35
3	B	302	CIT	O2-C1-O1	-2.26	117.53	123.33
3	E	302	CIT	O2-C1-O1	-2.25	117.56	123.33
3	E	302	CIT	O2-C1-C2	2.15	121.17	114.35
3	C	302	CIT	O2-C1-O1	-2.15	117.80	123.33
3	A	302	CIT	O2-C1-O1	-2.15	117.80	123.33
3	A	302	CIT	C3-C4-C5	-2.10	108.19	113.92
3	B	302	CIT	O4-C5-C4	2.04	120.80	114.35
3	F	302	CIT	O4-C5-O3	-2.03	118.11	123.33
3	C	302	CIT	C3-C4-C5	-2.02	108.41	113.92
3	D	302	CIT	O4-C5-O3	-2.01	118.16	123.33
3	D	302	CIT	O5-C6-C3	-2.01	118.21	122.09

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	302	CIT	C2-C3-C6-O5
3	D	302	CIT	O7-C3-C6-O5
3	D	302	CIT	O7-C3-C6-O6
3	F	302	CIT	C1-C2-C3-O7
3	F	302	CIT	C1-C2-C3-C6
3	D	302	CIT	C2-C3-C6-O6
3	F	302	CIT	C1-C2-C3-C4
3	B	302	CIT	O1-C1-C2-C3
3	B	302	CIT	O2-C1-C2-C3
3	B	302	CIT	C2-C3-C6-O5
3	D	302	CIT	C4-C3-C6-O6
3	F	302	CIT	O2-C1-C2-C3
3	F	302	CIT	O1-C1-C2-C3
3	B	302	CIT	C2-C3-C6-O6
3	B	302	CIT	C4-C3-C6-O5
3	B	302	CIT	C4-C3-C6-O6
3	E	302	CIT	C2-C3-C6-O5
3	E	302	CIT	C2-C3-C6-O6
3	E	302	CIT	C4-C3-C6-O5
3	E	302	CIT	C4-C3-C6-O6
3	B	302	CIT	C3-C4-C5-O4
3	B	302	CIT	C3-C4-C5-O3
3	B	302	CIT	O7-C3-C6-O5

There are no ring outliers.

6 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	302	CIT	2	0
3	F	302	CIT	1	0
3	D	302	CIT	1	0
3	E	302	CIT	1	0
3	C	302	CIT	1	0
3	B	302	CIT	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	248/252 (98%)	-0.29	1 (0%) 92 92	14, 25, 43, 65	0
1	B	244/252 (96%)	-0.33	1 (0%) 92 92	15, 25, 42, 53	0
1	C	245/252 (97%)	-0.28	2 (0%) 86 85	17, 26, 43, 69	0
1	D	247/252 (98%)	-0.22	3 (1%) 79 78	18, 28, 46, 60	0
1	E	246/252 (97%)	-0.16	4 (1%) 72 70	18, 30, 50, 70	0
1	F	246/252 (97%)	-0.32	1 (0%) 92 92	14, 23, 41, 64	0
All	All	1476/1512 (97%)	-0.27	12 (0%) 86 85	14, 26, 46, 70	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	4	VAL	3.6
1	E	4	VAL	3.5
1	C	5	ASN	3.1
1	D	4	VAL	3.0
1	E	5	ASN	2.8
1	A	2	GLU	2.6
1	D	3	VAL	2.4
1	C	241	ASP	2.2
1	B	22	ASP	2.1
1	E	241	ASP	2.1
1	D	241	ASP	2.1
1	E	131	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	CIT	D	302	13/13	0.77	0.24	28,44,54,55	13
3	CIT	B	302	13/13	0.78	0.26	27,39,49,52	13
3	CIT	E	302	13/13	0.79	0.23	28,43,50,54	13
3	CIT	A	302	13/13	0.80	0.22	21,38,47,53	13
3	CIT	F	302	13/13	0.80	0.24	17,41,45,47	13
3	CIT	C	302	13/13	0.83	0.22	25,42,53,53	13
2	FE	C	301	1/1	0.99	0.07	25,25,25,25	1
2	FE	D	301	1/1	0.99	0.06	26,26,26,26	1
2	FE	E	301	1/1	0.99	0.06	26,26,26,26	1
2	FE	F	301	1/1	0.99	0.06	26,26,26,26	1
2	FE	A	301	1/1	0.99	0.06	25,25,25,25	1
2	FE	B	301	1/1	1.00	0.05	22,22,22,22	1

6.5 Other polymers [i](#)

There are no such residues in this entry.