



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2024 – 09:16 PM EDT

PDB ID : 6XEC  
Title : STRUCTURE OF HUMAN HDAC2 IN COMPLEX WITH KETONE INHIBITOR (COMPOUND O)  
Authors : Klein, D.J.; Clausen, D.  
Deposited on : 2020-06-12  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	376	<div> <div>4%</div> <div>95%</div> </div>
1	B	376	<div> <div>5%</div> <div>93%</div> </div>
1	C	376	<div> <div>8%</div> <div>94%</div> </div>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 10267 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone deacetylase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	368	Total	C	N	O	S	0	15	0
			3012	1931	507	548	26			
1	B	366	Total	C	N	O	S	0	18	0
			3021	1937	503	553	28			
1	C	366	Total	C	N	O	S	0	6	0
			2975	1900	503	546	26			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Ca	0	0
			2	2		
4	B	2	Total	Ca	0	0
			2	2		

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	2	Total	Ca	0	0
			2	2		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		
5	C	1	Total	C	O	0	0
			7	4	3		
5	C	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is (1S)-N-{(1S)-1-[5-cyano-2-(4-fluorophenyl)-1H-imidazol-4-yl]-7,7-dihydroxynonyl}-6-methyl-6-azaspiro[2.5]octane-1-carboxamide (three-letter code: V1S) (formula:  $C_{28}H_{38}FN_5O_3$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	F	N	O	0	0
			37	28	1	5	3		
6	B	1	Total	C	F	N	O	0	0
			37	28	1	5	3		
6	C	1	Total	C	F	N	O	0	0
			37	28	1	5	3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	342	Total	O	0	0
			342	342		
7	B	400	Total	O	0	4
			404	404		
7	C	291	Total	O	0	3
			294	294		



- Molecule 1: Histone deacetylase 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.60Å 99.15Å 139.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.79 – 1.70 24.79 – 1.70	Depositor EDS
% Data completeness (in resolution range)	96.6 (24.79-1.70) 96.6 (24.79-1.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.57 (at 1.71Å)	Xtriage
Refinement program	BUSTER 2.11.7 (19-MAR-2020)	Depositor
R, $R_{free}$	0.175 , 0.203 0.182 , 0.214	Depositor DCC
$R_{free}$ test set	6798 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.3	Xtriage
Anisotropy	0.382	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 47.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	10267	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SO4, CA, V1S, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.43	0/3112	0.59	1/4201 (0.0%)
1	B	0.45	0/3136	0.60	0/4233
1	C	0.37	0/3059	0.55	1/4130 (0.0%)
All	All	0.42	0/9307	0.58	2/12564 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	305	THR	N-CA-C	-5.13	97.14	111.00
1	C	305	THR	N-CA-C	-5.05	97.38	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3012	0	2920	6	0
1	B	3021	0	2942	9	0
1	C	2975	0	2879	7	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	20	0	0	0	0
3	B	10	0	0	0	0
3	C	20	0	0	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
4	C	2	0	0	0	0
5	A	21	0	30	0	0
5	B	14	0	20	1	0
5	C	14	0	20	1	0
6	A	37	0	0	0	0
6	B	37	0	0	0	0
6	C	37	0	0	0	0
7	A	342	0	0	0	0
7	B	404	0	0	0	0
7	C	294	0	0	0	0
All	All	10267	0	8811	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:328:GLU:HG3	1:C:343:LYS:NZ	2.13	0.64
1:B:43[A]:LEU:HD12	1:B:49:TYR:HE1	1.64	0.63
1:B:241:ILE:HG22	1:B:364:LYS:HE2	1.83	0.59
1:B:43[A]:LEU:HD12	1:B:49:TYR:CE1	2.40	0.56
1:B:44:LEU:HB2	1:B:49:TYR:CZ	2.44	0.53
1:C:139:GLY:HA2	1:C:157:ILE:HD11	1.93	0.50
1:A:241:ILE:HG12	1:A:360:MET:CE	2.41	0.50
1:C:328:GLU:HG3	1:C:343:LYS:HZ3	1.76	0.49
1:A:241:ILE:HG12	1:A:360:MET:HE3	1.95	0.49
1:A:10:LYS:HB3	1:A:131:ASP:HB2	1.94	0.49
1:B:139:GLY:HA2	1:B:157:ILE:HD11	1.94	0.48
1:C:328:GLU:HG3	1:C:343:LYS:HZ2	1.79	0.47
1:C:51:LYS:NZ	1:C:323[A]:CYS:SG	2.89	0.45
1:A:139:GLY:HA2	1:A:157:ILE:HD11	1.99	0.44
1:B:72:GLU:HG2	1:B:95:PHE:HZ	1.82	0.44
1:A:267:LEU:HD11	1:A:312:CYS:HB2	1.99	0.44
5:B:406:PEG:O1	5:C:408:PEG:O4	2.28	0.43
1:B:49:TYR:HA	1:B:52[B]:MET:HG2	2.01	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:267:LEU:HD11	1:B:312:CYS:HB2	2.01	0.42
1:A:76:PHE:HE2	1:A:97:VAL:HG21	1.85	0.42
1:C:101:CYS:HB3	1:C:153:TYR:CE2	2.55	0.42
1:C:52:MET:HG2	1:C:321:LEU:HD21	2.02	0.41
1:B:48:LEU:O	1:B:52[B]:MET:SD	2.79	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	375/376 (100%)	371 (99%)	4 (1%)	0	100	100
1	B	379/376 (101%)	375 (99%)	4 (1%)	0	100	100
1	C	368/376 (98%)	362 (98%)	6 (2%)	0	100	100
All	All	1122/1128 (100%)	1108 (99%)	14 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	324/322 (101%)	323 (100%)	1 (0%)	92	89
1	B	330/322 (102%)	328 (99%)	2 (1%)	86	80

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	320/322 (99%)	317 (99%)	3 (1%)	78	70
All	All	974/966 (101%)	968 (99%)	6 (1%)	86	80

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	141	HIS
1	B	83	ASP
1	B	141	HIS
1	C	141	HIS
1	C	151	PHE
1	C	337	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 29 ligands modelled in this entry, 9 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	C	404	-	4,4,4	0.25	0	6,6,6	0.10	0
3	SO4	B	402	-	4,4,4	0.25	0	6,6,6	0.15	0
3	SO4	B	403	-	4,4,4	0.28	0	6,6,6	0.09	0
3	SO4	A	403	-	4,4,4	0.43	0	6,6,6	0.45	0
3	SO4	C	405	-	4,4,4	0.26	0	6,6,6	0.08	0
5	PEG	A	408	-	6,6,6	0.09	0	5,5,5	0.05	0
5	PEG	C	408	-	6,6,6	0.07	0	5,5,5	0.05	0
3	SO4	C	402	-	4,4,4	0.24	0	6,6,6	0.07	0
5	PEG	A	410	-	6,6,6	0.10	0	5,5,5	0.02	0
5	PEG	B	407	-	6,6,6	0.13	0	5,5,5	0.08	0
3	SO4	A	404	-	4,4,4	0.25	0	6,6,6	0.07	0
3	SO4	A	405	-	4,4,4	0.25	0	6,6,6	0.09	0
3	SO4	C	403	-	4,4,4	0.35	0	6,6,6	0.13	0
6	V1S	C	410	2	35,40,40	0.73	2 (5%)	35,58,58	0.55	0
6	V1S	A	411	2	35,40,40	0.75	1 (2%)	35,58,58	0.56	0
3	SO4	A	402	-	4,4,4	0.26	0	6,6,6	0.14	0
5	PEG	C	409	-	6,6,6	0.14	0	5,5,5	0.07	0
5	PEG	A	409	-	6,6,6	0.10	0	5,5,5	0.08	0
5	PEG	B	406	-	6,6,6	0.09	0	5,5,5	0.11	0
6	V1S	B	408	2	35,40,40	0.80	2 (5%)	35,58,58	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	V1S	A	411	2	-	0/24/50/50	0/4/4/4
6	V1S	C	410	2	-	0/24/50/50	0/4/4/4
5	PEG	A	410	-	-	2/4/4/4	-
5	PEG	B	407	-	-	2/4/4/4	-
5	PEG	C	409	-	-	0/4/4/4	-
5	PEG	B	406	-	-	0/4/4/4	-
5	PEG	A	409	-	-	2/4/4/4	-
5	PEG	C	408	-	-	0/4/4/4	-
6	V1S	B	408	2	-	0/24/50/50	0/4/4/4
5	PEG	A	408	-	-	1/4/4/4	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	408	V1S	C18-C26	-2.35	1.37	1.42
6	C	410	V1S	C19-N2	2.18	1.38	1.35
6	A	411	V1S	C19-N3	-2.04	1.33	1.35
6	C	410	V1S	C19-N3	-2.03	1.33	1.35
6	B	408	V1S	C19-N3	-2.02	1.33	1.35

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	410	PEG	C4-C3-O2-C2
5	A	408	PEG	C4-C3-O2-C2
5	A	409	PEG	C4-C3-O2-C2
5	A	409	PEG	C1-C2-O2-C3
5	B	407	PEG	C4-C3-O2-C2
5	B	407	PEG	C1-C2-O2-C3
5	A	410	PEG	C1-C2-O2-C3


There are no ring outliers.

2 monomers are involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	408	PEG	1	0
5	B	406	PEG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


## Ligand V1S C 410



## Ligand V1S A 411







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	368/376 (97%)	0.21	15 (4%)	37 41	18, 25, 40, 48	5 (1%)
1	B	366/376 (97%)	0.16	18 (4%)	29 33	16, 23, 39, 50	3 (0%)
1	C	366/376 (97%)	0.52	31 (8%)	10 12	21, 32, 48, 63	2 (0%)
All	All	1100/1128 (97%)	0.30	64 (5%)	23 25	16, 26, 44, 63	10 (0%)

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	49	TYR	11.5
1	C	337	TYR	5.7
1	C	350	ASN	5.2
1	B	10	LYS	5.1
1	A	49[A]	TYR	4.6
1	C	340	PRO	4.4
1	C	50	ARG	4.3
1	C	44	LEU	4.2
1	C	346	ILE	3.7
1	C	204	GLU	3.6
1	A	350	ASN	3.6
1	C	375	PRO	3.6
1	C	324	GLU	3.5
1	A	51	LYS	3.5
1	C	28	GLY	3.5
1	C	27	GLN	3.2
1	C	128	GLN	3.2
1	C	341	ASP	3.2
1	B	49	TYR	3.2
1	C	58	HIS	3.2
1	A	27	GLN	3.2
1	C	347	SER	3.2
1	A	167	TYR	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	348	PRO	3.1
1	A	236	GLU	3.0
1	C	322	ASP	2.9
1	A	375	PRO	2.8
1	C	90	LYS	2.7
1	A	363	ILE	2.7
1	C	87	GLU	2.7
1	C	342	PHE	2.7
1	B	51	LYS	2.7
1	B	176	ILE	2.7
1	C	240	GLN	2.7
1	B	197	VAL	2.7
1	C	88	TYR	2.7
1	A	50[A]	ARG	2.6
1	B	86	SER	2.6
1	C	99	GLU	2.5
1	B	27	GLN	2.5
1	A	72	GLU	2.4
1	A	227	PHE	2.4
1	B	11	LYS	2.4
1	B	174	ILE	2.4
1	A	324	GLU	2.3
1	C	86	SER	2.3
1	C	83	ASP	2.3
1	B	97	VAL	2.3
1	C	292	PHE	2.3
1	C	24	TYR	2.3
1	C	219	LYS	2.2
1	C	368	PHE	2.2
1	B	195	MET	2.2
1	B	173	TYR	2.2
1	B	225	VAL	2.2
1	B	83	ASP	2.1
1	A	174	ILE	2.1
1	C	55	TYR	2.1
1	B	259	VAL	2.1
1	A	254	GLN	2.1
1	A	86	SER	2.1
1	B	172	LEU	2.0
1	B	87	GLU	2.0
1	B	375	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SO4	C	404	5/5	0.53	0.31	121,121,122,122	0
5	PEG	B	407	7/7	0.58	0.25	67,67,68,68	0
5	PEG	A	410	7/7	0.64	0.18	69,69,69,69	0
3	SO4	A	404	5/5	0.64	0.36	120,120,121,121	0
5	PEG	A	409	7/7	0.71	0.25	62,62,62,62	0
5	PEG	C	408	7/7	0.74	0.19	61,61,61,61	0
5	PEG	A	408	7/7	0.75	0.27	65,65,65,65	0
5	PEG	C	409	7/7	0.75	0.38	69,70,70,70	0
3	SO4	A	405	5/5	0.76	0.24	117,117,117,117	0
3	SO4	A	403	5/5	0.81	0.31	40,41,42,43	0
6	V1S	C	410	37/37	0.82	0.16	26,34,41,41	0
5	PEG	B	406	7/7	0.83	0.19	78,78,78,78	0
3	SO4	C	403	5/5	0.91	0.24	58,58,59,59	0
6	V1S	B	408	37/37	0.91	0.12	19,25,35,36	0
3	SO4	B	403	5/5	0.91	0.29	91,91,91,91	0
6	V1S	A	411	37/37	0.92	0.11	20,25,33,33	0
3	SO4	C	402	5/5	0.93	0.13	115,115,115,115	0
3	SO4	C	405	5/5	0.93	0.30	89,89,89,89	0
3	SO4	B	402	5/5	0.93	0.18	68,68,68,68	0
3	SO4	A	402	5/5	0.94	0.14	74,74,74,74	0
4	CA	C	406	1/1	0.96	0.23	50,50,50,50	0
4	CA	A	407	1/1	0.98	0.09	50,50,50,50	0
2	ZN	C	401	1/1	0.99	0.04	25,25,25,25	0
4	CA	C	407	1/1	0.99	0.11	46,46,46,46	0
4	CA	B	404	1/1	0.99	0.10	40,40,40,40	0
4	CA	B	405	1/1	0.99	0.13	47,47,47,47	0
2	ZN	B	401	1/1	1.00	0.09	18,18,18,18	0

*Continued on next page...*




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZN	A	401	1/1	1.00	0.07	19,19,19,19	0
4	CA	A	406	1/1	1.00	0.16	36,36,36,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around V1S C 410:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around V1S B 408:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around V1S A 411:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers ⓘ

There are no such residues in this entry.