



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 07:37 PM EDT

PDB ID : 2Y98
Title : Structure of the mixed-function P450 MycG in complex with mycinamicin IV
in P21212 space group
Authors : Li, S.; Kells, P.M.; Sherman, D.H.; Podust, L.M.
Deposited on : 2011-02-12
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.36.2
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	417	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3715 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called P-450-LIKE PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	393	3203	2008	582	601	12	0	15	0

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP Q59523
A	-18	GLY	-	expression tag	UNP Q59523
A	-17	SER	-	expression tag	UNP Q59523
A	-16	SER	-	expression tag	UNP Q59523
A	-15	HIS	-	expression tag	UNP Q59523
A	-14	HIS	-	expression tag	UNP Q59523
A	-13	HIS	-	expression tag	UNP Q59523
A	-12	HIS	-	expression tag	UNP Q59523
A	-11	HIS	-	expression tag	UNP Q59523
A	-10	HIS	-	expression tag	UNP Q59523
A	-9	SER	-	expression tag	UNP Q59523
A	-8	SER	-	expression tag	UNP Q59523
A	-7	GLY	-	expression tag	UNP Q59523
A	-6	LEU	-	expression tag	UNP Q59523
A	-5	VAL	-	expression tag	UNP Q59523
A	-4	PRO	-	expression tag	UNP Q59523
A	-3	ARG	-	expression tag	UNP Q59523
A	-2	GLY	-	expression tag	UNP Q59523
A	-1	SER	-	expression tag	UNP Q59523
A	0	HIS	-	expression tag	UNP Q59523

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	
							0	0

- Molecule 3 is MYCINAMICIN IV (three-letter code: MIV) (formula: $C_{37}H_{61}NO_{11}$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O		
			49	37	1	11		
							0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	397	Total	O	0	0
			397	397		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: P-450-LIKE PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	56.28Å 162.03Å 50.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	162.02 – 1.65 53.17 – 1.65	Depositor EDS
% Data completeness (in resolution range)	84.7 (162.02-1.65) 84.8 (53.17-1.65)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.148 , 0.206 0.147 , 0.204	Depositor DCC
R_{free} test set	2430 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	20.3	Xtriage
Anisotropy	0.404	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3715	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, HEM, GOL, MIV, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.35	6/3270 (0.2%)	1.11	15/4450 (0.3%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	122	VAL	CB-CG2	6.20	1.65	1.52
1	A	71	GLU	CB-CG	-5.58	1.41	1.52
1	A	149	GLU	CB-CG	-5.47	1.41	1.52
1	A	358	VAL	CB-CG1	5.42	1.64	1.52
1	A	49	TYR	CE1-CZ	5.15	1.45	1.38
1	A	49	TYR	CG-CD2	5.01	1.45	1.39

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	96	ARG	NE-CZ-NH1	9.88	125.24	120.30
1	A	96	ARG	NE-CZ-NH2	-8.73	115.94	120.30
1	A	78	MET	CG-SD-CE	8.34	113.54	100.20
1	A	360	LEU	CB-CG-CD1	7.07	123.01	111.00
1	A	36	ARG	NE-CZ-NH1	6.50	123.55	120.30
1	A	58	ASP	CB-CG-OD1	6.09	123.78	118.30
1	A	58	ASP	CB-CG-OD2	-5.90	112.99	118.30
1	A	288	ARG	NE-CZ-NH2	-5.72	117.44	120.30
1	A	149	GLU	CA-CB-CG	5.65	125.83	113.40
1	A	297	ARG	NE-CZ-NH2	-5.63	117.48	120.30
1	A	95	ARG	NE-CZ-NH2	-5.60	117.50	120.30
1	A	93	ARG	CG-CD-NE	-5.49	100.28	111.80
1	A	106	ARG	NE-CZ-NH1	5.24	122.92	120.30
1	A	346	CYS	CA-CB-SG	-5.23	104.59	114.00
1	A	293	ASP	CB-CG-OD1	5.08	122.87	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3203	0	3163	46	0
2	A	43	0	30	2	0
3	A	49	0	61	0	0
4	A	1	0	0	0	0
5	A	10	0	0	0	0
6	A	12	0	16	1	0
7	A	397	0	0	3	0
All	All	3715	0	3270	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:285[B]:ALA:O	1:A:286[B]:PHE:HB2	1.40	1.12
1:A:67[A]:MET:HA	1:A:67[A]:MET:CE	1.87	1.02
1:A:67[A]:MET:HA	1:A:67[A]:MET:HE2	1.39	0.98
1:A:285[B]:ALA:O	1:A:286[B]:PHE:CB	2.14	0.94
1:A:67[A]:MET:CE	1:A:80:LYS:HB3	2.05	0.87
1:A:67[A]:MET:HE1	1:A:80:LYS:HB3	1.61	0.80
1:A:96:ARG:HD2	1:A:214[B]:ASP:OD2	1.83	0.78
1:A:67[A]:MET:CE	1:A:67[A]:MET:CA	2.67	0.71
1:A:15:HIS:HE1	1:A:383:GLU:OE2	1.74	0.70
1:A:67[A]:MET:HE1	1:A:80:LYS:CB	2.25	0.67
1:A:250:LEU:HD12	1:A:256:LEU:HG	1.77	0.67
1:A:67[A]:MET:HE2	1:A:80:LYS:HB3	1.78	0.65
1:A:192[B]:ASP:OD2	1:A:196:ARG:NH1	2.30	0.65
1:A:215[B]:GLN:CD	1:A:216:GLN:N	2.50	0.64
1:A:67[A]:MET:CE	1:A:80:LYS:CB	2.75	0.64
1:A:67[A]:MET:HA	1:A:67[A]:MET:HE3	1.79	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9:TYR:OH	1:A:284[B]:THR:HG23	2.03	0.59
1:A:67[A]:MET:CA	1:A:67[A]:MET:HE3	2.33	0.57
1:A:71:GLU:HG3	1:A:287:PRO:HG2	1.85	0.57
1:A:257:ARG:NH1	1:A:370:ILE:O	2.37	0.54
1:A:130:GLN:NE2	1:A:397:TRP:H	2.08	0.52
1:A:318:GLN:NE2	1:A:324:ALA:H	2.08	0.51
1:A:318:GLN:HE22	1:A:324:ALA:H	1.59	0.50
1:A:71:GLU:CG	1:A:287:PRO:HG2	2.42	0.50
1:A:13:ASP:OD1	1:A:15:HIS:HD2	1.95	0.49
1:A:67[A]:MET:HE2	1:A:80:LYS:CB	2.42	0.48
1:A:53:ARG:CD	7:A:2073:HOH:O	2.60	0.48
1:A:285[B]:ALA:O	2:A:450:HEM:O2A	2.31	0.48
1:A:9:TYR:OH	1:A:284[B]:THR:CG2	2.61	0.48
1:A:9:TYR:O	1:A:38:PRO:HD3	2.15	0.47
1:A:86:MET:SD	1:A:94:LEU:HD12	2.55	0.46
1:A:345[B]:HIS:HE1	7:A:2353:HOH:O	1.99	0.45
1:A:67[A]:MET:HE3	1:A:67[A]:MET:N	2.31	0.45
1:A:96:ARG:HD3	7:A:2234:HOH:O	2.17	0.44
1:A:347:LEU:C	1:A:347:LEU:HD12	2.38	0.43
1:A:15:HIS:CE1	1:A:383:GLU:OE2	2.64	0.42
1:A:389:GLY:C	6:A:1402:GOL:H32	2.40	0.42
1:A:113:ARG:HD2	1:A:113:ARG:HA	1.92	0.42
1:A:144:VAL:HG11	1:A:162:THR:HG23	2.01	0.42
1:A:235:GLY:HA2	2:A:450:HEM:C2C	2.55	0.41
1:A:380:ARG:HD3	1:A:392:GLU:OE1	2.20	0.41
1:A:9:TYR:CG	1:A:10:PRO:HA	2.56	0.41
1:A:278:VAL:HA	1:A:279:PRO:HD3	1.94	0.41
1:A:284[A]:THR:O	1:A:285[A]:ALA:HB3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	406/417 (97%)	392 (97%)	10 (2%)	4 (1%)	15 3

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	285[A]	ALA
1	A	285[B]	ALA
1	A	286[A]	PHE
1	A	286[B]	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	337/346 (97%)	325 (96%)	12 (4%)	35 11

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	67[A]	MET
1	A	67[B]	MET
1	A	80	LYS
1	A	105	ARG
1	A	155[A]	SER
1	A	155[B]	SER
1	A	180	GLN
1	A	211	GLN
1	A	250	LEU
1	A	279	PRO
1	A	360	LEU
1	A	365	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	HIS
1	A	130	GLN
1	A	211	GLN
1	A	318	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	450	7,1	42,50,50	2.45	14 (33%)	46,82,82	2.28	19 (41%)
5	SO4	A	1400	-	4,4,4	0.64	0	6,6,6	0.63	0
5	SO4	A	1399	-	4,4,4	0.40	0	6,6,6	0.78	0
6	GOL	A	1402	-	5,5,5	0.40	0	5,5,5	1.07	0
6	GOL	A	1401	-	5,5,5	0.71	0	5,5,5	0.81	0
3	MIV	A	460	-	51,51,51	1.37	3 (5%)	64,71,71	1.35	12 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MIV	A	460	-	-	6/55/91/91	0/2/3/3
6	GOL	A	1402	-	-	2/4/4/4	-
2	HEM	A	450	7,1	-	0/12/54/54	-
6	GOL	A	1401	-	-	2/4/4/4	-

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	450	HEM	C3D-C2D	7.12	1.52	1.36
3	A	460	MIV	O6-C24	6.24	1.47	1.34
2	A	450	HEM	C3C-C2C	-5.49	1.32	1.40
2	A	450	HEM	C3C-CAC	4.81	1.58	1.47
2	A	450	HEM	C3C-C4C	4.35	1.47	1.41
2	A	450	HEM	CMA-C3A	3.83	1.59	1.51
2	A	450	HEM	CHA-C4D	3.77	1.43	1.34
2	A	450	HEM	CHB-C1B	3.65	1.43	1.34
2	A	450	HEM	C4A-CHB	-3.06	1.32	1.41
2	A	450	HEM	CAA-C2A	3.02	1.59	1.52
2	A	450	HEM	CBA-CGA	2.90	1.57	1.50
3	A	460	MIV	C8-C9	2.82	1.58	1.54
3	A	460	MIV	O5-C24	2.70	1.26	1.21
2	A	450	HEM	CAB-C3B	2.69	1.54	1.47
2	A	450	HEM	CMC-C2C	2.46	1.57	1.51
2	A	450	HEM	CBD-CGD	2.28	1.55	1.50
2	A	450	HEM	C2C-C1C	2.22	1.47	1.42

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	450	HEM	CMA-C3A-C4A	-5.55	120.33	128.46
2	A	450	HEM	CMD-C2D-C1D	4.70	132.38	125.03
2	A	450	HEM	C3C-C4C-NC	-4.43	102.58	110.94
2	A	450	HEM	CMA-C3A-C2A	4.21	132.87	124.94
3	A	460	MIV	O2-C11-C20	-3.88	103.14	110.93
2	A	450	HEM	C2C-C3C-C4C	3.50	109.34	106.90
2	A	450	HEM	C1D-C2D-C3D	-3.40	103.41	106.98
2	A	450	HEM	CMC-C2C-C3C	3.39	131.46	124.68
2	A	450	HEM	CHB-C1B-NB	-3.08	120.56	124.37
2	A	450	HEM	CBD-CAD-C3D	-2.97	104.33	112.53
3	A	460	MIV	C37-C36-C35	2.84	118.27	113.08
2	A	450	HEM	CMB-C2B-C1B	2.82	129.44	125.03

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	450	HEM	C2D-C1D-ND	2.77	113.10	109.90
3	A	460	MIV	C20-C22-C23	-2.69	119.18	126.30
2	A	450	HEM	C2B-C1B-NB	2.69	112.94	109.84
2	A	450	HEM	C3B-C4B-NB	2.63	111.36	109.47
2	A	450	HEM	C4B-CHC-C1C	2.61	126.01	122.56
2	A	450	HEM	CAD-C3D-C4D	2.53	129.11	124.70
2	A	450	HEM	C1B-NB-C4B	-2.50	102.25	105.21
2	A	450	HEM	C4C-CHD-C1D	2.48	125.83	122.56
2	A	450	HEM	C3B-C2B-C1B	-2.48	104.55	106.41
3	A	460	MIV	C17-C14-N	-2.42	108.78	115.59
3	A	460	MIV	C6-C8-C9	2.41	119.41	115.09
3	A	460	MIV	C28-C1-C2	-2.38	119.81	125.40
3	A	460	MIV	C25-O6-C24	2.18	120.71	117.50
3	A	460	MIV	C18-C17-C14	2.16	114.14	110.46
3	A	460	MIV	O3-C13-C14	2.15	113.81	109.87
3	A	460	MIV	O6-C24-O5	-2.14	119.91	123.34
3	A	460	MIV	O10-C35-C36	2.13	114.44	109.74
3	A	460	MIV	O11-C30-O7	-2.06	105.17	110.04
2	A	450	HEM	CBA-CAA-C2A	-2.00	109.17	112.54

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	460	MIV	O11-C30-O7-C29
3	A	460	MIV	C31-C30-O7-C29
6	A	1401	GOL	O1-C1-C2-C3
6	A	1402	GOL	C1-C2-C3-O3
6	A	1402	GOL	O2-C2-C3-O3
3	A	460	MIV	C28-C29-O7-C30
6	A	1401	GOL	O1-C1-C2-O2
3	A	460	MIV	O6-C25-C26-C27
3	A	460	MIV	C2-C1-C28-C29
3	A	460	MIV	C33-C31-O8-C32


There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	450	HEM	2	0
6	A	1402	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	393/417 (94%)	-0.27	3 (0%) 86 88	12, 18, 30, 42	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	285[A]	ALA	6.6
1	A	216	GLN	2.4
1	A	177	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MIV	A	460	49/49	0.89	0.15	22,29,38,40	13
6	GOL	A	1402	6/6	0.96	0.10	18,22,25,26	0
4	CL	A	1398	1/1	0.97	0.05	44,44,44,44	0
5	SO4	A	1399	5/5	0.98	0.14	41,44,47,47	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	SO4	A	1400	5/5	0.98	0.07	22,22,28,29	0
6	GOL	A	1401	6/6	0.98	0.08	12,14,16,17	0
2	HEM	A	450	43/43	0.98	0.09	10,12,16,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.