



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 12, 2025 – 02:16 PM JST

PDB ID : 8YDL / pdb\_00008ydl  
Title : Crystal structure of single-chain dimer for RSV F protein  
Authors : Qi, J.X.; Li, J.; Zhao, J.R.; Guo, W.J.  
Deposited on : 2024-02-20  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.006 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.45.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3775 (2.60-2.60)
Clashscore	180529	4181 (2.60-2.60)
Ramachandran outliers	177936	4129 (2.60-2.60)
Sidechain outliers	177891	4129 (2.60-2.60)
RSRZ outliers	164620	3775 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	491	<div> <div>4%</div> <div> <div></div> <div>77%</div> <div>13%</div> <div>• 9%</div> </div> </div>
1	B	491	<div> <div>3%</div> <div> <div></div> <div>76%</div> <div>14%</div> <div>• 9%</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7090 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fusion glycoprotein F2, F1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	446	Total	C	N	O	S	0	4	0
			3456	2190	574	670	22			
1	B	446	Total	C	N	O	S	0	5	0
			3461	2193	575	671	22			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	89	Total	O	0	0
			89	89		
2	B	84	Total	O	0	0
			84	84		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fusion glycoprotein F2, F1



- Molecule 1: Fusion glycoprotein F2, F1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.66Å 59.91Å 102.42Å 76.38° 73.70° 76.77°	Depositor
Resolution (Å)	46.44 – 2.60 46.44 – 2.60	Depositor EDS
% Data completeness (in resolution range)	95.9 (46.44-2.60) 86.0 (46.44-2.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.61 (at 2.61Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.234 , 0.265 0.245 , 0.275	Depositor DCC
$R_{free}$ test set	1880 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.2	Xtriage
Anisotropy	0.105	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 20.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.118 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	7090	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 98.08 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.6076e-11. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.20	0/3504	0.42	2/4746 (0.0%)
1	B	0.20	0/3512	0.41	2/4757 (0.0%)
All	All	0.20	0/7016	0.42	4/9503 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	47	ALA	CA-C-N	-5.93	113.24	122.37
1	B	47	ALA	C-N-CA	-5.93	113.24	122.37
1	A	288	SER	CA-C-N	-5.87	112.77	122.39
1	A	288	SER	C-N-CA	-5.87	112.77	122.39

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3456	0	3628	34	0
1	B	3461	0	3634	38	0
2	A	89	0	0	1	0
2	B	84	0	0	3	0
All	All	7090	0	7262	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:77:LYS:O	1:B:81:GLN:HG3	1.89	0.72
1:A:46:SER:HB3	1:A:245:GLN:HE22	1.54	0.71
1:A:228:THR:HB	1:A:231:GLN:HG3	1.77	0.66
1:A:46:SER:CB	1:A:245:GLN:HE22	2.09	0.65
1:B:379:LYS:HE3	2:B:673:HOH:O	1.96	0.65
1:A:46:SER:C	1:A:48:LEU:H	2.08	0.62
1:A:228:THR:HG22	1:A:231:GLN:H	1.64	0.61
1:A:240:GLN:H	1:A:240:GLN:CD	2.09	0.60
1:A:460:LEU:HD23	1:A:481:VAL:HG13	1.86	0.58
1:B:336:GLN:O	1:B:340:GLN:HG3	2.04	0.57
1:B:227:ILE:HG13	1:B:232:LYS:HG3	1.85	0.57
1:B:482:GLN:NE2	2:B:601:HOH:O	2.29	0.56
1:A:319:LYS:O	1:A:323:GLN:HG3	2.06	0.56
1:B:240:GLN:H	1:B:240:GLN:CD	2.14	0.56
1:A:228:THR:HG23	2:A:660:HOH:O	2.05	0.55
1:B:460:LEU:HD23	1:B:481:VAL:HG13	1.89	0.55
1:B:302:GLU:HB2	1:B:397:ASP:HA	1.90	0.54
1:B:319:LYS:O	1:B:323:GLN:HG3	2.08	0.54
1:A:235:MET:HE2	1:A:267:TYR:HD2	1.72	0.53
1:A:77:LYS:O	1:A:81:GLN:HG3	2.08	0.53
1:A:471:ASN:OD1	1:B:136:ASN:ND2	2.42	0.53
1:A:336:GLN:O	1:A:340:GLN:HG3	2.09	0.52
1:B:315:ASP:OD2	1:B:317:LYS:NZ	2.42	0.52
1:A:304:SER:HB2	1:A:399:LYS:HA	1.92	0.52
1:A:93:LEU:O	1:A:97:MET:HG3	2.10	0.52
1:B:335:LEU:HG	1:B:437:THR:HG23	1.93	0.51
1:A:235:MET:HE2	1:A:267:TYR:CD2	2.47	0.50
1:A:61:LEU:HD11	1:A:258:LEU:HB2	1.94	0.50
1:A:106:GLY:O	1:A:291:ARG:NH1	2.40	0.49
1:A:228:THR:CB	1:A:231:GLN:HG3	2.43	0.48
1:A:171:GLN:HB2	1:A:174:SER:OG	2.14	0.48
1:B:171:GLN:HB2	1:B:174:SER:OG	2.13	0.48
1:B:235:MET:HE2	1:B:267:TYR:CD2	2.49	0.48
1:B:321:ILE:HD11	1:B:423:VAL:HA	1.95	0.47
1:B:64:ILE:HD12	1:B:165:LEU:HD11	1.95	0.47
1:B:228:THR:OG1	1:B:231:GLN:HG3	2.14	0.47
1:A:483:ILE:O	1:A:487:GLN:HG2	2.15	0.46
1:A:411:ASN:O	1:A:412:LYS:HB2	2.16	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:228:THR:CG2	1:A:230:ASP:H	2.29	0.46
1:A:470:THR:OG1	1:A:473:GLN:HG3	2.16	0.46
1:B:235:MET:HE2	1:B:267:TYR:HD2	1.82	0.45
1:B:304:SER:HB2	1:B:399:LYS:HA	1.98	0.45
1:A:228:THR:HG23	1:A:230:ASP:H	1.82	0.45
1:B:303:LEU:O	1:B:399:LYS:HB2	2.16	0.45
1:B:73:ASP:O	1:B:77:LYS:HG3	2.17	0.45
1:A:64:ILE:O	1:A:64:ILE:HG13	2.17	0.44
1:B:483:ILE:O	1:B:487:GLN:HG2	2.17	0.44
1:A:420:ILE:O	1:A:424:ILE:HG12	2.17	0.44
1:B:305:ASN:N	2:B:603:HOH:O	2.49	0.44
1:B:311:CYS:HB2	1:B:414:SER:HA	1.98	0.44
1:A:158:ASN:O	1:A:162:LYS:HB2	2.17	0.44
1:B:160:ILE:HA	1:B:164:LEU:HB3	1.99	0.44
1:B:82:GLU:HA	1:B:85:LYS:HD2	1.99	0.44
1:A:335:LEU:HG	1:A:437:THR:HG23	2.01	0.43
1:B:60:GLU:HB2	1:B:155:ASP:HA	2.00	0.43
1:B:470:THR:OG1	1:B:473:GLN:HG3	2.18	0.43
1:B:298:VAL:HB	1:B:392:THR:HG22	2.01	0.43
1:B:494:ILE:HD11	1:B:496:LYS:HD3	2.00	0.43
1:A:421:GLU:OE1	1:A:421:GLU:N	2.39	0.42
1:B:303:LEU:HD11	1:B:500:LEU:HB2	2.01	0.42
1:B:463:LEU:HD22	1:B:506:LEU:HD22	2.00	0.42
1:A:504:VAL:HG12	1:A:506:LEU:CD1	2.50	0.42
1:B:46:SER:CB	1:B:245:GLN:HE22	2.33	0.41
1:B:221:LEU:HD22	1:B:264:LEU:HD22	2.01	0.41
1:A:430:ASN:O	1:A:434:LEU:HG	2.21	0.41
1:B:66:GLU:H	1:B:66:GLU:HG2	1.71	0.41
1:B:312:ASN:HB2	1:B:414:SER:HB3	2.03	0.41
1:A:221:LEU:HD22	1:A:264:LEU:HD22	2.02	0.41
1:B:178:ILE:O	1:B:182:ILE:HG12	2.21	0.41
1:A:227:ILE:HG13	1:A:232:LYS:HG3	2.03	0.40
1:B:504:VAL:HG12	1:B:506:LEU:CD1	2.51	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	446/491 (91%)	433 (97%)	10 (2%)	3 (1%)	19	38
1	B	447/491 (91%)	432 (97%)	13 (3%)	2 (0%)	30	52
All	All	893/982 (91%)	865 (97%)	23 (3%)	5 (1%)	22	43

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	412	LYS
1	A	413	GLN
1	B	412	LYS
1	B	413	GLN
1	A	47	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	410/445 (92%)	393 (96%)	17 (4%)	26	51
1	B	411/445 (92%)	403 (98%)	8 (2%)	52	75
All	All	821/890 (92%)	796 (97%)	25 (3%)	35	63

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	95	LEU
1	A	99	SER
1	A	134	SER
1	A	173	CYS
1	A	174	SER
1	A	228	THR
1	A	262	VAL
1	A	329	LYS
1	A	364	GLU
1	A	372	SER
1	A	374	LEU
1	A	376	SER
1	A	400	ASN
1	A	405	GLN
1	A	415	CYS
1	A	465	ASN
1	B	46	SER
1	B	64	ILE
1	B	99	SER
1	B	173	CYS
1	B	400	ASN
1	B	412	LYS
1	B	415	CYS
1	B	448	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	81	GLN
1	A	245	GLN
1	A	340	GLN
1	A	465	ASN
1	A	486	GLN
1	B	223	ASN
1	B	323	GLN
1	B	471	ASN
1	B	473	GLN
1	B	482	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	446/491 (90%)	-1.14	19 (4%) 40 34	9, 38, 103, 194	4 (0%)
1	B	446/491 (90%)	-1.13	14 (3%) 51 46	13, 38, 110, 208	5 (1%)
All	All	892/982 (90%)	-1.13	33 (3%) 45 39	9, 38, 111, 208	9 (1%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	413	GLN	4.9
1	B	346	GLY	4.3
1	A	343	PRO	3.7
1	B	316	ALA	3.3
1	B	344	ALA	3.3
1	B	348	GLY	3.2
1	A	414	SER	3.1
1	A	413	GLN	3.1
1	B	311	CYS	3.0
1	A	72	THR	2.9
1	A	100	THR	2.8
1	A	345	THR	2.7
1	B	410	LEU	2.7
1	A	348	GLY	2.6
1	B	70	ASN	2.6
1	A	346	GLY	2.6
1	A	347	SER	2.5
1	A	73	ASP	2.5
1	B	100	THR	2.4
1	A	342	THR	2.3
1	A	410	LEU	2.3
1	A	344	ALA	2.2
1	A	311	CYS	2.1
1	B	345	THR	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	73	ASP	2.1
1	B	72	THR	2.1
1	A	171	GLN	2.1
1	A	313	GLY	2.1
1	B	414	SER	2.1
1	A	418	PRO	2.0
1	A	315	ASP	2.0
1	A	316	ALA	2.0
1	B	315	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.