



## wwPDB EM Validation Summary Report ⓘ

Oct 13, 2024 – 09:20 am BST

PDB ID : 6YEJ  
EMDB ID : EMD-10793  
Title : Cryo-EM structure of the Full-length disease type human Huntingtin  
Authors : Tame, G.; Jung, T.; Dal Perraro, M.; Hebert, H.; Song, J.  
Deposited on : 2020-03-24  
Resolution : 18.20 Å(reported)  
Based on initial model : 6EZ8

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113  
MolProbity : 4.02b-467  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 18.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	3199	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 15999 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Huntingtin.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	2046	Total	C	N	O	S	0	0
			15999	10276	2718	2907	98		

There are 76 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	MET	-	initiating methionine	UNP P42858
A	-4	ALA	-	expression tag	UNP P42858
A	-3	THR	-	expression tag	UNP P42858
A	-2	LEU	-	expression tag	UNP P42858
A	-1	GLU	-	expression tag	UNP P42858
A	0	LYS	-	expression tag	UNP P42858
A	1	LEU	-	expression tag	UNP P42858
A	2	MET	-	expression tag	UNP P42858
A	3	LYS	-	expression tag	UNP P42858
A	4	ALA	-	expression tag	UNP P42858
A	5	PHE	-	expression tag	UNP P42858
A	6	GLU	-	expression tag	UNP P42858
A	7	SER	-	expression tag	UNP P42858
A	8	LEU	-	expression tag	UNP P42858
A	9	LYS	-	expression tag	UNP P42858
A	10	SER	-	expression tag	UNP P42858
A	11	PHE	-	expression tag	UNP P42858
A	12	GLN	-	expression tag	UNP P42858
A	13	GLN	-	expression tag	UNP P42858
A	14	GLN	-	expression tag	UNP P42858
A	15	GLN	-	expression tag	UNP P42858
A	16	GLN	-	expression tag	UNP P42858
A	17	GLN	-	expression tag	UNP P42858
A	18	GLN	-	expression tag	UNP P42858
A	19	GLN	-	expression tag	UNP P42858
A	20	GLN	-	expression tag	UNP P42858
A	21	GLN	-	expression tag	UNP P42858
A	22	GLN	-	expression tag	UNP P42858

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	23	GLN	-	expression tag	UNP P42858
A	24	GLN	-	expression tag	UNP P42858
A	25	GLN	-	expression tag	UNP P42858
A	26	GLN	-	expression tag	UNP P42858
A	27	GLN	-	expression tag	UNP P42858
A	28	GLN	-	expression tag	UNP P42858
A	29	GLN	-	expression tag	UNP P42858
A	30	GLN	-	expression tag	UNP P42858
A	31	GLN	-	expression tag	UNP P42858
A	32	GLN	-	expression tag	UNP P42858
A	33	GLN	-	expression tag	UNP P42858
A	34	GLN	-	expression tag	UNP P42858
A	35	GLN	-	expression tag	UNP P42858
A	36	GLN	-	expression tag	UNP P42858
A	37	GLN	-	expression tag	UNP P42858
A	38	GLN	-	expression tag	UNP P42858
A	39	GLN	-	expression tag	UNP P42858
A	40	GLN	-	expression tag	UNP P42858
A	41	GLN	-	expression tag	UNP P42858
A	42	GLN	-	expression tag	UNP P42858
A	43	GLN	-	expression tag	UNP P42858
A	44	GLN	-	expression tag	UNP P42858
A	45	GLN	-	expression tag	UNP P42858
A	46	GLN	-	expression tag	UNP P42858
A	47	GLN	-	expression tag	UNP P42858
A	48	GLN	-	expression tag	UNP P42858
A	49	GLN	-	expression tag	UNP P42858
A	50	GLN	-	expression tag	UNP P42858
A	51	GLN	-	expression tag	UNP P42858
A	52	GLN	-	expression tag	UNP P42858
A	53	GLN	-	expression tag	UNP P42858
A	54	GLN	-	expression tag	UNP P42858
A	55	GLN	-	expression tag	UNP P42858
A	56	GLN	-	expression tag	UNP P42858
A	57	GLN	-	expression tag	UNP P42858
A	58	GLN	-	expression tag	UNP P42858
A	59	GLN	-	expression tag	UNP P42858
A	60	GLN	-	expression tag	UNP P42858
A	61	GLN	-	expression tag	UNP P42858
A	62	GLN	-	expression tag	UNP P42858
A	63	GLN	-	expression tag	UNP P42858
A	64	GLN	-	expression tag	UNP P42858


*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	65	GLN	-	expression tag	UNP P42858
A	66	GLN	-	expression tag	UNP P42858
A	67	GLN	-	expression tag	UNP P42858
A	68	GLN	-	expression tag	UNP P42858
A	1289	ARG	LYS	conflict	UNP P42858
A	2360	HIS	TYR	conflict	UNP P42858









## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	21781	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	33.6	Depositor
Minimum defocus (nm)	400	Depositor
Maximum defocus (nm)	400	Depositor
Magnification	47170	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.016	Depositor
Minimum map value	-0.001	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.0058	Depositor
Map size (Å)	254.4, 254.4, 254.4	wwPDB
Map dimensions	240, 240, 240	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.06, 1.06, 1.06	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.52	18/16315 (0.1%)	0.67	24/22179 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1768	GLY	C-O	-14.52	1.00	1.23
1	A	2980	GLY	C-O	-14.50	1.00	1.23
1	A	1373	THR	C-O	-12.07	1.00	1.23
1	A	2379	SER	C-O	-12.07	1.00	1.23
1	A	2526	LEU	C-O	-12.07	1.00	1.23

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1405	LEU	CA-CB-CG	7.78	133.20	115.30
1	A	2980	GLY	CA-C-O	-6.43	109.02	120.60
1	A	1768	GLY	CA-C-O	-6.42	109.04	120.60
1	A	907	LEU	CA-CB-CG	5.99	129.07	115.30
1	A	1554	LEU	CA-CB-CG	5.74	128.50	115.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1100	VAL	Peptide
1	A	1369	THR	Peptide
1	A	2323	SER	Peptide
1	A	351	VAL	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	15999	0	16385	185	0
All	All	15999	0	16385	185	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 185 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1157:ARG:HD3	1:A:2679:ILE:CD1	1.17	1.64
1:A:1157:ARG:HD2	1:A:2679:ILE:CG2	1.10	1.58
1:A:1157:ARG:CD	1:A:2679:ILE:CD1	1.87	1.48
1:A:1157:ARG:CD	1:A:2679:ILE:HG21	1.44	1.45
1:A:1157:ARG:CZ	1:A:2679:ILE:HD13	1.47	1.42

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	2010/3199 (63%)	1872 (93%)	133 (7%)	5 (0%)	44 78

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2569	VAL
1	A	332	SER
1	A	2568	PRO
1	A	1101	PRO
1	A	2321	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	1800/2824 (64%)	1784 (99%)	16 (1%)	75 83

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2851	LYS
1	A	2583	ARG
1	A	2211	GLN
1	A	2567	VAL
1	A	2188	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2291	HIS
1	A	2667	GLN
1	A	3140	ASN
1	A	3033	GLN
1	A	2618	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-10793. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)

#### 6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map



X Index: 120



Y Index: 120




Z Index: 120

The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)

### 6.3.1 Primary map



X Index: 160



Y Index: 134




Z Index: 110


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0058. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

## 6.6 Mask visualisation [i](#)


This section was not generated. No masks/segmentation were deposited.



## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 466  $\text{nm}^3$ ; this corresponds to an approximate mass of 421 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of 0.055 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.055  $\text{\AA}^{-1}$

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	18.20	-	-
Author-provided FSC curve	17.09	20.70	17.99
Unmasked-calculated*	-	-	-

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-10793 and PDB model 6YEJ. Per-residue inclusion information can be found in section [3](#) on page [6](#).


### 9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.0058 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0058).

## 9.4 Atom inclusion [i](#)



At the recommended contour level, 82% of all backbone atoms, 81% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0058) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div><div></div></div> 0.8100	<div><div></div></div> 0.0360
A	<div><div></div></div> 0.8100	<div><div></div></div> 0.0360

