



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2024 – 12:23 PM EDT

PDB ID : 4YV6
Title : X-ray crystal structure of Streptococcus dysgalactiae SHP pheromone receptor Rgg2
Authors : Neiditch, M.B.; Parashar, V.
Deposited on : 2015-03-19
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2684 (2.08-2.04)
Clashscore	141614	2801 (2.08-2.04)
Ramachandran outliers	138981	2768 (2.08-2.04)
Sidechain outliers	138945	2768 (2.08-2.04)
RSRZ outliers	127900	2646 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	284	
1	B	284	
1	C	284	
1	D	284	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 18109 atoms, of which 8860 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulator.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	260	Total	C	H	N	O	S	0	0	0
			4343	1401	2173	353	410	6			
1	B	271	Total	C	H	N	O	S	0	0	0
			4514	1455	2253	371	429	6			
1	C	267	Total	C	H	N	O	S	0	0	0
			4451	1437	2221	366	421	6			
1	D	266	Total	C	H	N	O	S	0	0	0
			4432	1430	2213	361	422	6			

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	110	Total	O	0	0
			110	110		
3	B	65	Total	O	0	0
			65	65		
3	C	75	Total	O	0	0
			75	75		
3	D	84	Total	O	0	0
			84	84		

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	68.10Å 99.08Å 99.89Å 90.00° 97.03° 90.00°	Depositor
Resolution (Å)	39.96 – 2.05 39.96 – 2.05	Depositor EDS
% Data completeness (in resolution range)	93.8 (39.96-2.05) 93.6 (39.96-2.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.64 (at 2.05Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.208 , 0.243 0.212 , 0.245	Depositor DCC
R_{free} test set	2000 reflections (2.45%)	wwPDB-VP
Wilson B-factor (Å ²)	37.2	Xtriage
Anisotropy	0.514	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 44.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	18109	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	0/2211	0.37	0/2975
1	B	0.22	0/2307	0.36	0/3108
1	C	0.22	0/2275	0.34	0/3064
1	D	0.23	0/2262	0.35	0/3046
All	All	0.23	0/9055	0.35	0/12193

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2170	2173	2173	4	0
1	B	2261	2253	2253	6	0
1	C	2230	2221	2221	5	0
1	D	2219	2213	2213	7	0
2	B	10	0	0	0	0
2	C	5	0	0	0	0
2	D	20	0	0	1	0
3	A	110	0	0	0	0
3	B	65	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	75	0	0	0	0
3	D	84	0	0	1	0
All	All	9249	8860	8860	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:31:SER:OG	1:B:35:ARG:NH1	2.27	0.68
1:C:81:ARG:NH2	1:D:175:GLU:OE1	2.30	0.64
1:D:22:SER:O	3:D:401:HOH:O	2.15	0.64
1:D:31:SER:OG	1:D:35:ARG:NH1	2.31	0.64
1:C:245:ASP:OD1	1:C:248:ARG:NH2	2.32	0.62
1:A:138:GLU:OE1	1:A:139:GLN:N	2.40	0.55
1:A:158:ASN:OD1	1:D:203:HIS:NE2	2.42	0.51
1:C:268:GLU:OE2	1:C:272:LYS:NZ	2.43	0.51
1:B:65:THR:O	1:B:67:SER:N	2.41	0.51
1:C:40:GLU:OE1	1:C:41:SER:N	2.45	0.49
1:D:34:SER:OG	1:D:38:ARG:NH1	2.46	0.48
1:A:245:ASP:OD1	1:A:248:ARG:NH2	2.44	0.47
1:A:35:ARG:O	1:A:36:PHE:HB3	2.15	0.46
1:D:202:ASP:OD1	1:D:202:ASP:N	2.46	0.45
1:B:202:ASP:N	1:B:202:ASP:OD1	2.51	0.43
1:D:35:ARG:NH2	2:D:301:SO4:O2	2.51	0.43
1:B:105:GLU:OE1	1:B:105:GLU:N	2.53	0.42
1:B:245:ASP:OD1	1:B:248:ARG:NH2	2.53	0.41
1:C:125:GLU:OE1	1:C:125:GLU:N	2.49	0.41
1:B:8:THR:HG21	1:B:66:HIS:HB2	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	254/284 (89%)	249 (98%)	3 (1%)	2 (1%)	19	9
1	B	269/284 (95%)	258 (96%)	10 (4%)	1 (0%)	34	25
1	C	263/284 (93%)	256 (97%)	6 (2%)	1 (0%)	34	25
1	D	262/284 (92%)	252 (96%)	9 (3%)	1 (0%)	34	25
All	All	1048/1136 (92%)	1015 (97%)	28 (3%)	5 (0%)	29	19

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	219	LEU
1	A	36	PHE
1	C	219	LEU
1	A	219	LEU
1	D	63	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/265 (92%)	241 (99%)	3 (1%)	71	69
1	B	254/265 (96%)	251 (99%)	3 (1%)	71	69
1	C	250/265 (94%)	247 (99%)	3 (1%)	71	69
1	D	249/265 (94%)	247 (99%)	2 (1%)	81	81
All	All	997/1060 (94%)	986 (99%)	11 (1%)	73	72

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LEU
1	A	71	THR
1	A	138	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	117	GLU
1	B	168	VAL
1	B	274	VAL
1	C	10	ARG
1	C	30	LYS
1	C	113	LEU
1	D	4	GLU
1	D	241	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	D	301	-	4,4,4	0.13	0	6,6,6	0.10	0
2	SO4	B	301	-	4,4,4	0.14	0	6,6,6	0.04	0
2	SO4	C	301	-	4,4,4	0.14	0	6,6,6	0.06	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	D	302	-	4,4,4	0.14	0	6,6,6	0.06	0
2	SO4	B	302	-	4,4,4	0.14	0	6,6,6	0.05	0
2	SO4	D	303	-	4,4,4	0.15	0	6,6,6	0.05	0
2	SO4	D	304	-	4,4,4	0.13	0	6,6,6	0.05	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	301	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	260/284 (91%)	0.63	21 (8%)	12 12	26, 42, 88, 126	0
1	B	271/284 (95%)	0.79	24 (8%)	9 10	34, 51, 87, 105	0
1	C	267/284 (94%)	0.73	34 (12%)	3 3	30, 49, 88, 114	0
1	D	266/284 (93%)	0.57	14 (5%)	26 27	30, 46, 76, 104	0
All	All	1064/1136 (93%)	0.68	93 (8%)	10 10	26, 48, 87, 126	0

All (93) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	36	PHE	7.0
1	A	38	ARG	5.9
1	D	271	ARG	5.8
1	D	241	VAL	5.1
1	A	40	GLU	4.9
1	C	198	TYR	4.8
1	C	28	LEU	4.7
1	D	272	LYS	4.6
1	B	94	LYS	4.5
1	C	241	VAL	4.4
1	A	42	GLU	4.4
1	B	201	PHE	4.3
1	C	272	LYS	4.1
1	A	241	VAL	4.0
1	B	236	LEU	3.9
1	D	274	VAL	3.9
1	A	33	ILE	3.9
1	C	11	ARG	3.6
1	C	20	ILE	3.6
1	A	37	GLU	3.6
1	A	31	SER	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	239	GLY	3.4
1	C	10	ARG	3.3
1	C	239	GLY	3.3
1	C	274	VAL	3.3
1	B	116	ILE	3.2
1	C	21	SER	3.2
1	A	20	ILE	3.1
1	B	202	ASP	3.1
1	C	39	GLY	3.1
1	C	176	GLN	3.0
1	A	11	ARG	2.9
1	A	19	SER	2.9
1	A	18	VAL	2.8
1	C	236	LEU	2.8
1	B	99	TYR	2.8
1	C	41	SER	2.7
1	B	69	THR	2.7
1	A	35	ARG	2.6
1	A	22	SER	2.6
1	A	30	LYS	2.6
1	C	33	ILE	2.6
1	D	4	GLU	2.6
1	C	200	TYR	2.5
1	A	21	SER	2.5
1	A	23	LEU	2.5
1	C	37	GLU	2.5
1	C	38	ARG	2.5
1	A	41	SER	2.5
1	A	9	LEU	2.4
1	D	219	LEU	2.4
1	C	35	ARG	2.4
1	C	199	SER	2.4
1	C	26	GLU	2.4
1	B	92	LEU	2.4
1	D	244	LYS	2.4
1	B	199	SER	2.4
1	C	219	LEU	2.4
1	C	42	GLU	2.4
1	D	65	THR	2.3
1	D	268	GLU	2.3
1	B	200	TYR	2.3
1	A	65	THR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	205	HIS	2.3
1	B	183	LEU	2.3
1	B	274	VAL	2.3
1	C	238	GLN	2.3
1	C	66	HIS	2.2
1	C	24	ALA	2.2
1	B	198	TYR	2.2
1	B	97	GLU	2.2
1	B	271	ARG	2.2
1	B	10	ARG	2.2
1	C	271	ARG	2.2
1	D	264	TYR	2.2
1	D	7	LYS	2.2
1	C	9	LEU	2.2
1	B	219	LEU	2.1
1	B	89	VAL	2.1
1	D	267	LYS	2.1
1	A	28	LEU	2.1
1	B	100	ALA	2.1
1	C	40	GLU	2.1
1	C	70	HIS	2.1
1	B	68	LYS	2.1
1	C	7	LYS	2.0
1	B	84	TYR	2.0
1	C	29	SER	2.0
1	D	248	ARG	2.0
1	C	244	LYS	2.0
1	C	25	ASP	2.0
1	B	222	TYR	2.0
1	B	5	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	D	302	5/5	0.89	0.27	56,57,62,67	0
2	SO4	D	301	5/5	0.90	0.19	56,60,61,67	0
2	SO4	B	302	5/5	0.93	0.14	53,57,62,65	0
2	SO4	C	301	5/5	0.95	0.10	52,57,60,64	0
2	SO4	B	301	5/5	0.96	0.16	51,53,55,58	0
2	SO4	D	303	5/5	0.96	0.12	53,53,54,60	0
2	SO4	D	304	5/5	0.97	0.10	52,55,57,59	0

6.5 Other polymers [i](#)

There are no such residues in this entry.