



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 25, 2025 – 01:26 PM JST

PDB ID : 8Z2C  
Title : Crystal structure of apo Aspergillus terreus glutamate dehydrogenase (At-GDH) with open and partially closed conformations (form I)  
Authors : Godsora, B.K.J.; Bhaumik, P.  
Deposited on : 2024-04-12  
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.41.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2096 (2.04-2.04)
Clashscore	180529	2229 (2.04-2.04)
Ramachandran outliers	177936	2217 (2.04-2.04)
Sidechain outliers	177891	2217 (2.04-2.04)
RSRZ outliers	164620	2096 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	460	<div> <div>3%</div> <div> <div></div> <div>84%</div> <div>14%</div> <div>.</div> </div> </div>
1	B	460	<div> <div>5%</div> <div> <div></div> <div>84%</div> <div>15%</div> <div>.</div> </div> </div>

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 7538 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	459	Total	C	N	O	S	0	19	0
			3599	2265	628	690	16			
1	A	458	Total	C	N	O	S	0	23	0
			3627	2289	631	694	13			

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	135	Total 135	O 135	0	0
3	A	161	Total 161	O 161	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamate dehydrogenase



- Molecule 1: Glutamate dehydrogenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	156.12Å 156.12Å 106.07Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.77 – 2.05 29.77 – 2.05	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.77-2.05) 99.9 (29.77-2.05)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.21 (at 2.04Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.176 , 0.203 0.202 , 0.230	Depositor DCC
$R_{free}$ test set	4683 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.5	Xtriage
Anisotropy	0.058	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 36.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.54$ , $\langle L^2 \rangle = 0.38$	Xtriage
Estimated twinning fraction	0.000 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	7538	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.88	0/3697	0.96	2/5002 (0.0%)
1	B	0.86	1/3669 (0.0%)	0.96	4/4962 (0.1%)
All	All	0.87	1/7366 (0.0%)	0.96	6/9964 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	128	GLU	CD-OE2	-5.11	1.20	1.25

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	131	ARG	NE-CZ-NH2	-10.42	115.09	120.30
1	B	45	ARG	NE-CZ-NH2	-8.59	116.01	120.30
1	A	131	ARG	NE-CZ-NH1	8.47	124.53	120.30
1	B	45	ARG	NE-CZ-NH1	6.66	123.63	120.30
1	B	142	ARG	NE-CZ-NH2	-5.59	117.51	120.30
1	B	50	ARG	NE-CZ-NH1	5.50	123.05	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3627	0	3568	59	0
1	B	3599	0	3531	56	0
2	A	8	0	6	0	0
2	B	8	0	6	1	0
3	A	161	0	0	12	0
3	B	135	0	0	11	0
All	All	7538	0	7111	112	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (112) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:36:ALA:O	1:A:40[A]:VAL:HG23	1.62	0.99
1:A:403:GLU:HB3	3:A:744:HOH:O	1.77	0.84
1:A:25:THR:HG23	3:A:605:HOH:O	1.82	0.78
1:A:40[B]:VAL:HG12	1:A:453:MET:SD	2.24	0.77
1:B:396:ARG:O	3:B:601:HOH:O	2.01	0.77
1:B:78:LYS:HG2	3:B:695:HOH:O	1.85	0.76
1:A:246:ARG:NH2	3:A:601:HOH:O	2.21	0.73
1:A:236:ALA:O	1:A:240[B]:VAL:HG22	1.87	0.73
1:B:126:ASP:HB2	3:B:718:HOH:O	1.89	0.72
1:B:45:ARG:HD3	1:A:139:GLU:OE1	1.94	0.68
1:B:351[A]:GLN:HA	1:B:351[A]:GLN:OE1	1.93	0.67
1:B:106:THR:HA	1:B:415[A]:CYS:SG	2.37	0.65
1:B:281:ASP:C	1:B:282[A]:ARG:HD2	2.17	0.65
1:A:259:VAL:HG22	3:A:629:HOH:O	1.95	0.64
1:A:160:ARG:HD3	3:A:715:HOH:O	1.98	0.63
1:A:268[A]:THR:HG23	3:A:739:HOH:O	1.98	0.63
1:B:282[B]:ARG:HH11	1:B:282[B]:ARG:HG2	1.64	0.63
1:B:286:ALA:HB2	1:B:300:TYR:CZ	2.34	0.63
1:B:57:LYS:N	1:B:57:LYS:HD2	2.14	0.62
1:B:450:ALA:HA	1:B:453[B]:MET:HE2	1.80	0.62
1:B:60:VAL:HG11	1:A:460:TRP:H	1.65	0.62
1:B:56:ASP:OD2	1:B:125:SER:OG	2.15	0.62
1:B:60:VAL:HG12	1:A:460:TRP:CE2	2.36	0.61
1:A:375:GLY:O	1:A:379[A]:ASN:ND2	2.34	0.59
1:B:39[A]:VAL:HG11	1:B:459:TRP:HB3	1.83	0.59
1:A:365:LYS:HE2	1:A:432:GLY:HA3	1.84	0.58
1:B:60:VAL:HG12	1:A:460:TRP:CD2	2.38	0.58
1:A:168:GLN:NE2	1:A:172[A]:LEU:HD11	2.19	0.58

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:192:ILE:HG22	1:B:192:ILE:O	2.04	0.58
1:B:195:GLU:HG3	1:B:235:TYR:CD1	2.39	0.57
1:B:282[B]:ARG:HG2	1:B:282[B]:ARG:NH1	2.20	0.56
1:B:78:LYS:CG	3:B:695:HOH:O	2.48	0.56
1:A:40[A]:VAL:HG22	1:A:453:MET:SD	2.46	0.56
1:A:36:ALA:O	1:A:40[B]:VAL:HG13	2.05	0.55
1:B:224:VAL:HG22	1:B:315:VAL:HB	1.89	0.55
1:A:224[A]:VAL:CG2	1:A:226[A]:ILE:HD11	2.38	0.54
1:A:358:GLU:OE1	1:A:436:SER:OG	2.21	0.54
1:A:409[A]:LYS:HE3	3:A:749:HOH:O	2.08	0.53
1:B:347[B]:MET:HE3	1:B:347[B]:MET:HA	1.91	0.53
1:B:310[B]:VAL:CG1	1:B:310[B]:VAL:O	2.57	0.53
1:B:114:LYS:NZ	1:B:154:ASP:OD2	2.28	0.52
1:B:126:ASP:OD1	1:B:164:TYR:OH	2.28	0.52
1:A:70:PHE:HB3	1:A:100:ILE:HD11	1.92	0.51
1:A:383:VAL:HG22	3:A:615:HOH:O	2.11	0.51
1:B:279:LYS:CE	1:B:279:LYS:HA	2.41	0.51
1:B:279:LYS:HA	1:B:279:LYS:HE2	1.93	0.51
1:A:100:ILE:HG22	1:A:449:VAL:HG21	1.93	0.51
1:A:25:THR:HG21	1:A:29:LYS:NZ	2.27	0.50
1:B:102:LYS:HE2	1:B:376:LYS:HA	1.92	0.50
1:B:317:LEU:HB3	3:B:605:HOH:O	2.10	0.50
1:A:224[A]:VAL:O	1:A:224[A]:VAL:HG13	2.12	0.50
1:B:195:GLU:HG3	1:B:235:TYR:CE1	2.47	0.50
1:A:202:VAL:HG11	1:A:239:LYS:HG3	1.94	0.50
1:A:25:THR:CG2	3:A:605:HOH:O	2.51	0.49
1:A:16:GLU:O	1:A:20[B]:THR:HG23	2.13	0.49
1:A:224[A]:VAL:HG22	1:A:226[A]:ILE:CD1	2.42	0.49
1:A:361:ARG:HA	1:A:369:ALA:HB1	1.96	0.48
1:A:271:GLU:OE2	1:A:297:LYS:NZ	2.42	0.48
1:A:224[A]:VAL:HG22	1:A:226[A]:ILE:HD11	1.95	0.48
1:B:248:VAL:HG21	1:B:310[A]:VAL:CG2	2.44	0.48
1:B:82:ARG:O	1:B:116:GLY:HA3	2.14	0.47
1:B:326:SER:OG	1:B:329:GLU:HG2	2.13	0.47
1:A:25:THR:CB	3:A:605:HOH:O	2.61	0.47
1:A:98:GLU:HG2	3:A:730:HOH:O	2.13	0.47
1:A:219:PHE:CD1	1:A:240[B]:VAL:HG12	2.49	0.47
1:A:25:THR:HG21	1:A:29:LYS:HZ1	1.79	0.47
1:A:163:GLY:HA2	1:A:187:TRP:CH2	2.51	0.46
1:A:379[A]:ASN:HA	3:A:665:HOH:O	2.15	0.46
1:A:146:ALA:HB2	1:A:177[A]:GLU:HG3	1.98	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:168:GLN:O	1:A:172[A]:LEU:HD13	2.15	0.46
1:B:286:ALA:O	1:B:289:VAL:HG22	2.15	0.46
1:B:454:LYS:HD2	1:B:459:TRP:CH2	2.51	0.46
1:A:366:GLY:N	1:A:432:GLY:O	2.48	0.46
1:B:201:VAL:HG11	3:B:605:HOH:O	2.15	0.46
1:B:20:THR:HG21	1:B:442:ASN:HB3	1.98	0.45
1:B:248:VAL:HG21	1:B:310[A]:VAL:HG23	1.99	0.45
1:A:194:PRO:HB3	1:A:231:ASN:OD1	2.17	0.45
1:B:241:ILE:HG13	3:B:620:HOH:O	2.17	0.44
1:A:267:PHE:HB3	1:A:272:ILE:HD11	1.99	0.44
1:B:122:LYS:HE2	3:B:723:HOH:O	2.16	0.44
1:A:267:PHE:HA	1:A:271:GLU:OE1	2.18	0.44
1:B:240:VAL:HG12	3:B:620:HOH:O	2.16	0.43
1:B:387:GLY:HA2	1:B:390:MET:HE2	1.99	0.43
1:B:271:GLU:CD	1:B:297:LYS:HZ3	2.21	0.43
1:A:130:ARG:O	1:A:134:VAL:HG23	2.18	0.43
1:A:230:GLY:O	1:A:234:GLN:HG3	2.18	0.43
2:B:501:ACT:H3	3:B:732:HOH:O	2.18	0.43
1:B:316:ALA:O	1:B:341:ILE:HA	2.18	0.43
1:A:237:ALA:O	1:A:240[B]:VAL:HG23	2.19	0.43
1:A:224[A]:VAL:HA	1:A:315:VAL:O	2.19	0.43
1:B:310[B]:VAL:CG1	1:B:313:VAL:HB	2.49	0.42
1:B:394:SER:O	3:B:602:HOH:O	2.21	0.42
1:B:105:LEU:HD22	1:B:105:LEU:O	2.20	0.42
1:A:211:HIS:CE1	1:A:420:LEU:HD21	2.54	0.42
1:B:325:VAL:HA	1:B:329:GLU:OE2	2.19	0.41
1:B:387:GLY:HA2	1:B:390:MET:CE	2.50	0.41
1:A:237:ALA:HA	1:A:240[B]:VAL:CG2	2.50	0.41
1:A:364:ASN:O	1:A:369:ALA:HB2	2.19	0.41
1:A:365:LYS:CE	1:A:432:GLY:HA3	2.48	0.41
1:B:48:GLN:NE2	1:B:89:LEU:HD22	2.35	0.41
1:A:272:ILE:HD13	1:A:272:ILE:N	2.35	0.41
1:A:387:GLY:HA2	1:A:390[A]:MET:CE	2.51	0.41
1:B:376:LYS:HB2	1:B:376:LYS:NZ	2.35	0.41
1:A:429:PRO:HB2	1:A:433:ILE:HB	2.02	0.41
1:A:78:LYS:HD2	1:A:150:VAL:O	2.21	0.41
1:B:437:LEU:HD23	1:B:437:LEU:HA	1.87	0.40
1:A:236:ALA:O	1:A:240[A]:VAL:HG23	2.21	0.40
1:A:334:ILE:HG21	1:A:360:HIS:CD2	2.56	0.40
1:B:257:LEU:HB2	1:B:285:ILE:HG21	2.03	0.40
1:B:310[B]:VAL:O	1:B:310[B]:VAL:HG13	2.21	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:192:ILE:O	1:B:192:ILE:CG2	2.69	0.40
1:B:163:GLY:HA2	1:B:187:TRP:CH2	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	479/460 (104%)	464 (97%)	14 (3%)	1 (0%)	44	38
1	B	476/460 (104%)	454 (95%)	19 (4%)	3 (1%)	22	13
All	All	955/920 (104%)	918 (96%)	33 (4%)	4 (0%)	37	23

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	192	ILE
1	B	192	ILE
1	B	346[A]	ASN
1	B	346[B]	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	371/351 (106%)	354 (95%)	17 (5%)	23	17
1	B	369/351 (105%)	356 (96%)	13 (4%)	31	25
All	All	740/702 (105%)	710 (96%)	30 (4%)	33	20

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	3	ASN
1	B	39[A]	VAL
1	B	39[B]	VAL
1	B	70	PHE
1	B	94	PHE
1	B	105	LEU
1	B	219	PHE
1	B	254	GLN
1	B	282[A]	ARG
1	B	282[B]	ARG
1	B	321[A]	THR
1	B	321[B]	THR
1	B	325	VAL
1	A	16	GLU
1	A	19	SER
1	A	20[A]	THR
1	A	20[B]	THR
1	A	25	THR
1	A	70	PHE
1	A	98	GLU
1	A	268[A]	THR
1	A	268[B]	THR
1	A	278	LEU
1	A	279	LYS
1	A	291	ASP
1	A	379[A]	ASN
1	A	379[B]	ASN
1	A	383	VAL
1	A	394[A]	SER
1	A	394[B]	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	28	GLN
1	A	23	ASN
1	A	61	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ACT	A	501	-	3,3,3	1.02	0	3,3,3	0.71	0
2	ACT	B	502	-	3,3,3	1.09	0	3,3,3	0.97	0
2	ACT	B	501	-	3,3,3	0.77	0	3,3,3	1.39	0
2	ACT	A	502	-	3,3,3	0.90	0	3,3,3	0.95	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	ACT	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	458/460 (99%)	0.21	15 (3%)	49 51	9, 25, 46, 70	23 (5%)
1	B	459/460 (99%)	0.41	25 (5%)	32 34	10, 28, 49, 83	19 (4%)
All	All	917/920 (99%)	0.31	40 (4%)	39 42	9, 27, 48, 83	42 (4%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	290	THR	5.1
1	B	289	VAL	4.7
1	B	288	LEU	3.8
1	A	288	LEU	3.8
1	B	280[A]	VAL	3.8
1	A	220	ALA	3.3
1	B	335	ALA	3.2
1	A	3	ASN	3.1
1	B	334	ILE	2.9
1	B	286	ALA	2.8
1	B	351[A]	GLN	2.8
1	B	225	ALA	2.8
1	B	285	ILE	2.8
1	B	345[A]	SER	2.8
1	A	231	ASN	2.7
1	B	254	GLN	2.5
1	A	254	GLN	2.5
1	B	216	ALA	2.4
1	B	299[A]	THR	2.4
1	A	289	VAL	2.4
1	B	2	SER	2.4
1	B	322	GLN	2.4
1	A	286	ALA	2.4
1	A	293	ALA	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	306	PRO	2.4
1	A	433	ILE	2.4
1	B	363	ALA	2.4
1	A	259	VAL	2.3
1	A	278	LEU	2.2
1	B	294	PHE	2.2
1	B	108	LEU	2.2
1	B	296	ASP	2.2
1	B	278	LEU	2.1
1	A	299	THR	2.1
1	B	352	ALA	2.1
1	A	40[A]	VAL	2.1
1	A	98	GLU	2.1
1	B	283[A]	LYS	2.1
1	A	290	THR	2.1
1	B	274	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ACT	B	502	4/4	0.75	0.18	38,38,39,46	0
2	ACT	A	502	4/4	0.75	0.18	53,56,57,60	0
2	ACT	B	501	4/4	0.81	0.20	32,39,44,52	0
2	ACT	A	501	4/4	0.90	0.14	46,46,48,56	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around ACT B 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around ACT A 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around ACT B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.