



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 14, 2025 – 02:35 PM JST

PDB ID : 8Z4C / pdb\_00008z4c  
Title : The Fab fragment of anti-Fibrin monoclonal antibody 59D8  
Authors : Zhang, J.; Sun, C.  
Deposited on : 2024-04-17  
Resolution : 2.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3436 (2.08-2.04)
Clashscore	180529	3661 (2.08-2.04)
Ramachandran outliers	177936	3649 (2.08-2.04)
Sidechain outliers	177891	3649 (2.08-2.04)
RSRZ outliers	164620	3436 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	222	<div> <div>7%</div> <div>82%</div> <div>14%</div> <div>...</div> </div>
1	H	222	<div> <div>3%</div> <div>86%</div> <div>10%</div> <div>.</div> </div>
2	D	220	<div> <div>92%</div> <div>8%</div> </div>
2	L	220	<div> <div>14%</div> <div>77%</div> <div>20%</div> <div>.</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6966 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab 59D8 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	216	Total	C	N	O	S	0	0	0
			1636	1021	283	324	8			
1	A	218	Total	C	N	O	S	0	0	0
			1650	1028	286	328	8			

- Molecule 2 is a protein called Fab 59D8 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	220	Total	C	N	O	S	0	0	0
			1705	1074	285	341	5			
2	D	220	Total	C	N	O	S	0	0	0
			1705	1074	285	341	5			



- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	H	92	Total	O	0	0
			92	92		
3	L	64	Total	O	0	0
			64	64		
3	A	51	Total	O	0	0
			51	51		
3	D	63	Total	O	0	0
			63	63		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Fab 59D8 heavy chain



- Molecule 1: Fab 59D8 heavy chain



- Molecule 2: Fab 59D8 light chain



- Molecule 2: Fab 59D8 light chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.16Å 65.29Å 100.14Å 90.00° 97.90° 90.00°	Depositor
Resolution (Å)	43.50 – 2.06 43.50 – 2.06	Depositor EDS
% Data completeness (in resolution range)	99.9 (43.50-2.06) 99.9 (43.50-2.06)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.74 (at 2.07Å)	Xtriage
Refinement program	PHENIX (1.15.2_3472: ???)	Depositor
R, $R_{free}$	0.198 , 0.234 0.202 , 0.237	Depositor DCC
$R_{free}$ test set	2775 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.2	Xtriage
Anisotropy	0.273	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 45.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6966	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.33	0/1691	0.59	1/2298 (0.0%)
1	H	0.30	0/1677	0.56	1/2278 (0.0%)
2	D	0.30	0/1742	0.53	0/2363
2	L	0.34	0/1742	0.60	2/2363 (0.1%)
All	All	0.32	0/6852	0.57	4/9302 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	31	TYR	CB-CA-C	-5.94	98.52	110.40
2	L	36	THR	CB-CA-C	-5.91	95.63	111.60
1	A	181	LEU	CA-CB-CG	5.73	128.49	115.30
1	H	181	LEU	CA-CB-CG	5.06	126.94	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	132	ARG	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1650	0	1591	24	0
1	H	1636	0	1572	18	0
2	D	1705	0	1674	11	0
2	L	1705	0	1674	49	0
3	A	51	0	0	4	0
3	D	63	0	0	3	0
3	H	92	0	0	3	1
3	L	64	0	0	6	1
All	All	6966	0	6511	96	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:36:THR:CG2	2:L:56:VAL:HG23	2.06	0.86
2:L:36:THR:HG23	2:L:56:VAL:HG23	1.62	0.81
2:L:148:ARG:NH2	3:L:301:HOH:O	2.15	0.80
1:A:166:VAL:HG22	1:A:185:VAL:HG12	1.68	0.75
1:H:120:LYS:NZ	3:H:302:HOH:O	2.20	0.74
2:L:36:THR:HG22	2:L:38:LEU:H	1.53	0.74
1:A:101:GLY:O	3:A:301:HOH:O	2.06	0.73
2:D:41:LEU:HG	2:D:94:TRP:CZ3	2.23	0.73
1:H:216:HIS:O	3:H:301:HOH:O	2.06	0.72
2:L:41:LEU:HG	2:L:94:TRP:CZ3	2.25	0.71
2:L:2:ILE:HG21	2:L:29:LEU:HD21	1.71	0.70
2:L:11:LEU:HD21	2:L:19:ALA:HB1	1.76	0.68
2:L:39:ASN:ND2	2:L:54:TYR:O	2.27	0.68
2:L:36:THR:HG21	2:L:56:VAL:HG23	1.77	0.65
2:L:88:LEU:HD21	2:L:174:SER:HA	1.79	0.64
2:L:35:THR:HB	2:L:37:TYR:CE2	2.32	0.64
1:A:120:LYS:NZ	3:A:302:HOH:O	2.32	0.62
2:L:36:THR:HG21	2:L:56:VAL:N	2.18	0.59
1:A:98:ARG:O	1:A:103:TYR:HA	2.03	0.57

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:131:SER:C	1:A:133:SER:N	2.55	0.57
1:A:83:MET:HE2	1:A:86:LEU:HD21	1.88	0.56
1:A:40:THR:OG1	1:A:42:ASP:OD1	2.14	0.56
2:D:38:LEU:HD22	2:D:76:PHE:CG	2.41	0.56
1:H:130:CYS:SG	2:L:125:PRO:HG3	2.46	0.56
2:D:114:ARG:NH2	3:D:307:HOH:O	2.40	0.55
1:H:106:TRP:CZ2	2:L:41:LEU:HD11	2.42	0.54
2:L:72:SER:O	3:L:302:HOH:O	2.19	0.54
2:D:11:LEU:HD21	2:D:19:ALA:HB1	1.89	0.54
2:D:50:LYS:NZ	3:D:306:HOH:O	2.39	0.54
1:H:108:GLN:HA	2:L:48:SER:HG	1.72	0.54
1:A:196:THR:HB	1:A:213:ARG:CZ	2.38	0.54
1:H:166:VAL:HG22	1:H:185:VAL:HG12	1.90	0.53
1:A:32:TYR:O	1:A:72:ARG:NH2	2.35	0.53
1:H:83:MET:HE2	1:H:86:LEU:HD21	1.91	0.52
2:L:38:LEU:HD22	2:L:76:PHE:CG	2.45	0.51
1:A:134:THR:O	1:A:135:SER:HB2	2.11	0.51
1:A:216:HIS:CD2	1:A:216:HIS:N	2.79	0.51
1:A:191:SER:O	1:A:195:LYS:HG2	2.11	0.50
2:L:36:THR:HG22	2:L:38:LEU:N	2.23	0.50
2:L:170:THR:O	3:L:301:HOH:O	2.20	0.50
2:L:32:SER:C	2:L:34:GLY:H	2.14	0.50
2:L:11:LEU:HD23	2:L:109:LEU:HD12	1.95	0.49
1:H:60:TYR:HB2	1:H:65:LYS:HD3	1.94	0.48
1:A:106:TRP:CZ2	2:D:41:LEU:HD11	2.49	0.47
1:H:83:MET:HB3	1:H:86:LEU:HD21	1.96	0.47
1:A:197:TYR:O	1:A:213:ARG:HD3	2.15	0.47
2:L:2:ILE:HG21	2:L:29:LEU:CD2	2.43	0.47
2:L:13:VAL:HG21	2:L:19:ALA:HB2	1.96	0.47
2:L:131:LEU:O	2:L:189:LYS:HD3	2.14	0.47
2:L:51:ARG:HD2	2:L:54:TYR:HB3	1.97	0.47
1:A:212:LYS:HE2	1:A:212:LYS:HB2	1.66	0.47
2:L:45:PRO:HB3	2:L:171:GLU:HG3	1.96	0.46
2:L:32:SER:C	2:L:34:GLY:N	2.68	0.46
2:L:36:THR:HG21	2:L:56:VAL:H	1.79	0.46
1:H:11:LEU:HD21	1:H:117:ALA:O	2.17	0.45
1:H:29:PHE:CD2	1:H:77:ASN:HA	2.51	0.45
2:L:39:ASN:HB2	2:L:94:TRP:CZ3	2.51	0.45
2:L:37:TYR:H	2:L:55:LEU:HD22	1.81	0.45
2:L:37:TYR:HB2	2:L:97:THR:OG1	2.16	0.45
2:L:53:ILE:HG12	2:L:59:VAL:HA	1.99	0.45

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9:GLY:HA3	1:A:18:LEU:HD11	1.98	0.45
1:A:141:LEU:HD21	1:A:216:HIS:HA	1.99	0.45
2:L:53:ILE:HG22	2:L:55:LEU:O	2.17	0.44
2:L:66:ARG:NH1	2:L:87:ASP:OD2	2.44	0.44
2:L:207:LEU:HD13	2:L:211:VAL:HG22	1.98	0.44
1:A:137:SER:HB3	1:A:138:THR:H	1.46	0.44
2:L:36:THR:HA	3:L:304:HOH:O	2.18	0.43
2:D:113:ARG:HG3	2:D:177:SER:HB2	1.99	0.43
1:A:30:SER:HB3	3:A:308:HOH:O	2.17	0.43
2:L:197:VAL:HG22	2:L:216:ASN:OD1	2.19	0.43
2:D:66:ARG:NH1	2:D:87:ASP:OD2	2.42	0.43
1:A:1:GLN:HG3	3:A:348:HOH:O	2.18	0.43
1:H:146:LYS:NZ	1:H:174:GLN:OE1	2.52	0.42
1:H:1:GLN:OE1	1:H:1:GLN:N	2.35	0.42
1:H:216:HIS:HD2	3:H:362:HOH:O	2.03	0.42
2:L:2:ILE:HD13	2:L:29:LEU:CD2	2.48	0.42
2:L:54:TYR:CD2	2:L:55:LEU:HG	2.55	0.41
1:H:192:LEU:HD22	1:H:216:HIS:ND1	2.34	0.41
2:L:175:LYS:NZ	2:L:176:ASP:OD2	2.53	0.41
1:A:1:GLN:OE1	1:A:1:GLN:N	2.45	0.41
2:L:32:SER:OG	2:L:33:ASP:N	2.54	0.41
2:L:55:LEU:HB2	2:L:58:LYS:HG2	2.01	0.41
1:H:186:THR:HG21	2:L:143:ASN:ND2	2.35	0.41
2:L:167:GLU:HG3	3:L:331:HOH:O	2.20	0.41
1:A:131:SER:C	1:A:133:SER:H	2.22	0.41
1:A:132:ARG:O	1:A:133:SER:HB2	2.20	0.41
2:D:3:GLU:HB2	2:D:26:SER:HB3	2.02	0.41
2:L:14:ILE:HG12	2:L:112:LYS:HB3	2.03	0.40
1:A:29:PHE:O	1:A:72:ARG:NH2	2.54	0.40
2:D:186:THR:HG23	3:D:324:HOH:O	2.21	0.40
1:H:56:ARG:HH11	1:H:56:ARG:HG3	1.86	0.40
1:H:108:GLN:HA	2:L:48:SER:OG	2.20	0.40
2:L:88:LEU:HD13	3:L:340:HOH:O	2.21	0.40
2:D:45:PRO:HB3	2:D:171:GLU:HG3	2.02	0.40
2:L:39:ASN:HB2	2:L:94:TRP:CE3	2.57	0.40
2:L:42:LEU:HB2	2:L:52:LEU:HD11	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:H:387:HOH:O	3:L:345:HOH:O[2_545]	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	216/222 (97%)	207 (96%)	8 (4%)	1 (0%)	25	17
1	H	212/222 (96%)	205 (97%)	6 (3%)	1 (0%)	25	17
2	D	218/220 (99%)	213 (98%)	5 (2%)	0	100	100
2	L	218/220 (99%)	212 (97%)	6 (3%)	0	100	100
All	All	864/884 (98%)	837 (97%)	25 (3%)	2 (0%)	44	38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	103	TYR
1	A	135	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/192 (98%)	182 (97%)	6 (3%)	34	28
1	H	186/192 (97%)	184 (99%)	2 (1%)	70	70
2	D	196/196 (100%)	194 (99%)	2 (1%)	73	73

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	L	196/196 (100%)	191 (97%)	5 (3%)	41	37
All	All	766/776 (99%)	751 (98%)	15 (2%)	50	47

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	96	CYS
1	H	181	LEU
2	L	31	TYR
2	L	32	SER
2	L	35	THR
2	L	39	ASN
2	L	148	ARG
1	A	130	CYS
1	A	132	ARG
1	A	135	SER
1	A	136	GLU
1	A	137	SER
1	A	181	LEU
2	D	58	LYS
2	D	114	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	L	39	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	218/222 (98%)	0.45	16 (7%) 22 24	30, 45, 98, 127	0
1	H	216/222 (97%)	0.02	6 (2%) 55 56	24, 35, 71, 101	0
2	D	220/220 (100%)	0.30	1 (0%) 87 89	27, 48, 69, 90	0
2	L	220/220 (100%)	0.71	30 (13%) 8 8	26, 46, 93, 113	0
All	All	874/884 (98%)	0.37	53 (6%) 28 29	24, 43, 86, 127	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	35	THR	7.5
2	L	34	GLY	7.5
1	A	134	THR	6.9
2	L	55	LEU	6.4
2	L	36	THR	4.8
2	L	220	CYS	4.6
2	L	31	TYR	4.2
1	A	188	PRO	4.0
2	L	56	VAL	4.0
2	L	37	TYR	3.9
2	L	15	ILE	3.8
2	L	30	LEU	3.8
1	H	135	SER	3.8
2	L	32	SER	3.8
1	A	192	LEU	3.7
2	L	63	VAL	3.4
2	L	72	SER	3.4
1	A	141	LEU	3.3
2	L	29	LEU	3.3
1	H	194	THR	3.3
1	A	189	SER	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	L	64	PRO	3.1
1	A	217	HIS	3.0
1	H	217	HIS	2.9
1	A	193	GLY	2.9
1	A	187	VAL	2.9
2	L	54	TYR	2.9
2	D	220	CYS	2.9
1	A	196	THR	2.7
1	A	102	ASP	2.7
2	L	73	GLY	2.6
2	L	33	ASP	2.5
2	L	11	LEU	2.5
2	L	71	GLY	2.5
2	L	97	THR	2.5
2	L	76	PHE	2.4
2	L	62	GLY	2.4
1	H	216	HIS	2.3
1	A	103	TYR	2.3
1	H	130	CYS	2.3
2	L	28	SER	2.3
1	A	130	CYS	2.2
1	A	133	SER	2.2
1	A	137	SER	2.2
2	L	88	LEU	2.2
2	L	12	SER	2.2
1	A	213	ARG	2.1
2	L	52	LEU	2.1
2	L	14	ILE	2.1
1	H	218	HIS	2.1
1	A	131	SER	2.1
2	L	27	GLN	2.0
2	L	75	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.