



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 13, 2024 – 03:09 AM EDT

PDB ID : 3ZUQ
Title : Crystal structure of an engineered botulinum neurotoxin type B - derivative, LC-B-GS-Hn-B
Authors : Masuyer, G.; Stancombe, P.; Chaddock, J.A.; Acharya, K.R.
Deposited on : 2011-07-19
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	906	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6953 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BOTULINUM NEUROTOXIN TYPE B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	850	Total	C	N	O	S	0	0	0
			6908	4450	1102	1331	25			

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	894	LEU	-	expression tag	UNP P10844
A	895	GLU	-	expression tag	UNP P10844
A	896	ALA	-	expression tag	UNP P10844
A	897	LEU	-	expression tag	UNP P10844
A	898	ALA	-	expression tag	UNP P10844
A	899	SER	-	expression tag	UNP P10844
A	900	GLY	-	expression tag	UNP P10844
A	901	HIS	-	expression tag	UNP P10844
A	902	HIS	-	expression tag	UNP P10844
A	903	HIS	-	expression tag	UNP P10844
A	904	HIS	-	expression tag	UNP P10844
A	905	HIS	-	expression tag	UNP P10844
A	906	HIS	-	expression tag	UNP P10844

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	44	Total	O	0	0
			44	44		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	89.37Å 103.79Å 114.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	77.03 – 2.70 43.82 – 2.70	Depositor EDS
% Data completeness (in resolution range)	93.1 (77.03-2.70) 93.1 (43.82-2.70)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.83 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.248 , 0.281 0.240 , 0.272	Depositor DCC
R_{free} test set	1417 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	43.0	Xtriage
Anisotropy	0.191	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 33.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6953	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/7049	0.45	0/9530

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6908	0	6848	36	0
2	A	1	0	0	0	0
3	A	44	0	0	0	0
All	All	6953	0	6848	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:368:LYS:H	1:A:412:GLN:HE22	1.44	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:261:THR:HG22	1:A:262:ASP:O	1.98	0.64
1:A:437:CYS:HB3	1:A:479:LEU:HD21	1.79	0.63
1:A:492:ILE:HG23	1:A:742:THR:HG23	1.84	0.60
1:A:102:LYS:HE3	1:A:365:TYR:HA	1.85	0.57
1:A:572:ILE:HG22	1:A:663:LEU:HB3	1.85	0.57
1:A:693:LEU:HD11	1:A:822:MET:HG2	1.88	0.55
1:A:673:ASN:HD22	1:A:676:ASN:H	1.54	0.55
1:A:626:ALA:HA	1:A:786:ILE:HD11	1.89	0.53
1:A:653:SER:HB3	1:A:808:ASN:HD21	1.75	0.52
1:A:75:ASP:HB3	1:A:167:LEU:HD11	1.95	0.49
1:A:200:VAL:HG11	1:A:224:PRO:HG3	1.96	0.48
1:A:81:THR:HG22	1:A:83:ASP:H	1.78	0.48
1:A:399:ASN:HD21	1:A:412:GLN:HE21	1.61	0.48
1:A:312:ASP:HB3	1:A:315:ILE:HG12	1.95	0.47
1:A:399:ASN:HB3	1:A:406:GLU:HA	1.95	0.47
1:A:273:GLY:HA3	1:A:357:THR:CG2	2.46	0.46
1:A:572:ILE:HG13	1:A:573:PHE:N	2.31	0.45
1:A:786:ILE:HG22	1:A:787:ASP:N	2.31	0.45
1:A:666:GLY:HA2	1:A:688:GLU:HA	1.99	0.44
1:A:20:ILE:HG12	1:A:44:TRP:CZ3	2.53	0.44
1:A:38:LYS:HG2	1:A:40:THR:O	2.18	0.44
1:A:691:PRO:HB3	1:A:742:THR:CG2	2.48	0.44
1:A:10:TYR:H	1:A:86:ASN:HD22	1.66	0.43
1:A:357:THR:HG22	1:A:360:ASN:H	1.82	0.43
1:A:154:ILE:HA	1:A:541:GLU:O	2.18	0.43
1:A:693:LEU:HD21	1:A:822:MET:HE3	2.00	0.43
1:A:297:ARG:HG2	1:A:340:ILE:HD12	2.01	0.43
1:A:273:GLY:HA3	1:A:357:THR:HG21	2.01	0.42
1:A:207:GLN:H	1:A:207:GLN:HE21	1.66	0.41
1:A:10:TYR:H	1:A:86:ASN:ND2	2.18	0.41
1:A:206:VAL:HG21	1:A:752:GLU:OE1	2.21	0.41
1:A:779:LYS:H	1:A:779:LYS:HG3	1.68	0.41
1:A:213:SER:C	1:A:215:PHE:N	2.74	0.41
1:A:728:TRP:CE3	1:A:830:LEU:HD13	2.56	0.40
1:A:118:TYR:HA	1:A:324:PHE:CZ	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	844/906 (93%)	811 (96%)	29 (3%)	4 (0%)	29	54

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	146	GLY
1	A	213	SER
1	A	210	LYS
1	A	214	ILE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	776/801 (97%)	768 (99%)	8 (1%)	76	91

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	207	GLN
1	A	216	ASN
1	A	515	ASN
1	A	564	LYS
1	A	566	PHE
1	A	796	ASN
1	A	808	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	885	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	86	ASN
1	A	207	GLN
1	A	275	GLN
1	A	412	GLN
1	A	435	GLN
1	A	519	ASN
1	A	559	GLN
1	A	641	ASN
1	A	673	ASN
1	A	676	ASN
1	A	719	ASN
1	A	808	ASN
1	A	864	ASN
1	A	889	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	850/906 (93%)	0.24	43 (5%) 28 26	25, 42, 66, 80	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	146	GLY	11.0
1	A	147	GLU	5.0
1	A	149	GLU	4.7
1	A	145	PRO	4.4
1	A	892	ASN	4.3
1	A	211	GLY	4.3
1	A	785	ASN	4.2
1	A	307	LEU	4.2
1	A	314	ASN	4.1
1	A	566	PHE	3.5
1	A	310	ILE	3.5
1	A	652	ILE	3.3
1	A	150	ARG	3.2
1	A	651	ASP	3.2
1	A	653	SER	3.1
1	A	784	ILE	3.1
1	A	891	TYR	3.0
1	A	309	CYS	2.9
1	A	212	ALA	2.9
1	A	707	ILE	2.9
1	A	315	ILE	2.8
1	A	143	SER	2.8
1	A	516	TYR	2.7
1	A	517	ILE	2.5
1	A	144	ASN	2.5
1	A	308	VAL	2.5
1	A	783	ASN	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	439	ASP	2.4
1	A	708	ASP	2.4
1	A	786	ILE	2.4
1	A	330	PHE	2.3
1	A	320	TYR	2.2
1	A	148	VAL	2.2
1	A	311	SER	2.2
1	A	586	ARG	2.2
1	A	334	SER	2.2
1	A	338	TYR	2.1
1	A	141	LEU	2.1
1	A	341	ASP	2.1
1	A	118	TYR	2.0
1	A	335	GLU	2.0
1	A	882	THR	2.0
1	A	1	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ZN	A	1440	1/1	0.99	0.02	32,32,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.